

PSETs Landing Page*

Anish Krishna Lakkapragada

This is the documentation for using my PSET PDFs responsibly. I post these LaTeX'd PSETs (1) as an education resource for friends at other universities, fellow Yalies, and all those interested and (2) for quick reference. These PSETs are not to be used irresponsibly; only look at the solution after giving each problem an honest attempt. **If YOU USE THESE PSETS TO CHEAT, YOU ARE NOT ONLY STUPID BUT YOU ARE CHEATING YOURSELF OUT OF THE ABILITY TO FALL IN LOVE WITH MATH.** Furthermore, I am not smarter than you and my solutions did not always get a perfect score.

The general format for accessing the (one-indexed) `N`th assigned PSET PDF of a Yale course with course number `CODE` is:

`https://anish.lakkapragada.com/notes/TYPE-CODE/psets/N.pdf`

where `TYPE` is `stats` or `math`. Similarly, to access my solution for this PSET you can go to:

`https://anish.lakkapragada.com/notes/TYPE-CODE/sols/N.pdf`

These PSETs and associated solution PDFs are synchronized daily at 4:20AM with my computer files through a Cronjob Shell Script. If you want to contribute any corrections, please email `anish.lakkapragada@yale.edu`.

*Note that PDF here is referring to Portable Document Format, not to be confused with the veritable Probability Density Function.

Question 1

$$(a) \quad T_1 = \frac{(40 + 30 + T_3 + T_2)}{4} \rightarrow 4T_1 - T_3 - T_2 = 70$$

$$T_2 = (10 + 30 + T_4 + T_1) \rightarrow 4T_2 - T_4 - T_1 = 140$$

$$T_3 = \left(\frac{20 + 40 + T_4 + T_1}{4} \right) \rightarrow 4T_3 - T_4 - T_1 = 60$$

$$T_4 = \frac{(20 + 10 + T_2 + T_3)}{4} \rightarrow 4T_4 - T_3 - T_2 = 30$$

(b) $A = \begin{bmatrix} 4 & -1 & -1 & 0 \\ -1 & 4 & 0 & -1 \\ -1 & 0 & 4 & -1 \\ 0 & -1 & -1 & 4 \end{bmatrix}$

Applying G.E.

$$\begin{aligned} R_1 &= \frac{R_1}{4} \rightarrow \begin{bmatrix} 1 & -\frac{1}{4} & -\frac{1}{4} & 0 & | & \frac{1}{4} & 0 & 0 & 0 \\ -1 & 4 & 0 & -1 & | & 0 & 1 & 0 & 0 \\ -1 & 0 & 4 & -1 & | & 0 & 0 & 0 & 0 \\ 0 & -1 & -1 & 4 & | & 0 & 0 & 0 & 1 \end{bmatrix} \quad \begin{aligned} R_2 &= R_2 + R_1 \\ R_3 &= R_3 + R_1 \end{aligned} \rightarrow \begin{bmatrix} 1 & -\frac{1}{4} & -\frac{1}{4} & 0 & | & \frac{1}{4} & 0 & 0 & 0 \\ 0 & \frac{15}{4} & -\frac{1}{4} & -1 & | & \frac{1}{4} & 1 & 0 & 0 \\ 0 & -\frac{1}{4} & \frac{15}{4} & -1 & | & \frac{1}{4} & 0 & 1 & 0 \\ 0 & -1 & -1 & 4 & | & 0 & 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & -\frac{4}{15} & -\frac{1}{15} & \frac{4}{15} & \frac{1}{15} & 0 & 0 \\ 0 & 1 & -\frac{1}{15} & -\frac{4}{15} & \frac{1}{15} & \frac{4}{15} & 0 & 0 \\ 0 & 0 & \frac{5}{15} & \frac{1}{15} & \frac{15}{15} & \frac{15}{15} & 0 & 0 \\ 0 & 0 & -\frac{16}{15} & \frac{54}{15} & \frac{1}{5} & \frac{4}{15} & 0 & 1 \end{array} \right] \begin{array}{l} R_1 + \frac{R_2}{4} \rightarrow R_1 \\ R_3 + \frac{R_2}{4} \rightarrow R_3 \\ R_4 + R_2 \rightarrow R_4 \end{array}$$

$$\begin{aligned} & \left(\frac{15}{5} R_3 \Rightarrow R_3 \right. \\ & \left. \begin{array}{ccc|ccc} 1 & 0 & -\frac{4}{15} & \frac{4}{15} & -\frac{1}{15} & 0 & 0 \\ 0 & 1 & -\frac{1}{15} & \frac{1}{15} & \frac{4}{15} & 0 & 0 \\ 0 & 0 & 1 & -\frac{1}{15} & -\frac{2}{15} & \frac{15}{56} & 0 \\ 0 & 0 & -\frac{16}{15} & \frac{56}{15} & \frac{56}{15} & 0 & 1 \end{array} \right) \xrightarrow{\begin{array}{l} R_2 + \frac{R_3}{15} \rightarrow R_2 \\ R_1 + \frac{4R_3}{15} \rightarrow R_1 \\ R_4 + \frac{16}{5} R_3 \rightarrow R_4 \end{array}} \rightarrow \begin{array}{ccc|ccc} 1 & 0 & -\frac{1}{4} & \frac{2}{4} & \frac{1}{4} & \frac{1}{4} & 0 \\ 0 & 1 & -\frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & 0 \\ 0 & 0 & 1 & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & 0 \\ 0 & 0 & \frac{24}{4} & \frac{1}{4} & \frac{2}{4} & \frac{1}{4} & 1 \end{array}$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{7}{24} & \frac{1}{12} & \frac{1}{12} & \frac{1}{24} \\ \frac{1}{12} & \frac{7}{24} & \frac{1}{24} & \frac{1}{12} \\ \frac{1}{12} & \frac{1}{24} & \frac{7}{24} & \frac{1}{12} \\ \frac{1}{24} & \frac{1}{12} & \frac{1}{12} & \frac{7}{24} \end{bmatrix} \begin{matrix} \leftarrow R_1 + R_4 \rightarrow R_1 \\ \leftarrow R_2 + 2R_4 \end{matrix}$$

$$V(A^{-1})$$

$$R_3 + 2R_3 \xrightarrow{7} R_3$$

Question 1 continued

$$(c) \quad AT = X$$

$$T = A^{-1}X$$

$$T = \begin{bmatrix} \frac{7}{24} & \frac{1}{12} & \frac{1}{12} & \frac{1}{24} \\ \frac{1}{12} & \frac{7}{24} & \frac{1}{24} & \frac{1}{12} \\ \frac{1}{12} & \frac{1}{24} & \frac{7}{24} & \frac{1}{12} \\ \frac{1}{24} & \frac{1}{12} & \frac{1}{12} & \frac{7}{24} \end{bmatrix} \begin{bmatrix} 30 \\ 40 \\ 60 \\ 30 \end{bmatrix} = \begin{bmatrix} 30 \\ 22.5 \\ 27.5 \\ 20 \end{bmatrix}$$

Question 2

To find the basis of $N(L_A)$, we solve $AX=0$

$$\left(\begin{array}{cccc|c} 3 & 5 & -4 & 2 & 0 \\ 2 & 4 & -6 & 3 & 0 \\ 11 & 17 & -8 & 9 & 0 \end{array}\right)$$

$$\begin{array}{l} 3R_2 - 2R_1 \rightarrow R_2 \\ 3R_3 - 11R_1 \rightarrow R_3 \end{array}$$

$$\left(\begin{array}{cccc|c} 3 & 5 & -4 & 2 & 0 \\ 0 & 2 & -10 & 5 & 0 \\ 0 & -4 & 20 & -10 & 0 \end{array}\right)$$

$$\begin{array}{l} 2R_2 + R_3 \rightarrow R_3 \\ 2R_1 - 5R_2 \rightarrow R_1 \end{array}$$

$$\left(\begin{array}{cccc|c} 6 & 0 & 42 & -21 & 0 \\ 0 & 2 & -10 & 5 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array}\right)$$

$$x_1 + 7x_3 - 3.5x_4 = 0$$

$$2x_2 - 10x_3 + 5x_4 = 0$$

$$x_1 = 3.5x_4 - 7x_3$$

$$x_2 = 5x_3 - 2.5x_4$$

$$\begin{array}{c} x_1 \ x_2 \ x_3 \ x_4 \\ \left(\begin{array}{cccc|c} 1 & 0 & 7 & -3.5 & 0 \\ 0 & 2 & -10 & 5 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array}\right) \end{array}$$

$$\frac{R_1}{6} \rightarrow R_1$$

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 3.5t_4 - 7t_3 \\ 5t_3 - 2.5t_4 \\ t_3 \\ t_4 \end{pmatrix} = t_3 \begin{pmatrix} -7 \\ 5 \\ 1 \\ 0 \end{pmatrix} + t_4 \begin{pmatrix} 3.5 \\ -2.5 \\ 0 \\ 1 \end{pmatrix}$$

Thus, the basis for $N(L_A)$ can be given by $\left\{ \begin{pmatrix} -7 \\ 5 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 3.5 \\ -2.5 \\ 0 \\ 1 \end{pmatrix} \right\}$.

The set of solutions to the given system is given by the span of this basis.

Question 3

$$AX=b$$

$$A^T X = b$$

$$[A|b] = \left(\begin{array}{cccc|c} 1 & -3 & -3 & -14 & 8 \\ 2 & -6 & -3 & -1 & -5 \\ 3 & -9 & -5 & -6 & -4 \end{array} \right) \xrightarrow{\substack{R_2 - 2R_1 \rightarrow R_2 \\ R_3 - 3R_1 \rightarrow R_3}} \left(\begin{array}{cccc|c} 1 & -3 & -3 & -14 & 8 \\ 0 & 0 & 3 & 27 & -21 \\ 0 & 0 & 4 & 36 & -28 \end{array} \right)$$

$$\left(\begin{array}{cccc|c} 1 & -3 & 0 & 13 & -13 \\ 0 & 0 & 1 & 9 & -7 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right) \xleftarrow{\substack{R_2 \rightarrow R_2 \\ \frac{R_2}{3} \rightarrow R_2}} \left(\begin{array}{cccc|c} 1 & -3 & 0 & 13 & -13 \\ 0 & 0 & 3 & 27 & -21 \\ 0 & 0 & 4 & 36 & -28 \end{array} \right) \xleftarrow{R_1 + R_2 \rightarrow R_1}$$

$$3R_3 - 4R_2 \rightarrow R_3$$

$$\begin{cases} x_1 - 3x_2 + 13x_4 = -13 \Rightarrow x_1 = -13 + 3x_2 - 13x_4 = -13 + 3t_2 - 13t_4 \\ x_3 + 9x_4 = -7 \Rightarrow x_3 = -7 - 9t_4 \end{cases}$$

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} -13 + 3t_2 - 13t_4 \\ t_2 \\ -7 - 9t_4 \\ t_4 \end{pmatrix} = \begin{pmatrix} -13 \\ 0 \\ -7 \\ 0 \end{pmatrix} + t_2 \begin{pmatrix} 3 \\ 1 \\ 0 \\ 0 \end{pmatrix} + t_4 \begin{pmatrix} -13 \\ 0 \\ -9 \\ 1 \end{pmatrix}$$

Vectors in \mathbb{R}^4 whose image is equal to $b \in \mathbb{R}^3$ under linear map given in A.

The basis of $N(A)$ can be given by $\left\{ \begin{pmatrix} 3 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} -13 \\ 0 \\ 9 \\ 1 \end{pmatrix} \right\}$.

Question 4

(i)

$$[T]^\beta = \begin{bmatrix} 1 & -2 & 4 & -8 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & 1 & 1 \\ 1 & 2 & 4 & 8 \end{bmatrix}$$

$$T(1) = (1, 1, 1, 1)$$

$$T(x) = (-2, -1, 1, 2)$$

$$T(x^2) = (4, 1, 1, 4)$$

$$T(x^3) = (-8, -1, 1, 8)$$

(ii)

$$Tx = \begin{bmatrix} 1 \\ 3 \\ 3 \\ 33 \end{bmatrix}$$

$$\left[T \mid \begin{bmatrix} 1 \\ 3 \\ 3 \\ 33 \end{bmatrix} \right] \rightarrow \left[\begin{array}{cccc|c} 1 & -2 & 4 & -8 & 1 \\ 1 & -1 & 1 & -1 & 3 \\ 1 & 1 & 1 & 1 & 3 \\ 1 & 2 & 4 & 8 & 33 \end{array} \right] \xrightarrow{\substack{R_2 - R_1 \rightarrow R_2 \\ R_3 - R_1 \rightarrow R_3 \\ R_4 - R_1 \rightarrow R_4}} \left[\begin{array}{cccc|c} 1 & -2 & 4 & -8 & 1 \\ 0 & 1 & -3 & 7 & 2 \\ 0 & 3 & -3 & 9 & 12 \\ 0 & 4 & 0 & 16 & 32 \end{array} \right]$$

$$\left[\begin{array}{cccc|c} 3 & 0 & 0 & 6 & 21 \\ 0 & 2 & 0 & 2 & 10 \\ 0 & 0 & 6 & -12 & 6 \\ 0 & 0 & 0 & 12 & 12 \end{array} \right]$$

$$\begin{aligned} R_4 - 2R_3 &\rightarrow R_4 \\ 2R_2 + R_3 &\rightarrow R_2 \\ 3R_1 + R_3 &\rightarrow R_1 \end{aligned}$$

$$\left[\begin{array}{cccc|c} 1 & 0 & -2 & 6 & 5 \\ 0 & 1 & -3 & 7 & 2 \\ 0 & 0 & 6 & -12 & 18 \\ 0 & 0 & 12 & -12 & 24 \end{array} \right]$$

$$\begin{aligned} R_3 - 3R_2 &\rightarrow R_3 \\ R_4 - 4R_2 &\rightarrow R_4 \\ 2R_2 + R_1 &\rightarrow R_1 \end{aligned}$$

$$\left[\begin{array}{cccc|c} 6 & 0 & 0 & 0 & 30 \\ 0 & 12 & 0 & 0 & 48 \\ 0 & 0 & 6 & 6 & 18 \\ 0 & 0 & 0 & 12 & 12 \end{array} \right] \xrightarrow{\substack{R_1 \rightarrow R_1 \\ R_2 \rightarrow R_2 \\ R_3 \rightarrow R_3 \\ R_4 \rightarrow R_4}} \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 5 \\ 0 & 1 & 0 & 0 & 4 \\ 0 & 0 & 1 & 0 & 3 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right]$$

Thus, we get this polynomial as:

$$f(x) = 5 + 4x + 3x^2 + x^3$$

Question 5

$$V = \{ (x_1, x_2, x_3, x_4, x_5) : x_1 - 2x_2 + 3x_3 - x_4 + 2x_5 = 0 \}$$

is finally a subspace of \mathbb{R}^5 .

We first find the basis of V :

$$x_1 = 2x_2 - 3x_3 + x_4 - 2x_5$$

$$\begin{aligned} (x_1, x_2, x_3, x_4, x_5) &= (2t_2 - 3t_3 + t_4 - 2t_5, t_2, t_3, t_4, t_5) \\ &= t_2(2, 1, 0, 0, 0) + t_3(-3, 0, 1, 0, 0) + t_4(1, 0, 0, 1, 0) + t_5(-2, 0, 0, 0, 1) \end{aligned}$$

Thus, the basis for V , β_V is given by

$$\beta_V = \{ (2, 1, 0, 0, 0), (-3, 0, 1, 0, 0), (1, 0, 0, 1, 0), (-2, 0, 0, 0, 1) \}$$

Given set $S = \{v\} = \{ (1, 2, 1, 0, 0) \}$:

$$\begin{pmatrix} 1 & 2 & -3 & 1 & -2 \\ 2 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow{\substack{R_3 - R_1 \rightarrow R_3 \\ 2R_1 - R_2 \rightarrow R_2}} \begin{pmatrix} 1 & 2 & -3 & 1 & -2 \\ 0 & 3 & -6 & 2 & -4 \\ 0 & -2 & 4 & -1 & 2 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{aligned} &\xrightarrow{\substack{R_3 + 2R_5 \rightarrow R_3 \\ 2R_5 + R_4 \rightarrow R_5}} \begin{pmatrix} 3 & 0 & 3 & 0 & 0 \\ 0 & 3 & -6 & 0 & 0 \\ 0 & 0 & 0 & 1 & -2 \\ 0 & 0 & 0 & 0 & -2 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow{\substack{R_3 - R_4 \rightarrow R_3 \\ R_3 + R_1 \rightarrow R_1 \\ R_2 - 2R_3 \rightarrow R_2}} \begin{pmatrix} 3 & 0 & 3 & -1 & 2 \\ 0 & 3 & -6 & 2 & -4 \\ 0 & 0 & 0 & 1 & -2 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow{\substack{3R_1 - 2R_2 \rightarrow R_1 \\ 3R_3 + 2R_2 \rightarrow R_3}} \\ &\xrightarrow{\substack{R_1 \rightarrow R_1 \\ R_2 \rightarrow R_2 \\ R_4 \rightarrow R_4}} \begin{pmatrix} 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & -2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \end{aligned}$$

Question 5 continued

Thus,

$$\{(1, 2, 1, 0, 0), (2, 1, 0, 0, 0), (1, 0, 0, 1, 0), (-2, 0, 0, 0, 1)\}$$

is a basis for V that includes v .