

# PSETs Landing Page\*

Anish Krishna Lakkapragada

This is the documentation for using my PSET PDFs responsibly. I post these LaTeX'd PSETs (1) as an education resource for friends at other universities, fellow Yalies, and all those interested and (2) for quick reference. These PSETs are not to be used irresponsibly; only look at the solution after giving each problem an honest attempt. **If YOU USE THESE PSETS TO CHEAT, YOU ARE NOT ONLY STUPID BUT YOU ARE CHEATING YOURSELF OUT OF THE ABILITY TO FALL IN LOVE WITH MATH.** Furthermore, I am not smarter than you and my solutions did not always get a perfect score.

The general format for accessing the (one-indexed) `N`th assigned PSET PDF of a Yale course with course number `CODE` is:

`https://anish.lakkapragada.com/notes/TYPE-CODE/psets/N.pdf`

where `TYPE` is `stats` or `math`. Similarly, to access my solution for this PSET you can go to:

`https://anish.lakkapragada.com/notes/TYPE-CODE/sols/N.pdf`

These PSETs and associated solution PDFs are synchronized daily at 4:20AM with my computer files through a Cronjob Shell Script. If you want to contribute any corrections, please email `anish.lakkapragada@yale.edu`.

---

\*Note that PDF here is referring to Portable Document Format, not to be confused with the veritable Probability Density Function.

# MATH 241 PSET 1

September 11, 2024

---

1.

We prove the following three inequalities and for each provide a case when the inequality would become an equality.

$$\textcircled{1} \quad P(A) + P(B) - 1 \leq P(A \cap B)$$

$$P(A) + P(B) - 1 \leq P(A \cap B)$$

$$P(A) + P(B) - P(A \cap B) \leq 1$$

$$P(A \cup B) \leq 1$$

Because we know  $P(A \cup B) \leq 1$  by axioms of probability, the inequality in  $\textcircled{1}$  is proven. We now investigate the equality case in which  $P(A) + P(B) - 1 = P(A \cap B)$

$$P(A) + P(B) - 1 = P(A \cap B)$$

$$P(A) + P(B) - P(A \cap B) = 1$$

$$P(A \cup B) = 1$$

Because  $P(A \cup A^c) = P(S) = 1$ , **the inequality would be an equality if  $B = A^c$ .**

$$\textcircled{2} \quad P(A \cap B) \leq P(A \cup B)$$

$$P(A \cap B) \leq P(A \cup B)$$

$$P(A \cap B) \leq P(A) + P(B) - P(A \cap B)$$

$$2P(A \cap B) \leq P(A) + P(B)$$

$$2P(A \cap B) \leq P(A \cap B) + P(A \cap B^c) + P(B \cap A) + P(B \cap A^c)$$

$$P(A \cap B^c) + P(B \cap A^c) \geq 0$$

Because we know  $P(A \cap B^c) \geq 0$  and  $P(B \cap A^c) \geq 0$  by axioms of probability, the inequality in (2) is proven. We now investigate the equality case in which  $P(A \cap B) = P(A \cup B)$ .

$$\begin{aligned} P(A \cap B) &= P(A \cup B) \\ P(A \cap B) &= P(A) + P(B) - P(A \cap B) \\ P(A) + P(B) &= 2P(A \cap B) \end{aligned}$$

**The inequality would be an equality if  $A = B$ .** If  $A = B$ ,  $P(A) = P(B)$  and  $A \cap B = A$  so  $P(A \cap B) = P(A)$ . Thus  $P(A) + P(B) = 2P(A) = 2P(A \cap B)$ .

$$(3) \quad P(A \cup B) \leq P(A) + P(B)$$

$$\begin{aligned} P(A \cup B) &\leq P(A) + P(B) \\ P(A) + P(B) - P(A \cap B) &\leq P(A) + P(B) \\ -P(A \cap B) &\leq 0 \\ P(A \cap B) &\geq 0 \end{aligned}$$

Because we know  $P(A \cap B) \geq 0$  by axioms of probability, the inequality in (3) is proven. We now investigate the equality case in which  $P(A \cup B) = P(A) + P(B)$ .

$$\begin{aligned} P(A \cup B) &= P(A) + P(B) \\ P(A) + P(B) - P(A \cap B) &= P(A) + P(B) \\ P(A \cap B) &= 0 \end{aligned}$$

When  $P(A \cap B) = 0$ , the equality case holds. **The inequality would be an equality if  $A$  and  $B$  are disjoint events.**

2.

Given events  $A$  and  $B$  where  $A \subseteq B$ , the set of elements of  $B$  not in  $A$  is given by  $B - A = A^c \cap B$ .

$$\begin{aligned} P(B - A) &= P(A^c \cap B) = P(A^c) + P(B) - P(A^c \cup B) \\ P(B - A) &= P(A^c \cap B) = 1 - P(A) + P(B) - P(A^c \cup B) \end{aligned}$$

Because  $P(A^c \cup B) = 1$  as  $A^c \cup B = S$  and  $P(S) = 1$ , we know  $P(B - A) = P(A^c \cap B) = 1 - P(A) + P(B) - 1$ . Thus,  $P(B - A) = P(A^c \cap B) = P(B) - P(A)$ .

3.

$$P(S|LC) = \frac{P(LC \cap S)}{P(LC)} = \frac{P(S) \cdot P(LC|S)}{P(S)P(LC|S) + P(S')P(LC|S')}$$

Given  $P(S) = 0.216$ ,  $P(S') = 1 - 0.216 = 0.784$ ,  $P(LC|S) = 23P(LC|S')$ , we can re-express  $P(S|LC)$  as:

$$P(S|LC) = \frac{23P(S)P(LC|S')}{23P(S)P(LC|S') + P(S')P(LC|S')} = \frac{23P(S)}{23P(S) + P(S')}$$

This gives us  $P(S|LC) = 0.8637$ .

4.

a)

$$P(K|R) = \frac{P(K \cap R)}{P(R)} = \frac{P(R|K) \cdot P(K)}{P(R)} = \frac{P(R|K) \cdot P(K)}{P(R|K)P(K) + P(R|K')P(K')}$$

Given  $P(R|K) = 1$ ,  $P(K) = p$ ,  $P(K') = 1 - P(K) = 1 - p$ , and  $P(R|K') = \frac{1}{n}$  as the answer is chosen randomly in the case Fred doesn't know the answer,  $P(K|R) = \frac{p}{p + \frac{1-p}{n}}$ .

b) We show  $P(K|R) \geq p$  below.

$$\begin{aligned} P(K|R) &\geq p \\ \frac{p}{p + \frac{1-p}{n}} &\geq p \\ p + \frac{1-p}{n} &\leq 1 \\ \frac{1-p}{n} &\leq 1-p \\ \frac{1-p}{1-p} &\leq n \\ n &\geq 1 \end{aligned}$$

Because all tests have at least one question,  $n \geq 1$ , and thus  $P(K|R) \geq p$ . This makes sense as  $P(K|R)$  refers to the probability that Fred knows the answer to a question given that he got the question right. Intuitively,  $P(K|R) \geq p$  as we would expect a drastically higher percentage of the questions he gets correct to be because he knew the answer to them as opposed to choosing the question's correct answer by luck.

The only case in which  $P(K|R) = p$  is when  $n = 1$ .

5.

$$P(A|D) = \frac{P(A \cap D)}{P(D)} = \frac{P(D|A) \cdot P(A)}{P(D)} = \frac{P(D|A) \cdot P(A)}{P(D|A)P(A) + P(D|B)P(B) + P(D|C)P(C)}$$

Given that  $P(A) = 0.5, P(B) = 0.3, P(C) = 0.2$  and  $P(D|A) = 0.01, P(D|B) = 0.02, P(D|C) = 0.03, P(A|D) = 0.294$ .

6. Anish Lakkapragada. I worked independently.