

Discretionary Note

Anish Krishna Lakkapragada

IF YOU USE THIS FILE TO CHEAT, YOU ARE NOT ONLY STUPID BUT YOU ARE CHEATING YOURSELF OUT OF THE ABILITY TO FALL IN LOVE WITH MATH. Furthermore, I am not smarter than you and my solutions did not always get a perfect score.

CONTENT STARTS ON NEXT PAGE.

To access the general instructions for this repository head [here](#).

S&DS 242/542: Homework 2

Due Wednesday, January 29, at 1PM

1. **Binomial MGF.** Let $X \sim \text{Binomial}(n, p)$. Derive a simple expression for the moment generating function of X in terms of n and p .

2. **Construction of the bivariate normal.** Let Z_1 and Z_2 be two independent $\mathcal{N}(0, 1)$ variables, and consider the bivariate normal vector (X_1, X_2) given by

$$\begin{aligned}X_1 &= c_1 Z_1 + d_1 Z_2 + e_1 \\X_2 &= c_2 Z_1 + d_2 Z_2 + e_2\end{aligned}$$

where $c_1, c_2, d_1, d_2, e_1, e_2 \in \mathbb{R}$ are fixed constants.

Given any values $\mu_1, \mu_2 \in \mathbb{R}$, $\sigma_1^2, \sigma_2^2 > 0$, and $\rho \in [-1, 1]$, show how one may choose $c_1, c_2, d_1, d_2, e_1, e_2$ so that X_1 and X_2 have means μ_1 and μ_2 , variances σ_1^2 and σ_2^2 , and correlation ρ .

3. **Monte Carlo integration (based on Rice 5.21 and 5.22).** For a given function $f : [a, b] \rightarrow \mathbb{R}$, suppose we wish to numerically approximate

$$I(f) = \int_a^b f(x) dx.$$

A Monte Carlo approach to doing this is the following: Let g be the PDF of any continuous random variable taking values in $[a, b]$, where $g(x) > 0$ for all $x \in [a, b]$. Generate independent random draws X_1, \dots, X_n from g . Then approximate $I(f)$ by

$$\hat{I}_n(f) = \frac{1}{n} \sum_{i=1}^n \frac{f(X_i)}{g(X_i)}.$$

(a) Show that $\mathbb{E}[\hat{I}_n(f)] = I(f)$. Assuming that $\text{Var}[f(X_i)/g(X_i)] < \infty$, explain why $\hat{I}_n(f) \rightarrow I(f)$ in probability as $n \rightarrow \infty$.

(b) Derive a formula for $\text{Var}[\hat{I}_n(f)]$. Show that for some $c_n \in \mathbb{R}$, $c_n(\hat{I}_n(f) - I(f)) \rightarrow \mathcal{N}(0, 1)$ in distribution as $n \rightarrow \infty$.

(c) Consider concretely the task of approximating

$$I(f) = \int_0^1 \cos(2\pi x) dx.$$

If g is the uniform distribution on $[0, 1]$, provide an explicit form for the above value c_n .

4. **Simulating a sample median.** Let $X_1, \dots, X_{99} \stackrel{IID}{\sim} \mathcal{N}(0, 1)$. The sample median X_{median} is the 50th largest value among X_1, \dots, X_{99} .

Compute X_{median} for 5000 simulations of X_1, \dots, X_{99} . What is the mean of X_{median} across your simulations? What is the standard deviation? Plot a histogram of the 5000 values of X_{median} — what does the sampling distribution of X_{median} look like?

Derive (analytically) the standard deviation of the sample mean $\bar{X} = \frac{X_1 + \dots + X_{99}}{99}$, and compare this with your simulated standard deviation of X_{median} . According to your simulation, is X_{median} more or less variable than \bar{X} ? Include both your code and your histogram in your homework submission.

If you are new to programming, the following will walk you through how to do this in R:

- Install R from <https://www.r-project.org/>. Launch R and select “New Document” from the “File” drop-down menu.

- We will write our code in this document. First, let’s create a numeric vector of length 5000 that will save the results from the 5000 simulations. Call it `X.median`:

```
X.median = numeric(5000)
```

- To repeat a series of commands 5000 times, we’ll use a `for` loop:

```
for (i in 1:5000) {  
  ...  
}
```

We can fill in any commands in the “...” section above, and these will be executed once for each value of `i` from 1 to 5000.

- Let’s fill in the ... section. We can simulate 99 independent samples from $\mathcal{N}(0, 1)$ using the `rnorm` function in R, and save it to a temporary vector variable `X`:

```
X = rnorm(99, mean=0, sd=1)
```

(The `mean` and `sd` arguments indicate the mean and standard deviation of the normal distribution.) We can then use the `median` function in R to compute the sample median of the values in `X`. We will save this as `X.median[i]`, referring to element `i` of the numeric vector we created above:

```
X.median[i] = median(X)
```

(Hence, in the first loop iteration the sample median is saved as `X.median[1]`, in the second iteration it is saved as `X.median[2]`, etc.) Let's put the above two lines of code inside the `for` loop.

- After the `for` loop, we can now write some commands that compute and print the mean and standard deviation of the values in `X.median`, and plot a histogram of these values:

```
print(mean(X.median))
print(sd(X.median))
hist(X.median)
```

- Let's save our file as `sample_median.R`. Then go back to the main R Console, and select "Source File..." under the "File" drop-down menu. Select our file `sample_median.R`, and voila! You should see the the mean and standard deviation of the 5000 sample medians printed in the R Console, and a separate plot window displaying the histogram.

We'll use more built-in functions in R as we go. To see what a function does and how to use it, type `?` followed by the function name in the R console to pull up the help page. For example, entering

```
?median
```

into the R console pulls up a page about the `median` function used above.