

## Discretionary Note

Anish Krishna Lakkapragada

**IF YOU USE THIS FILE TO CHEAT, YOU ARE NOT ONLY STUPID BUT YOU ARE CHEATING YOURSELF OUT OF THE ABILITY TO FALL IN LOVE WITH MATH.** Furthermore, I am not smarter than you and my solutions did not always get a perfect score.

**CONTENT STARTS ON NEXT PAGE.**

To access the general instructions for this repository head [here](#).

# STATS 242 HW 2

January 29, 2025

---

**Number of late days: 0; Collaborators: None.**

1.

For  $1 \leq i \leq n$ , let us define r.v.  $B_i \sim \text{Bern}(p)$ , where for  $i \neq j$ ,  $B_i$  and  $B_j$  are independent. Then, we have that  $X = \sum_{i=1}^n B_i$ . Because all  $B_i$ s are independent, we have that:

$$M_X(t) = M_{\sum_{i=1}^n B_i}(t) = \prod_{i=1}^n M_{B_i}(t)$$

The MGF of r.v.  $B_i$  can be computed as:  $M_{B_i}(t) = \mathbb{E}[e^{tB_i}] = e^{t(1)}P(B_i = 1) + e^{t(0)}P(B_i = 0) = pe^t + (1 - p) = 1 + p(e^t - 1)$ . Thus, we have  $M_X(t)$  as:

$$M_X(t) = \prod_{i=1}^n M_{B_i}(t) = \prod_{i=1}^n [1 + p(e^t - 1)] = [1 + p(e^t - 1)]^n$$

2.

We first start by computing the distributions of  $X_1$  and  $X_2$ . Note that because  $Z_1$  and  $Z_2$  are independent normal distributions, their sum forms a normal distribution as well.

## 1. Distribution of $X_1$

Since  $c_1 Z_1 \sim \mathcal{N}(0, c_1^2)$  and  $d_1 Z_1 \sim \mathcal{N}(0, d_1^2)$ ,  $c_1 Z_1 + d_1 Z_2 \sim \mathcal{N}(0, c_1^2 + d_1^2)$ . Finally,  $e_1$  is just a constant and so it doesn't affect the variance so  $X_1 = c_1 Z_1 + d_1 Z_2 + e_1 \sim \mathcal{N}(e_1, c_1^2 + d_1^2)$ . Therefore we can set  $e_1 = \mu_1$ , the mean of  $X_1$ . Furthermore, the variance  $\sigma_1^2$  of  $X_1$  is equal to  $c_1^2 + d_1^2$ .

## 2. Distribution of $X_2$

We use identical reasoning as with before.  $c_2 Z_1 \sim \mathcal{N}(0, c_2^2)$  and  $d_2 Z_2 \sim \mathcal{N}(0, d_2^2)$ , so we have  $c_2 Z_2 + d_2 Z_2 \sim \mathcal{N}(0, c_2^2 + d_2^2)$ . Thus,  $X_2 = c_2 Z_2 + d_2 Z_2 + e_2 \sim \mathcal{N}(e_2, c_2^2 + d_2^2)$ . Therefore we can set  $e_2 = \mu_2$ , the mean of  $X_2$ . Furthermore, the variance  $\sigma_2^2$  of  $X_2$  is equal to  $c_2^2 + d_2^2$ .

We now have  $c_1, c_2, d_1, d_2$  remaining to assign. We compute the correlation  $\rho$  between  $X_1$  and  $X_2$ , starting by computing the covariance between  $X_1$  and  $X_2$ :

$$\begin{aligned}\text{Cov}(X_1, X_2) &= \text{Cov}(c_1 Z_1 + d_1 Z_2 + e_1, c_2 Z_1 + d_2 Z_2 + e_2) \\ &= c_1 \text{Cov}(Z_1, c_2 Z_1 + d_2 Z_2 + e_2) + d_1 \text{Cov}(Z_2, c_2 Z_1 + d_2 Z_2 + e_2) + \text{Cov}(e_1, c_2 Z_1 + d_2 Z_2 + e_2)\end{aligned}$$

Note that because  $e_1$  is a fixed constant,  $\text{Cov}(e_1, c_2 Z_1 + d_2 Z_2 + e_2) = 0$ . Also note that because  $Z_1$  and  $Z_2$  are independent,  $\text{Cov}(Z_1, Z_2) = 0$ .

$$\begin{aligned}\text{Cov}(X_1, X_2) &= c_1[c_2 \text{Cov}(Z_1, Z_1) + d_2 \text{Cov}(Z_1, Z_2) + \text{Cov}(Z_1, e_2)] + d_1 \text{Cov}(Z_2, c_2 Z_1 + d_2 Z_2 + e_2) \\ &= c_1[c_2 + 0 + 0] + d_1[c_2 \text{Cov}(Z_2, Z_1) + d_2 \text{Cov}(Z_2, Z_2) + \text{Cov}(Z_2, e_2)] = c_1 c_2 + d_1[0 + d_2 + 0] \\ &= c_1 c_2 + d_1 d_2\end{aligned}$$

and now we compute the correlation  $\rho$  between  $X_1$  and  $X_2$  as:

$$\rho = \frac{\text{Cov}(X_1, X_2)}{\sqrt{\text{Var}(X_1)\text{Var}(X_2)}} = \frac{c_1 c_2 + d_1 d_2}{\sigma_1 \sigma_2}$$

This gives us three equations:

$$\begin{aligned}\rho \sigma_1 \sigma_2 &= c_1 c_2 + d_1 d_2 \\ c_1^2 + d_1^2 &= \sigma_1^2 \\ c_2^2 + d_2^2 &= \sigma_2^2\end{aligned}$$

to solve for four variables. As such, we arbitrarily choose to set  $c_1 = 0$ , giving us  $d_1 = \sigma_1$  and  $\rho \sigma_1 \sigma_2 = d_1 d_2 = \sigma_1 d_2 \implies d_2 = \rho \sigma_2$ . Finally, we can solve for  $c_2^2 = \sigma_2^2 - d_2^2 = \sigma_2^2 - \rho^2 \sigma_2^2 = \sigma_2^2(1 - \rho^2) \implies c_2 = \sigma_2 \sqrt{1 - \rho^2}$ .

As a summary of my answer,

$$\begin{aligned}e_1 &= \mu_1 \\ e_2 &= \mu_2 \\ c_1 &= 0 \\ c_2 &= \sigma_2 \sqrt{1 - \rho^2} \\ d_1 &= \sigma_1 \\ d_2 &= \rho \sigma_2\end{aligned}$$

3.

- (a) We first show  $\mathbb{E}[\hat{I}_n(f)] = I(f)$ . Note that  $\frac{f(X_i)}{g(X_i)}$  is a random variable and so its expectation is given to us by  $\mathbb{E}[\frac{f(X_i)}{g(X_i)}] = \int_{-\infty}^{\infty} \frac{f(u)}{g(u)} g(u) du = \int_a^b \frac{f(u)}{g(u)} g(u) du = \int_a^b f(u) du = I(f)$ . Thus:

$$\mathbb{E}[\hat{I}_n(f)] = \mathbb{E}[\frac{1}{n} \sum_{i=1}^n \frac{f(X_i)}{g(X_i)}] = \frac{1}{n} \cdot n \cdot \mathbb{E}[\frac{f(X_i)}{g(X_i)}] = I(f)$$

Notice that  $\hat{I}_n(f)$  is essentially an average of  $n$  random variables, each of with an expectation of  $I(f)$ . Thus,  $\hat{I}_n(f) \rightarrow I(f)$  in probability as  $n \rightarrow \infty$  due to the (Weak) Law of Large Numbers.

- b) We first compute  $\text{Var}[\hat{I}_n(f)]$ . We begin by computing the variance of random variable  $\frac{f(X_i)}{g(X_i)}$ :

$$\text{Var}[\frac{f(X_i)}{g(X_i)}] = \mathbb{E}[(\frac{f(X_i)}{g(X_i)})^2] - \mathbb{E}[\frac{f(X_i)}{g(X_i)}]^2 = \int_a^b \frac{f^2(u)}{g^2(u)} g(u) du - I^2(f) = \int_a^b \frac{f^2(u)}{g(u)} du - I^2(f)$$

Let us call define this quantity to be  $\sigma^2 = \int_a^b \frac{f^2(u)}{g(u)} du - I^2(f) \in \mathbb{R}$ . and so we have:

$$\text{Var}[\hat{I}_n(f)] = \text{Var}[\frac{1}{n} \sum_{i=1}^n \frac{f(X_i)}{g(X_i)}] = \frac{1}{n^2} \sum_{i=1}^n \text{Var}[\frac{f(X_i)}{g(X_i)}] = \frac{1}{n} \sigma^2$$

Let us define  $c_n = \frac{\sqrt{n}}{\sigma} \in \mathbb{R}$ . Thus, by the Central Limit Theorem, we have that:

$$c_n(\hat{I}_n(f) - I(f)) \rightarrow \mathcal{N}(0, 1) \text{ as } n \rightarrow \infty$$

- c) In this problem,  $a = 0$  and  $b = 1$ ,  $f(x) = \cos(2\pi x)$ , and  $g(x) = 1$  if  $x \in [0, 1]$  and 0 otherwise. We first compute  $I(f)$ :

$$I(f) = \int_0^1 \cos(2\pi x) dx = \sin(2\pi x) \Big|_0^1 = \sin(2\pi) - \sin(0) = 0 - 0 = 0$$

and then  $\sigma^2$ :


$$\begin{aligned} \sigma^2 &= \int_0^1 \frac{\cos^2(2\pi x)}{1} dx - I^2(f) = \int_0^1 [\frac{1}{2} + \frac{\cos(4\pi x)}{2}] dx = [\frac{x}{2} + \frac{\sin(4\pi x)}{8}] \Big|_0^1 \\ &= \frac{1}{2} + \frac{1}{8}(\sin(4\pi) - \sin(0)) = \frac{1}{2} \end{aligned}$$

and so  $c_n$  is given by:

$$c_n = \frac{\sqrt{n}}{\sigma} = \sqrt{\frac{n}{\sigma^2}} = \sqrt{2n}$$

4.

From my simulation, the mean and standard deviation of  $X_{median}$  are given by 0.00178 and 0.1265 respectively. Below is the histogram of the 5000 values of  $X_{median}$  from my simulation:



Based on the above histogram, we can see that the sampling distribution of  $X_{median}$  follows a normal distribution. We now derive the standard deviation of the sample mean  $\bar{X}$ . Note that 99 observations is enough for us to apply the Central Limit Theorem with confidence. Thus<sup>1</sup>,  $\bar{X} = \frac{X_1 + \dots + X_{99}}{99} \sim \mathcal{N}(\mathbb{E}[X_1], \frac{\text{Var}(X_1)}{99})$  or  $\bar{X} \sim \mathcal{N}(0, \frac{1}{99})$ . Thus the analytically-computed standard deviation of sample mean  $\bar{X}$  is  $\sqrt{\frac{1}{99}}$  or 0.1005, which is less than my calculated simulated standard deviation for  $X_{median}$  (0.1265). According to my simulation,  $X_{median}$  is more variable than  $\bar{X}$ .

```

1  # %%
2
3  # Run all imports first and then write helper functions.
4
5  import numpy as np
6  import math
7  import matplotlib.pyplot as plt
8
9  def get_N_obs_iid_standard_normal(N):
10     return np.random.normal(0, 1, N)
11
12  def compute_median(arr):
13     # assume arr is numpy
14     return np.median(arr)

```

<sup>1</sup>Note that  $X_1 \dots X_{99}$  are identical distributions and so they all have the same mean and variances.

```

15 # %%
16 N_SIMULATIONS = 5000
17 X_medians = []
18 for _ in range(N_SIMULATIONS):
19     samples = get_N_obs_iid_standard_normal(99)
20     X_median_curr = compute_median(samples)
21     X_medians.append(X_median_curr)
22
23 X_medians = np.array(X_medians)
24 X_median_mean = np.mean(X_medians)
25 X_median_std = np.std(X_medians)
26
27 print(f"X_median mean: {X_median_mean} and std: {X_median_std}")
28 # %%
29
30 """
31 Plot a histogram.
32 """
33
34 plt.hist(X_medians, bins=100, edgecolor='black', alpha=0.7)
35
36 # Add labels and title
37 plt.xlabel('X_median values')
38 plt.ylabel('Frequency')
39 plt.title('Histogram of 5000 values of X_median')
40
41 # %%

```