

PSETs Landing Page*

Anish Krishna Lakkapragada

This is the documentation for using my PSET PDFs responsibly. I post these LaTeX'd PSETs (1) as an education resource for friends at other universities, fellow Yalies, and all those interested and (2) for quick reference. These PSETs are not to be used irresponsibly; only look at the solution after giving each problem an honest attempt. **If YOU USE THESE PSETS TO CHEAT, YOU ARE NOT ONLY STUPID BUT YOU ARE CHEATING YOURSELF OUT OF THE ABILITY TO FALL IN LOVE WITH MATH.** Furthermore, I am not smarter than you and my solutions did not always get a perfect score.

The general format for accessing the (one-indexed) `N`th assigned PSET PDF of a Yale course with course number `CODE` is:

`https://anish.lakkapragada.com/notes/TYPE-CODE/psets/N.pdf`

where `TYPE` is `stats` or `math`. Similarly, to access my solution for this PSET you can go to:

`https://anish.lakkapragada.com/notes/TYPE-CODE/sols/N.pdf`

These PSETs and associated solution PDFs are synchronized daily at 4:20AM with my computer files through a Cronjob Shell Script. If you want to contribute any corrections, please email anish.lakkapragada@yale.edu.

*Note that PDF here is referring to Portable Document Format, not to be confused with the veritable Probability Density Function.

MATH 255 HW 1

January 23, 2025

1. Exercise 1.1 (5 points)

- (1) We prove this by contrapositive and thus assume f is not injective $\Rightarrow \exists x, x' \in A$ s.t. $x \neq x'$ and $f(x) = f(x')$. Let us define $u = f(x) = f(x')$. Then, h is not injective as $h(x) = g(f(x)) = g(u)$ and $h(x') = g(f(x')) = g(u)$ and so $\exists x, x' \in A$ s.t. $x \neq x'$ and $h(x) = h(x')$.
- (2) We prove this by contrapositive and thus assume that g is not surjective $\Rightarrow \exists c \in C$ s.t. $\nexists b \in B$ where $g(b) = c$. Because no element in B maps to C by g , this means $\forall x \in A, h(x) = g(f(x)) \neq c$ and so because $\nexists x \in A$ s.t. $h(x) = c$, h is not surjective.

2. Exercise 1.2 (5 points; Rudin 1.1)

- (1) Prove $r + x \notin \mathbb{Q}$

Let us prove this by contradiction and assume that $r + x \in \mathbb{Q}$. By negation rule, $r \in \mathbb{Q} \Rightarrow -r \in \mathbb{Q}$. By addition rule, $(-r) + (r + x) \in \mathbb{Q}$ and so this means $(-r) + (r + x) = (-r + r) + x = 0 + x = x \in \mathbb{Q}$, which is a contradiction.

- (2) Prove $rx \notin \mathbb{Q}$

Let us prove this by contradiction and assume that $rx \in \mathbb{Q}$. Because $r \neq 0$, by the inversion rule we have that r has an inverse r^{-1} . By multiplication rule, $(r^{-1}) \cdot rx \in \mathbb{Q}$ or $(r^{-1}) \cdot rx = (r^{-1} \cdot r)x = 1 \cdot x = x \in \mathbb{Q}$, which is a contradiction.

3. Exercise 1.3 (10 points; Rudin 1.3)

- (1) Because $x \neq 0$, by the inversion rule we know $\exists x^{-1}$ s.t. $x^{-1} \cdot x = 1$. Thus:

$$\begin{aligned}xy &= xz \\x^{-1}xy &= x^{-1}xz \\1 \cdot y &= 1 \cdot z \\y &= z\end{aligned}$$

(2) Because $x \neq 0$, we can apply the inversion rule again:

$$\begin{aligned} xy &= x \\ x^{-1}xy &= x^{-1}x \\ 1 \cdot y &= 1 \\ y &= 1 \end{aligned}$$

(3) We first prove that $0 \cdot y = 0$:

$$\begin{aligned} 0 \cdot y + 0 \cdot y &= (0 + 0) \cdot y = 0 \cdot y \\ 0 \cdot y + 0 \cdot y &= 0 \cdot y \end{aligned}$$

Because $0 \cdot y \in F$, its additive inverse is given by $-0 \cdot y$:

$$\begin{aligned} 0 \cdot y + 0 \cdot y - 0 \cdot y &= 0 \cdot y - 0 \cdot y \\ 0 \cdot y + (0 \cdot y - 0 \cdot y) &= 0 \\ 0 \cdot y + 0 &= 0 \\ 0 \cdot y &= 0 \end{aligned}$$

To prove that if $xy = 1 \implies x \neq 0$ we proceed by contradiction. If $x = 0$, then $xy = 0 \cdot y = 0 \neq 1$. We now show $y = x^{-1}$. Thus, $x \neq 0$. Because $x \neq 0$, we can apply the inversion rule again:

$$\begin{aligned} xy &= 1 \\ x^{-1}xy &= x^{-1} \\ 1 \cdot y &= x^{-1} \\ y &= x^{-1} \end{aligned}$$

(4) Let us define inverse of x^{-1} to be u , where by definition $x^{-1} \cdot u = 1$. Because $x^{-1} \cdot x = 1$ by definition, then we have that $u = x$, or that the inverse of x^{-1} is x . Expressed as an equation, we have shown: $(x^{-1})^{-1} = x$.

4. Exercise 1.4 (10 points)

(1) Let us suppose $x = \frac{p}{q} \in \mathbb{Q}$, where $p, q \in \mathbb{Z}$ and $\frac{p}{q}$ are in lowest terms. For proof by contradiction, we assume $x^2 = 3$. This means $\frac{p^2}{q^2} = 3$ or $p^2 = 3q^2$ and thus p^2 has a factor of three.

We now prove that p has a factor of three. Because p^2 has a factor of three, the prime factorization of p^2 can be given by $3^\alpha \dots$ where $\alpha \geq 1 \in \mathbb{Z}$. Let us define the prime factorization of $p = 3^\beta \dots$ where $\beta \geq 0 \in \mathbb{Z}$. Note that since $p^2 = (3^\beta \dots)^2$,

$\alpha = 2\beta$. The lowest possible integer value of α s.t. $\alpha \geq 1$ and $\beta \in \mathbb{Z}$ is then $\alpha = 2$ and $\beta = 1$, so we are guaranteed that p has a factor of three. This means we can express $p = 3k$, where $k \in \mathbb{Z}$ and so $p^2 = (3k)^2 = 9k^2 = 3q^2$ or $q^2 = 3k^2$. Using the same logic as before because q^2 has a factor of three, so does q . Thus, p and q both have a factor of three and so this contradicts the assumption that $\frac{p}{q}$ are in lowest terms.

(2) We show that these provided operations define $\mathbb{Q}(\sqrt{3})$ as a field:

1. **Zero & One Element**

Our zero and one element in $\mathbb{Q}(\sqrt{3})$ are given by $0+0\sqrt{3}$ and $1+0\sqrt{3}$ respectively. Written as ordered pairs, they are given by $(0, 0)$ and $(1, 0)$ respectively.

2. **Negation Law**

For an element $u = a + b\sqrt{3} \in \mathbb{Q}(\sqrt{3})$, $-u = (-a) + (-b)\sqrt{3} \in \mathbb{Q}(\sqrt{3})$.

3. **Inversion Law**

For an element $u = a + b\sqrt{3} \in \mathbb{Q}$, u^{-1} is given by:

$$\begin{aligned} uu^{-1} &= 1_{\mathbb{Q}\sqrt{3}} = 1 + 0\sqrt{3} \\ u^{-1} &= \frac{1 + 0\sqrt{3}}{u} = \frac{1 + 0\sqrt{3}}{a + b\sqrt{3}} = \frac{(1 + 0\sqrt{3})(a - b\sqrt{3})}{(a + b\sqrt{3})(a - b\sqrt{3})} = \frac{(1 + 0\sqrt{3})(a - b\sqrt{3})}{(a^2 - 3b^2) + (-ab + ba)\sqrt{3}} \\ &= \frac{a - b\sqrt{3}}{a^2 - 3b^2} = \frac{a}{a^2 - 3b^2} + \left(\frac{-b}{a^2 - 3b^2}\right)\sqrt{3} \end{aligned}$$

4. $\forall x, y \in \mathbb{Q}\sqrt{3}, x + y = y + x$

Define $x = a + b\sqrt{3}, y = a' + b'\sqrt{3} \in \mathbb{Q}\sqrt{3}$. $x + y$ is given by:

$$x + y = (a + b\sqrt{3}) + (a' + b'\sqrt{3}) = (a + a') + (b + b')\sqrt{3}$$

and $y + x$ is given by:

$$y + x = (a' + b'\sqrt{3}) + (a + b\sqrt{3}) = (a' + a) + (b' + b)\sqrt{3} = (a' + a) + (b + b')\sqrt{3}$$

and so $x + y = y + x$.

5. $\forall x, y, z \in \mathbb{Q}\sqrt{3}, (x + y) + z = x + (y + z)$

We define x, y the same as above. We define $z = a'' + b''\sqrt{3} \in \mathbb{Q}\sqrt{3}$. Then $(x + y) + z$ is given by:

$$\begin{aligned} (x + y) + z &= [(a + a') + (b + b')\sqrt{3}] + z = ((a + a') + (b + b')\sqrt{3}) + (a'' + b''\sqrt{3}) \\ &= ((a + a') + a'') + ((b + b') + b'')\sqrt{3} = (a + a' + a'') + (b + b' + b'')\sqrt{3} \end{aligned}$$

and $x + (y + z)$ is given by:

$$\begin{aligned} x + (y + z) &= x + ((a' + b'\sqrt{3}) + (a'' + b''\sqrt{3})) = x + ((a' + a'') + (b' + b'')\sqrt{3}) \\ &= (a + b\sqrt{3}) + ((a' + a'') + (b' + b'')\sqrt{3}) = (a + a' + a'') + (b + b' + b'')\sqrt{3} \end{aligned}$$

and so $(x + y) + z = x + (y + z)$.

6. $\forall x \in \mathbb{Q}\sqrt{3}, 0 + x = x$

Using the previous definition of x :

$$0_{\mathbb{Q}\sqrt{3}} + x = (0 + 0\sqrt{3}) + (a + b\sqrt{3}) = (0 + a) + (0 + b)\sqrt{3} = a + b\sqrt{3} = x$$

7. $\forall x \in \mathbb{Q}\sqrt{3}, -x + x = 0$.

Using the previous definition of x :

$$-x + x = ((-a) + (-b)\sqrt{3}) + (a + b\sqrt{3}) = ((-a + a) + (-b + b)\sqrt{3}) = 0 + 0\sqrt{3} = 0_{\mathbb{Q}\sqrt{3}}$$

8. $\forall x, y \in \mathbb{Q}\sqrt{3}, xy = yx$

We define x, y the same as before. This gives us xy as:

$$xy = (a + b\sqrt{3})(a' + b'\sqrt{3}) = (aa' + 3bb') + (ab' + ba')\sqrt{3}$$

and yx as:

$$\begin{aligned} yx &= (a' + b'\sqrt{3})(a + b\sqrt{3}) = (a'a + 3b'b) + (a'b + b'a)\sqrt{3} = (aa' + 3bb') + (ba' + ab')\sqrt{3} \\ &= (aa' + 3bb') + (ab' + ba')\sqrt{3} \end{aligned}$$

and so $xy = yx$.

9. $\forall x, y, z \in \mathbb{Q}\sqrt{3}, (xy)z = x(yz)$

We use the previous definitions of x, y, z as before. This gives us $(xy)z$ as:

$$\begin{aligned} (xy)z &= ((aa' + 3bb') + (ab' + ba')\sqrt{3})(a'' + b''\sqrt{3}) \\ &= (a''(aa' + 3bb') + 3(ab' + ba')b'') + ((aa' + 3bb')b'' + (ab' + ba')a'')\sqrt{3} \end{aligned}$$

and $x(yz)$ as:

$$\begin{aligned} x(yz) &= x((a' + b'\sqrt{3})(a'' + b''\sqrt{3})) = x((a'a'' + 3b'b'') + (a'b'' + b'a'')\sqrt{3}) \\ &= (a + b\sqrt{3})((a'a'' + 3b'b'') + (a'b'' + b'a'')\sqrt{3}) \\ &= (a(a'a'' + 3b'b'') + 3b(a'b'' + b'a'')) + (a(a'b'' + b'a'') + b(a'a'' + 3b'b''))\sqrt{3} \\ &= (a''(aa' + 3bb') + 3(ab' + ba')b'') + ((aa' + 3bb')b'' + (ab' + ba')a'')\sqrt{3} \end{aligned}$$

and so $(xy)z = x(yz)$.

10. $\forall x \in \mathbb{Q}, 1 \cdot x = x$

Using previous definition of x :

$$1_{\mathbb{Q}\sqrt{3}} \cdot x = (1 + 0\sqrt{3})(a + b\sqrt{3}) = (1 \cdot a + 3 \cdot 0 \cdot b) + (1 \cdot b + 0 \cdot a)\sqrt{3} = a + b\sqrt{3} = x$$

11. $\forall x \in \mathbb{Q}\sqrt{3}$ with $x \neq 0, x \cdot x^{-1} = 1$

Using the previous definition of x :

$$\begin{aligned} x \cdot x^{-1} &= (a + b\sqrt{3})\left(\frac{a}{a^2 - 3b^2} + \left(\frac{-b}{a^2 - 3b^2}\right)\sqrt{3}\right) = \\ &= \left(\frac{a^2}{a^2 - 3b^2} - \frac{3b^2}{a^2 - 3b^2}\right) + \left(\frac{-ab}{a^2 - 3b^2} + \frac{ba}{a^2 - 3b^2}\right)\sqrt{3} \\ &= \frac{a^2 - 3b^2}{a^2 - 3b^2} + \left(\frac{ab - ab}{a^2 - 3b^2}\right)\sqrt{3} = 1 + 0\sqrt{3} = 1_{\mathbb{Q}\sqrt{3}} \end{aligned}$$

12. $\forall x, y, z \in \mathbb{Q}\sqrt{3}, x(y + z) = xy + xz$

We use the previous definitions for $x, y, z \in \mathbb{Q}\sqrt{3}$. This gives us $x(y + z)$ as:

$$\begin{aligned} x(y + z) &= x((a' + b'\sqrt{3}) + (a'' + b''\sqrt{3})) = x((a' + a'') + (b' + b'')\sqrt{3}) \\ &= (a + b\sqrt{3})((a' + a'') + (b' + b'')\sqrt{3}) \\ &= (a(a' + a'') + 3b(b' + b'')) + (a(b' + b'') + b(a' + a''))\sqrt{3} \end{aligned}$$

and $xy + xz$ as:

$$\begin{aligned} xy + xz &= ((aa' + 3bb') + (ab' + ba')\sqrt{3}) + xz \\ &= ((aa' + 3bb') + (ab' + ba')\sqrt{3}) + ((a + b\sqrt{3})(a'' + b''\sqrt{3})) \\ &= ((aa' + 3bb') + (ab' + ba')\sqrt{3}) + ((aa'' + 3bb'') + (ab'' + ba'')\sqrt{3}) \\ &= ((aa' + 3bb' + aa'' + 3bb'') + (ab' + ba' + ab'' + ba'')\sqrt{3}) \\ &= (a(a' + a'') + 3b(b' + b'')) + (a(b' + b'') + b(a' + a''))\sqrt{3} \end{aligned}$$

and so $x(y + z) = xy + xz$.

(3) Given these addition and product laws, the inversion rule we defined in (2) for $\mathbb{Q}\sqrt{3}$ looks like:

$$(a + b\sqrt{3})^{-1} = \frac{a}{a^2 - 3b^2} + \left(\frac{-b}{a^2 - 3b^2}\right)\sqrt{3}$$

and the zero element was given by $0 + 0\sqrt{3}$.

Now consider element $x = 2 + 0\sqrt{3} \neq 0 + 0\sqrt{3}$, where $x \in \mathbb{Z}\sqrt{3}$. The inverse of x is given by $x^{-1} = \frac{2}{4} + 0\sqrt{3}$. Because $\frac{2}{4} \notin \mathbb{Z}, x^{-1} \notin \mathbb{Z}$ and so the inversion rule does not apply for $\mathbb{Z}\sqrt{3}$ with the provided addition and product rules.

5. **Exercise 1.5 (5 points)**

We prove that \prec does not make \mathbb{Q} into an ordered set by a contradictory example. Consider $x = \frac{1}{6}$ and $y = \frac{2}{3}$. For these values, $x \not\prec y$ and $y \not\prec x$. Furthermore, $x \neq y$ because they are not the same element in the set \mathbb{Q} (this set is reduced to lowest terms, so two elements are equivalent only if their numerator and denominator are the same.) Thus, we have shown for $x, y \in \mathbb{Q}$, none of the following statements are true: $x \prec y, y \prec x, x = y$ and so \prec does not make \mathbb{Q} an ordered set.

6. **Exercise 1.6 (5 points)**

Theorem 0.1 *If S is an ordered set with elements $x, y, z \in S$, $x \leq y$ and $y \leq z \implies x \leq z$.*

Proof: We do casework:

1. **Case 1:** $x < y$

We investigate the two subcases:

1. **Subcase 1:** $y < z$

By the transitivity property of ordered sets, $x < y$ and $y < z \implies x < z \implies x \leq z$.

2. **Subcase 2:** $y = z$

We are given $x < y$. Because $y = z$, $x < z \implies x \leq z$.

2. **Case 2:** $x = y$

We are given $y \leq z$. Because $x = y$, we can conclude $x \leq z$.

Because A is a non-empty set, $\exists x \in A$. Pick any $x \in A$. Because α is a lower bound and β is an upper bound $\alpha \leq x \leq \beta \implies \alpha \leq \beta$ by **Theorem 0.1**.