

# PSETs Landing Page\*

Anish Krishna Lakkapragada

This is the documentation for using my PSET PDFs responsibly. I post these LaTeX'd PSETs (1) as an education resource for friends at other universities, fellow Yalies, and all those interested and (2) for quick reference. These PSETs are not to be used irresponsibly; only look at the solution after giving each problem an honest attempt. **If YOU USE THESE PSETS TO CHEAT, YOU ARE NOT ONLY STUPID BUT YOU ARE CHEATING YOURSELF OUT OF THE ABILITY TO FALL IN LOVE WITH MATH.** Furthermore, I am not smarter than you and my solutions did not always get a perfect score.

The general format for accessing the (one-indexed) `N`th assigned PSET PDF of a Yale course with course number `CODE` is:

`https://anish.lakkapragada.com/notes/TYPE-CODE/psets/N.pdf`

where `TYPE` is `stats` or `math`. Similarly, to access my solution for this PSET you can go to:

`https://anish.lakkapragada.com/notes/TYPE-CODE/sols/N.pdf`

These PSETs and associated solution PDFs are synchronized daily at 4:20AM with my computer files through a Cronjob Shell Script. If you want to contribute any corrections, please email `anish.lakkapragada@yale.edu`.

---

\*Note that PDF here is referring to Portable Document Format, not to be confused with the veritable Probability Density Function.

---

## Math 225- HW 11 Due: Dec 9 by Midnight

---

Submit the first two problems, along with any three additional problems of your choice.

1. • Prove that if  $U$  and  $T$  simultaneously diagonalizable then  $U$  and  $T$  commute. i.e.  $UT = TU$

If  $U$  and  $T$  are simultaneously diagonalizable, this means that  $[T]_\beta$  and  $[U]_\beta$  are diagonal matrices. The product of two diagonal matrices is obviously commutative (i.e. if matrices  $X$  and  $Y$  are diagonal,  $XY = YX$ ). Thus, if  $U$  and  $T$  are simultaneously diagonalizable then:

$$\begin{aligned}[T]_\beta[U]_\beta &= [U]_\beta[T]_\beta \\ [TU]_\beta &= [UT]_\beta \\ TU &= UT\end{aligned}$$

- Conclude that if matrices  $A, B$  are simultaneously diagonalizable then  $A, B$  commute  
If  $A$  and  $B$  are simultaneously diagonalizable then we know that  $\exists Q$  invertible s.t.  $Q^{-1}AQ$  and  $Q^{-1}BQ$  are diagonal and so, using the logic that the product of two diagonal matrices is commutative, we have:

$$\begin{aligned}Q^{-1}AQQ^{-1}BQ &= Q^{-1}BQQ^{-1}AQ \\ Q^{-1}ABQ &= Q^{-1}BAQ \\ AB &= BA\end{aligned}$$

- Let  $T$  be diagonalizable linear operator on a finite dimensional vector space, then  $T$  and  $T^m$  are simultaneously diagonalizable for any  $m$  positive integer.

Because  $T$  is diagonalizable,  $[T]_\beta$  is a diagonal matrix. Thus,  $[T^m]_\beta = \underbrace{[T]_\beta \dots [T]_\beta}_{m \text{ times}} = \underbrace{[T]_\beta \dots [T]_\beta}_{m \text{ times}} = \prod_{i=1}^m [T]_\beta$ . Because  $[T]_\beta$  is diagonal,  $[T^m]_\beta = \prod_{i=1}^m [T]_\beta$  is also diagonal  $\Rightarrow T$  and  $T^m$  are simultaneously diagonalizable.

2. a) For any vector  $w \in E_\lambda$ ,  $T(w) = \lambda w \in E_\lambda$ . Let us define  $u = \lambda w$ . Because  $T(u) = T(\lambda w) = \lambda T(w) = \lambda^2 w = \lambda u$ ,  $u = \lambda w \in E_\lambda$ . Thus we have shown  $\forall w \in E_\lambda, T(w) \in E_\lambda \Rightarrow E_\lambda$  is a  $T$ -invariant subspace of  $V$ .
- b) Let us define this  $T$ -cyclic subspace generated by  $v$  as  $W \leq V$ .  $W$  can be expressed as  $\text{Span}\{v, T(v), \dots, T^n(v)\}$ . We now show that  $\forall w \in W, T(w) \in W$ . By definition,  $\forall w \in W$  can be expressed as  $w = \sum_{i=0}^n c_i T^i(v)$  and so  $T(w) = c_n T^{n+1}(v) + \sum_{i=1}^n c_i T^i(v)$ . Because  $T^{n+1}(v)$  can be expressed as a linear combination of  $\{v, T(v), \dots, T^n(v)\}$ , this means that  $T(w)$  can be expressed as a linear combination of  $\{v, T(v), \dots, T^n(v)\} \Rightarrow T(w) \in \text{Span}\{v, T(v), \dots, T^n(v)\} \Rightarrow T(w) \in W$ . Thus, we have shown  $\forall w \in W, T(w) \in W \Rightarrow W$  is a  $T$ -invariant subspace of  $V$ .
- c) The  $T$ -cyclic subspace  $W$  can be given by  $W = \text{Span}\{v, T(v), \dots, T^n(v)\}$ . We now prove both directions of this statement:

1. If  $w \in W$ ,  $w = g(T)v$

If  $w \in W$ ,  $w$  can be expressed as  $\sum_{i=0}^n c_i T^i(v) = U(v)$ , where  $U = \sum_{i=0}^n c_i T^i$  is an operator. Defining  $g(x) = \sum_{i=0}^n c_i x^i$ ,  $U = g(T)$  and so we have that  $w = g(T)v$ .

2. If  $w = g(T)v$ ,  $w \in W$

We can express polynomial  $g$  as  $g(x) = \sum_{i=0}^n c_i x^i$ . Thus, we have that  $w = g(T)v = \sum_{i=0}^n c_i T^i(v) \Rightarrow w \in \text{Span}\{v, T(v), \dots, T^n(v)\} \Rightarrow w \in W$ .

d) Because  $V$  is a  $T$ -cyclic subspace of itself, we can express  $V = \text{Span}(\{v, T(v), T^2(v), \dots, T^n(v)\})$ . Thus, this means  $\forall z \in V$ ,  $z = \sum_{i=0}^n c_i T^i(v) \Rightarrow$  because  $U(v) \in V$ ,  $U(v)$  can be expressed as a linear combination of  $T^i(v)$ . Note that if  $U$  commutes with  $T$ , that means  $UT^2 = UTT = TUT = TTU = T^2U$ , or more generally  $UT^\alpha = T^\alpha U$  for  $\alpha \geq 0$ . Thus, we have that for  $i \geq 0$ :

$$\begin{aligned} UT^i &= T^i U \\ UT^i(v) &= T^i U(v) \\ U(T^i(v)) &= T^i(\sum_{k=0}^n c_k T^k(v)) \\ U(T^i(v)) &= \sum_{k=0}^n c_k T^{i+k}(v) \end{aligned}$$

Setting  $a = T^i(v)$ , we have:

$$U(a) = \sum_{k=0}^n c_k T^k(a)$$

Thus, we can clearly see that  $U = g(T)$ , where polynomial  $g$  is given by  $g(x) = \sum_{k=0}^n c_k x^k$ .

e) There are two cases in this scenario: (1) all vectors in  $V$  are eigenvectors or (2) not all vectors in  $V$  are eigenvectors. We address both cases below:

1. All vectors in  $V$  are eigenvectors

This means that  $\forall v \in V, T(v) = \lambda v$  where  $\lambda \in \mathbb{F}$ . Let us define two vectors  $a, b \in V$  and compute  $T(a+b)$ :

$$T(a+b) = \lambda_{a+b}(a+b)$$

However, by linearity, we also have that  $T(a+b) = T(a) + T(b) = \lambda_a a + \lambda_b b$ . Thus, we have that:

$$\begin{aligned} T(a+b) &= T(a+b) \\ \lambda_{a+b}(a+b) &= \lambda_a a + \lambda_b b \end{aligned}$$

This means that  $\lambda_{a+b} = \lambda_a = \lambda_b \Rightarrow \forall v \in V, T(v) = \lambda_a v \Rightarrow T = cI$  where  $c \in \mathbb{F}$ .

2. Not all vectors in  $V$  are eigenvectors

This means that  $\exists v \neq 0 \in V$  s.t.  $T(v) \neq \lambda v$ ,  $\forall \lambda \in \mathbb{F}$ . Consider the set  $\{v, T(v)\}$ . In order for the set of vectors  $\{a, b\}$  to be linearly independent, neither  $a$  nor  $b$  can be expressed as a scalar multiple of the either vector. Because we know that  $\forall \lambda \in \mathbb{F}$ ,  $T(v) \neq \lambda v$ ,  $\{v, T(v)\}$  are a linearly independent set of two vectors  $\Rightarrow$  because  $\dim(V) = 2$ ,  $\{v, T(v)\}$  serve as a basis for  $V \Rightarrow V = \text{Span}(\{v, T(v)\}) \Rightarrow V$  is a  $T$ -cyclic subspace of itself.

3. I didn't do this question.

4. (a) We use induction to prove this statement.

1. Base Case: Single element  $v_1$

If  $n = 1$ , then given  $\sum_{i=1}^n v_i \in W \Rightarrow v_1 \in W$ .

2. Inductive Step: Given  $v_1, \dots, v_{k-1} \in W$ , prove that  $v_k \in W$   
 For proof by contrapositive, let us assume that  $v_k \notin W$ . Let us define  $v = v_1 + \dots + v_n$ . We start with our given:

$$\begin{aligned} v &= v_1 + \dots + v_n \in W \\ v &= (v_1 + \dots + v_{k-1}) + v_k + (v_k + \dots + v_n) \\ v_1 + \dots + v_{k-1} &= v - v_k - (v_k + \dots + v_n) \end{aligned}$$

Because  $W$  is a subspace, it is closed under addition. Thus, because  $v_k \notin W \Rightarrow v - v_k - (v_k + \dots + v_n) \notin W \Rightarrow v_1 + \dots + v_{k-1} \notin W$ . By proof by contrapositive, we have proven if  $v_1, \dots, v_{k-1}$ , then  $\in W, v_k \in W$ .

- (b) Let us define  $U$  as a non-trivial  $T$ -invariant subspace of  $V$ . If  $T$  is a diagonalizable linear operator, that means its eigenvectors  $v_1, v_2, \dots, v_n$  form a basis for  $V$ . Because  $U$  is a non-trivial subspace,  $\exists v \neq 0 \in U$ . Furthermore, given that  $\forall v \neq 0 \in U \leq V$ ,  $v$  can be written as a linear combination of  $\{v_1, \dots, v_n\}$ , we can define the nonempty set of eigenvectors which all elements of  $U$  are a linear combination of as  $\{u_1, \dots, u_k\} \Rightarrow \text{Span}(\{u_1, \dots, u_k\}) = U$ . Note that  $\{u_1, \dots, u_k\}$  are all part of the basis  $\{v_1, \dots, v_n\}$  for  $V$  and so they are all linearly independent. Thus, we can conclude the linearly independent and generating set of eigenvectors  $\{u_1, \dots, u_k\}$  forms a basis for  $U$  and so  $T|_U$  is diagonalizable.
- (c) Because  $v_1, v_2, \dots, v_n \in V$  all correspond to distinct eigenvalues, they are all linearly independent. Given these  $n$  linearly independent vectors and that  $\dim(V) = n$ , we can conclude that the eigenvectors  $v_1, v_2, \dots, v_n$  form a basis for  $V$ . This means that  $V = \text{Span}(\{v_1, v_2, \dots, v_n\})$ . Let us define vector  $v = v_1 + v_2 + \dots + v_n$ . Note that  $\text{Span}(\{v, T(v), \dots, T^n(v)\}) = \text{Span}(\{\sum_{i=1}^n v_i, \sum_{i=1}^n \lambda_i v_i, \dots, \sum_{i=1}^n \lambda_i^n v_i\})$ . We can write out this transformation from the eigenvectors  $v_1, \dots, v_n$  to  $\sum_{i=1}^n v_i, \sum_{i=1}^n \lambda_i v_i, \dots, \sum_{i=1}^n \lambda_i^n v_i$  as such:

$$\begin{bmatrix} 1 & 1 & \dots & 1 \\ \lambda_1 & \lambda_2 & \dots & \lambda_n \\ \vdots & \ddots & \ddots & \vdots \\ \lambda_1^n & \lambda_2^n & \dots & \lambda_n^n \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} = \begin{bmatrix} \sum_{i=1}^n v_i \\ \sum_{i=1}^n \lambda_i v_i \\ \vdots \\ \sum_{i=1}^n \lambda_i^n v_i \end{bmatrix}$$

Note that the leftmost matrix above, which I refer to as  $V$ , is the Vandermonde matrix (pg 230.) Because all  $\forall 0 \leq i < j \leq n, \lambda_i \neq \lambda_j$ ,  $\det(V) = \prod_{0 \leq i < j \leq n} (\lambda_j - \lambda_i) \neq 0 \Rightarrow V$  is invertible  $\Rightarrow$  because  $\{v_1, v_2, \dots, v_n\}$  serve as a basis for  $V$ , so does  $\{\sum_{i=1}^n v_i, \sum_{i=1}^n \lambda_i v_i, \dots, \sum_{i=1}^n \lambda_i^n v_i\} \Rightarrow V = \text{Span}(\{v, T(v), \dots, T^n(v)\}) \Rightarrow V$  is a  $T$ -cyclic subspace of itself.

5. (a) We prove both directions of this statement below:

1. If  $T$  is diagonalizable,  $V$  is the direct sum of one-dimensional  $T$ -invariant subspaces

If  $T$  is diagonalizable, that means that eigenvectors  $v_1, v_2, \dots, v_n \in V$  serve as a basis for  $V$ . This means  $V = \text{Span}(\{v_1, \dots, v_n\}) = \{\sum_{i=1}^n c_i v_i : c_i \in \mathbb{R}\} = \{\sum_{i=1}^n T(\frac{c_i}{\lambda_i} v_i)\}$ . Let us define the set  $W_j = \{cv_j : c \in \mathbb{F}\}$  for a given eigenvector  $v_j$ . Note that  $W_j$  is a one-dimensional subspace as  $W_j$  is composed of scalar multiples of one unique vector,  $v_j$ . Furthermore,  $W_j$  is a  $T$ -invariant subspace

as  $\forall w \in W_j, T(w)$  is equal to a scalar multiple of  $v_j \Rightarrow w \in W_j$ . Furthermore, because  $\{v_1, \dots, v_n\}$ , serve as a basis, that means that all the eigenvectors are linearly independent  $\Rightarrow$  for  $0 \leq i < j \leq n$ ,  $W_i \cap W_j = \emptyset$  and so we have that  $V = \{\sum_{i=1}^n c_i v_i : c_i \in \mathbb{R}\} = W_1 \oplus W_2 \cdots \oplus W_n$ .

2. If  $V$  is the direct sum of one-dimensional  $T$ -invariant subspaces,  $T$  is diagonalizable

Let us define  $V = W_1 \oplus W_2 \cdots \oplus W_k$  and the one basis vector for the  $j$ th subspace  $W_j$  as  $v_j$ . Because  $W_j$  is one-dimensional and a subspace (thus closed under addition and scalar multiplication),  $W_j = \{cv_j : c \in \mathbb{F}\}$ . Furthermore, because  $W_j$  is  $T$ -invariant and  $v_j \in W_j$ ,  $T(v_j) \in W_j \Rightarrow T(v_j) \in \{cv_j : c \in \mathbb{F}\} \Rightarrow v_j$  is an eigenvector of  $T_{W_j}$  as  $T(v_j) = kv_j$  where  $k \in \mathbb{F}$ . Because  $V$  is a *direct* sum of  $W_1, \dots, W_k$ , the individual basis vector  $v_j$  for each subspace is linearly independent from all of the vectors in  $\{v_1, \dots, v_{j-1}\} \cup \{v_{j+1}, \dots, v_k\} \Rightarrow \{v_1, \dots, v_k\}$  are linearly independent<sup>1</sup>. Furthermore,  $V = W_1 \oplus W_2 \cdots \oplus W_k$  means that  $V$  contains all possible linear combinations of  $\{v_1, \dots, v_k\} \Rightarrow V = \text{Span}(\{v_1, \dots, v_k\})$ . Thus we can conclude that the linearly independent and generating eigenvectors  $\{v_1, \dots, v_k\}$  forms a basis for  $V$  and so  $T$  is diagonalizable.

- b) Let us define the unordered basis for the  $T$ -invariant subspace  $W_j$  as  $\beta_j$ . This means that the ordered basis  $\gamma$  for vector space  $V$  can be given as  $\gamma = \beta_1 \cup \beta_2 \cdots \cup \beta_k$ . We now try to understand what the matrix  $[T]_\beta$  looks like. Note that  $\forall v \in \beta_j, v \in W_j$  and so  $T(v) = T_{W_j}(v) \in W_j \Rightarrow T(v)$  can be expressed as a linear combination of  $\beta_j$ . Thus,  $[T]_\beta$  will be given as a collection of block matrices  $[T_{W_j}]_{\beta_j}$  along the diagonal:

$$[T]_\beta = \begin{bmatrix} [T_{W_1}]_{\beta_1} & 0 & \cdots & 0 \\ 0 & [T_{W_2}]_{\beta_2} & \cdots & 0 \\ 0 & \ddots & \ddots & 0 \\ 0 & 0 & 0 & [T_{W_k}]_{\beta_k} \end{bmatrix}$$

From this matrix, it is obvious that:

$$\begin{aligned} \det(T) &= \det([T]_\beta) = \prod_{i=1}^k \det([T_{W_i}]_{\beta_i}) = \prod_{i=1}^k \det(T_{W_i}) \\ \det(T) &= \prod_{i=1}^k \det(T_{W_i}) \end{aligned}$$

6. To prove this law, we compare the LHS with the RHS. The LHS can be given as:

$$\begin{aligned} \|x + y\|^2 + \|x - y\|^2 &= \langle x + y, x + y \rangle + \langle x - y, x - y \rangle = \langle x, x + y \rangle + \langle y, x + y \rangle + \langle x, x - y \rangle - \langle y, x - y \rangle \\ &= \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle + \langle x, x \rangle - \langle x, y \rangle - \langle y, x \rangle + \langle y, y \rangle = 2\langle x, x \rangle + 2\langle y, y \rangle \end{aligned}$$

We now compare this with the RHS:

---

<sup>1</sup>This can be trivially proven by induction by the following proof:  $\{v_1\}$  is a linearly independent set,  $\{v_1, v_2\}$  is a linearly independent set,  $\{v_1, v_2, v_3\}$  is a linearly independent set, and so on until  $\{v_1, \dots, v_n\}$  is a linearly independent set. I believe we did this proof in class.

$$2\|x\|^2 + 2\|y\|^2 = 2\langle x, x \rangle + 2\langle y, y \rangle$$

Thus, we can clearly see that the LHS = RHS and so we have proven this law.