

PSETs Landing Page*

Anish Krishna Lakkapragada

This is the documentation for using my PSET PDFs responsibly. I post these LaTeX'd PSETs (1) as an education resource for friends at other universities, fellow Yalies, and all those interested and (2) for quick reference. These PSETs are not to be used irresponsibly; only look at the solution after giving each problem an honest attempt. **If YOU USE THESE PSETS TO CHEAT, YOU ARE NOT ONLY STUPID BUT YOU ARE CHEATING YOURSELF OUT OF THE ABILITY TO FALL IN LOVE WITH MATH.** Furthermore, I am not smarter than you and my solutions did not always get a perfect score.

The general format for accessing the (one-indexed) `N`th assigned PSET PDF of a Yale course with course number `CODE` is:

`https://anish.lakkapragada.com/notes/TYPE-CODE/psets/N.pdf`

where `TYPE` is `stats` or `math`. Similarly, to access my solution for this PSET you can go to:

`https://anish.lakkapragada.com/notes/TYPE-CODE/sols/N.pdf`

These PSETs and associated solution PDFs are synchronized daily at 4:20AM with my computer files through a Cronjob Shell Script. If you want to contribute any corrections, please email `anish.lakkapragada@yale.edu`.

*Note that PDF here is referring to Portable Document Format, not to be confused with the veritable Probability Density Function.

STATS 242 HW 6

March 5, 2025

Number of late days: 0; Collaborators: Matt Sprintson

1.

We first compute the first moment μ_1 of $X \sim \text{Geom}(p)$, which is given by $\mu_1 = \mathbb{E}[X] = \frac{1}{p}$. We now solve for the MoM estimator \hat{p} where the observed sample moment $\hat{\mu}_1 = \frac{1}{n}(X_1 + \dots + X_n) = \bar{X}$ is equal to theoretical moment μ_1 with the estimated \hat{p} parameter:

$$\begin{aligned}\hat{\mu}_1 = \bar{X} &= \frac{1}{\hat{p}} \\ \hat{p} &= \frac{1}{\bar{X}}\end{aligned}$$

Thus, $\hat{p} = \frac{1}{\bar{X}}$ is the MoM estimator \hat{p} for p . We now compute the MLE of p . We first start by finding the log-likelihood function $\ell_n(p)$:

$$\begin{aligned}\ell_n(p) = \log[\text{lik}(p)] &= \sum_{i=1}^n \log f(X_i|p) = \sum_{i=1}^n \log(p(1-p)^{x_i-1}) = \sum_{i=1}^n \log(p) + (x_i - 1)\log(1-p) \\ &= n\log(p) + \sum_{i=1}^n (x_i - 1)\log(1-p)\end{aligned}$$

We now solve for the MLE \hat{p} by solving $\ell'_n(\hat{p}) = 0$:

$$\begin{aligned}
\ell'_n(\hat{p}) &= 0 \\
(n \log(\hat{p}) + \sum_{i=1}^n (x_i - 1) \log(1 - \hat{p}))' &= 0 \\
\frac{n}{\hat{p}} - \sum_{i=1}^n \frac{(x_i - 1)}{1 - \hat{p}} &= 0 \\
\frac{n}{\hat{p}} - \frac{1}{1 - \hat{p}} (n\bar{X} - n) &= 0 \\
\frac{n(1 - \hat{p}) - \hat{p}(n\bar{X} - n)}{\hat{p}(1 - \hat{p})} &= 0 \\
n(1 - \hat{p}) - \hat{p}(n\bar{X} - n) &= 0 \\
n - n\hat{p} - \hat{p}n\bar{X} + n\hat{p} &= 0 \\
n &= n\hat{p}\bar{X} \\
1 &= \hat{p}\bar{X} \\
\hat{p} &= \frac{1}{\bar{X}}
\end{aligned}$$

Thus the MLE of p is $\hat{p} = \frac{1}{\bar{X}}$. We now find the sampling distribution of the MLE of p , under large n . We begin by computing the Fisher Information $I(p)$:

$$\begin{aligned}
I(p) &= -\mathbb{E}_p \left[\frac{\partial^2}{\partial p^2} \log f(X|p) \right] = -\mathbb{E}_p \left[\frac{\partial^2}{\partial p^2} (\log(p) + (x-1)\log(1-p)) \right] \\
&= -\mathbb{E}_p \left[\frac{\partial}{\partial p} \left(\frac{1}{p} + (1-x)\frac{1}{1-p} \right) \right] = -\mathbb{E}_p \left[-\frac{1}{p^2} + \frac{(1-x)}{(1-p)^2} \right] = -\left(-\frac{1}{p^2} + \frac{1}{(1-p)^2} - \frac{\mathbb{E}_p[x]}{(1-p)^2} \right) \\
&= \frac{1}{p^2} - \frac{1}{(1-p)^2} + \frac{1}{p(1-p)^2} = \frac{(1-p)^2 - p^2 + p}{p^2(1-p)^2} = \frac{1-p}{p^2(1-p)^2} = \frac{1}{p^2(1-p)}
\end{aligned}$$

So this means that for the MLE \hat{p} , $\sqrt{n}(\hat{p} - p) \rightarrow \mathcal{N}(0, \frac{1}{I(p)})$ or $\sqrt{n}(\hat{p} - p) \rightarrow \mathcal{N}(0, p^2(1-p))$ meaning $\hat{p} \rightarrow \mathcal{N}(p, \frac{p^2(1-p)}{n})$ as $n \rightarrow \infty$.

2.

We first compute the MoM estimate of p . To do so, we compute the first moment μ_1 of $X \sim \text{NegBinom}(r, p)$ which is given by $\mu_1 = \mathbb{E}[X] = \frac{pr}{1-p}$. We now solve for the MoM estimator \hat{p} where the observed sample moment $\hat{\mu}_1 = \frac{1}{n}(X_1 + \dots + X_n) = \bar{X}$ is equal to theoretical moment μ_1 with the estimated \hat{p} parameter:

$$\begin{aligned}
\hat{\mu}_1 &= \bar{X} = \frac{\hat{p}r}{1 - \hat{p}} \\
\bar{X}(1 - \hat{p}) &= \hat{p}r \\
\bar{X} &= \hat{p}\bar{X} + \hat{p}r \\
\bar{X} &= \hat{p}(\bar{X} + r) \\
\hat{p} &= \frac{\bar{X}}{\bar{X} + r}
\end{aligned}$$

Thus the MoM estimator of p is given by $\hat{p} = \frac{\bar{X}}{\bar{X} + r}$. We now compute the MLE of p . We first start by finding the log-likelihood function $\ell_n(p)$:

$$\begin{aligned}
\ell_n(p) &= \log[\text{lik}(p)] = \sum_{i=1}^n \log f(x_i|p) = \sum_{i=1}^n \log \left(\binom{x_i + r - 1}{x_i} (1-p)^r p^{x_i} \right) \\
&= \sum_{i=1}^n \log \left(\binom{x_i + r - 1}{x_i} \right) + r \log(1-p) + x_i \log(p) = nr \log(1-p) + \sum_{i=1}^n \log \left(\binom{x_i + r - 1}{x_i} \right) + x_i \log(p)
\end{aligned}$$

We now solve for the MLE \hat{p} by solving $\ell'_n(\hat{p}) = 0$:

$$\begin{aligned}
\ell'_n(\hat{p}) &= 0 \\
(nr \log(1 - \hat{p}) + \sum_{i=1}^n \log \left(\binom{x_i + r - 1}{x_i} \right) + x_i \log(\hat{p}))' &= 0 \\
\frac{-nr}{1 - \hat{p}} + \sum_{i=1}^n \frac{x_i}{\hat{p}} &= 0 \\
-\frac{nr\hat{p}}{\hat{p}(1 - \hat{p})} + \frac{n\bar{X}(1 - \hat{p})}{\hat{p}(1 - \hat{p})} &= 0 \\
\frac{n\bar{X}(1 - \hat{p}) - nr\hat{p}}{\hat{p}(1 - \hat{p})} &= 0 \\
n\bar{X}(1 - \hat{p}) - nr\hat{p} &= 0 \\
\bar{X}(1 - \hat{p}) - r\hat{p} &= 0 \\
\bar{X} - \hat{p}\bar{X} - \hat{p}r &= 0 \\
\bar{X} &= \hat{p}(\bar{X} + r) \\
\hat{p} &= \frac{\bar{X}}{\bar{X} + r}
\end{aligned}$$

Thus the MLE of p is given by $\hat{p} = \frac{\bar{X}}{\bar{X} + r}$. We now find the sampling distribution of the MLE of p , under large n . We begin by computing the Fisher Information $I(p)$:

$$\begin{aligned}
I(p) &= \text{Var}_p\left[\frac{\partial}{\partial p} \log f(x|p)\right] = \text{Var}_p\left[\frac{\partial}{\partial p} \left[\log\left(\binom{x+r-1}{x}\right) + r\log(1-p) + x\log(p)\right]\right] \\
&= \text{Var}_p\left[\frac{-r}{1-p} + \frac{x}{p}\right] = \frac{1}{p^2} \text{Var}_p[x] = \frac{r}{p(1-p)^2}
\end{aligned}$$

So this means that for the MLE \hat{p} , $\sqrt{n}(\hat{p} - p) \rightarrow \mathcal{N}(0, \frac{1}{I(p)})$ or $\sqrt{n}(\hat{p} - p) \rightarrow \mathcal{N}(0, \frac{p(1-p)^2}{r})$ so $\hat{p} \rightarrow \mathcal{N}(p, \frac{p(1-p)^2}{nr})$.

3.

- (a) The PDF of $\text{Pareto}(\alpha, x_m)$ is given by $f(x|\alpha, x_m) = \frac{\alpha x_m^\alpha}{x^{\alpha+1}}$. Thus, the PDF of $\text{Pareto}(\theta, 1)$ is given by:

$$f(x|\theta, 1) = \frac{\theta \cdot 1^\theta}{x^{\theta+1}} = \frac{\theta}{x^{\theta+1}}$$

Expressed differently, we can set functions $T(x) = \log(\frac{1}{x})$, $A(\theta) = \log(\frac{1}{\theta})$, $h(x) = \frac{1}{x}$ so the PDF

$$f(x|\theta) = e^{\theta T(x) - A(\theta)} h(x) = \frac{e^{\theta T(x)}}{x e^{A(\theta)}} = \frac{(e^{\log(\frac{1}{x})})^\theta}{x e^{\log \frac{1}{\theta}}} = \frac{1}{x^\theta \cdot x \cdot \frac{1}{\theta}} = \frac{\theta}{x^{\theta+1}}$$

is identical to the aforementioned derived $\text{Pareto}(\theta, 1)$ PDF.

- (b) We differentiate both sides of the identity $1 = \int_{\mathcal{X}} f(x|\theta) dx$ to compute $\mathbb{E}_\theta[T(X)]$:

$$\begin{aligned}
&\int_{\mathcal{X}} f(x|\theta) dx = 1 \\
\frac{d}{d\theta} \int_{\mathcal{X}} f(x|\theta) dx &= \int_{\mathcal{X}} \frac{d}{d\theta} f(x|\theta) dx = \frac{d}{d\theta}(1) = 0 \\
&\int_{\mathcal{X}} \frac{d}{d\theta} [e^{\theta T(x) - A(\theta)} h(x)] dx = 0 \\
\int_{\mathcal{X}} [(e^{\theta T(x) - A(\theta)})' h(x) + h'(x) e^{\theta T(x) - A(\theta)}] dx &= \int_{\mathcal{X}} [(e^{\theta T(x) - A(\theta)})(T(x) - A'(\theta)) h(x) + 0] dx = 0 \\
&\int_{\mathcal{X}} [e^{\theta T(x) - A(\theta)} h(x)] (T(x) - A'(\theta)) dx = 0 \\
\int_{\mathcal{X}} f(x|\theta) (T(x) - A'(\theta)) dx &= \mathbb{E}_\theta[T(x) - A'(\theta)] = 0 \\
\mathbb{E}_\theta[T(x)] &= \mathbb{E}_\theta[A'(\theta)] = A'(\theta)
\end{aligned}$$

So we have $\mathbb{E}_\theta[T(x)] = A'(\theta)$. We now verify this is the case for our Pareto($\theta, 1$) model with its defined functions $T(x) = \log(\frac{1}{x})$, $A(\theta) = \log(\frac{1}{\theta})$, $h(x) = \frac{1}{x}$ in part (a). We first compute $\mathbb{E}_\theta[T(x)]$:

$$\mathbb{E}_\theta[T(x)] = \mathbb{E}_\theta[\log(\frac{1}{x})] = \int_{-\infty}^{\infty} f(x|\theta) \log(\frac{1}{x}) dx = \int_1^{\infty} \frac{\theta}{x^{\theta+1}} \log(\frac{1}{x}) dx$$

We can use a u-substitution of $u = \frac{1}{x}$ so $-u^{\theta-1} du = -\frac{x}{x^\theta} \cdot -\frac{1}{x^2} dx = \frac{1}{x^{\theta+1}} dx$. Thus, this integral can be re-written as:

$$\int_1^0 \log(u) \cdot \theta \cdot (-u^{\theta-1}) du = \theta \int_0^1 \log(u) \cdot u^{\theta-1} du$$

Using integration by parts, we can set $a = \log(u)$, $da = \frac{1}{u} du$, $db = u^{\theta-1} du$, $b = \frac{1}{\theta} u^\theta$. As such, we get:

$$\begin{aligned} \theta \int_0^1 \log(u) \cdot u^{\theta-1} du &= \theta [ab \Big|_{u=0}^{u=1} - \int_0^1 b da] \\ &= \theta \left[\frac{\log(u) \cdot u^\theta}{\theta} \right] \Big|_0^1 - \theta \int_0^1 \frac{1}{\theta} u^{\theta-1} du = 0 - \int_0^1 u^{\theta-1} du = -\frac{1}{\theta} u^\theta \Big|_0^1 = -\frac{1}{\theta} + 0 = -\frac{1}{\theta} \end{aligned}$$

So $\mathbb{E}_\theta[T(x)] = -\frac{1}{\theta}$. We now compute $A'(\theta)$:

$$A'(\theta) = \frac{1}{\frac{1}{\theta}} \cdot \frac{-1}{\theta^2} = \frac{-\theta}{\theta^2} = -\frac{1}{\theta}$$

So because $\mathbb{E}_\theta[T(x)] = -\frac{1}{\theta} = A'(\theta)$, we have verified for our Pareto model in part (a) that $\mathbb{E}_\theta[T(x)] = A'(\theta)$.

(c) We first find the log-likelihood function $\ell_n(\theta)$:

$$\begin{aligned} \ell_n(\theta) &= \log[\text{lik}(\theta)] = \sum_{i=1}^n \log f(x_i|\theta) = \sum_{i=1}^n \log[e^{\theta T(x_i) - A(\theta)} h(x_i)] = \\ &= \sum_{i=1}^n \log[e^{\theta T(x_i)}] - \log[e^{A(\theta)}] + \log[h(x_i)] = -nA(\theta) + \sum_{i=1}^n \theta T(x_i) + \log[h(x_i)] \end{aligned}$$

To get the MLE of θ we solve for the solution to $0 = \ell'_n(\theta)$:

$$\begin{aligned} \ell'_n(\theta) &= 0 \\ (-nA(\theta) + \sum_{i=1}^n \theta T(x_i) + \log[h(x_i)])' &= 0 \\ -nA'(\theta) + \sum_{i=1}^n T(x_i) &= 0 \\ nA'(\theta) &= \sum_{i=1}^n T(x_i) \\ A'(\theta) &= \frac{1}{n} \sum_{i=1}^n T(x_i) \\ \mathbb{E}_\theta[T(x)] = A'(\theta) &= \frac{1}{n} \sum_{i=1}^n T(x_i) \end{aligned}$$

So therefore the MLE of θ is the unique solution to $\mathbb{E}_\theta[T(x)] = \frac{1}{n}\sum_{i=1}^n T(x_i)$. Because the generalized MoM estimator of θ is also the solution to $\mathbb{E}_\theta[T(x)] = \frac{1}{n}\sum_{i=1}^n T(x_i)$, and we are guaranteed this solution is unique, the generalized MoM estimator of θ is equivalent to the MLE of θ .

An example of this occurring would be the generalized MoM estimator based on $T(x) = \log(x)$ for the $\text{Pareto}(\theta, 1)$ model coinciding with the MLE.

4.

- (a) By the CLT, $\sqrt{n}(\hat{p} - p) \rightarrow \mathcal{N}(0, p(1 - p))$. Informally, this gives us that $\text{Var}[\sqrt{n}(\hat{p} - p)] \approx \hat{p}(1 - \hat{p}) \implies n\text{Var}(\hat{p} - p) \approx \hat{p}(1 - \hat{p}) \implies \text{Var}(p) \approx \frac{\hat{p}(1 - \hat{p})}{n}$. Using this approximation for the variance of p , we can compute a 95% confidence interval for p as $\hat{p} \pm z^{\frac{0.05}{2}} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$ where $z^{\frac{0.05}{2}}$ is the upper- $\frac{0.05}{2}$ point of the standard normal distribution.
- (b) We solve $\sqrt{n}(\hat{p} - p) = \pm \sqrt{p(1 - p)}z^{(\alpha/2)}$ below to create a confidence interval for p :

$$\begin{aligned}\sqrt{n}(\hat{p} - p) &= \pm \sqrt{p(1 - p)}z^{(\alpha/2)} \\ n(\hat{p} - p)^2 &= p(1 - p)(z^{(\alpha/2)})^2 \\ n\hat{p}^2 - 2n\hat{p}p + np^2 &= (z^{(\alpha/2)})^2 p - (z^{(\alpha/2)})^2 p^2 \\ (n + (z^{(\alpha/2)})^2)p^2 - (2n\hat{p} + (z^{(\alpha/2)})^2)p + n\hat{p}^2 &= 0\end{aligned}$$

Using the quadratic formula, we construct the following confidence interval for p :

$$p = \frac{(2n\hat{p} + (z^{(\alpha/2)})^2) \pm \sqrt{(2n\hat{p} + (z^{(\alpha/2)})^2)^2 - 4n\hat{p}^2(n + (z^{(\alpha/2)})^2)}}{2(n + (z^{(\alpha/2)})^2)}$$

- (c) For our confidence interval in part (a), we report the simulated coverage probabilities below:

	$n = 10$	$n = 40$	$n = 100$
$p = 0.1$	0.65061	0.91435	0.93267
$p = 0.3$	0.83926	0.92955	0.94984
$p = 0.5$	0.8922	0.91919	0.94205

We similarly report the simulated coverage probabilities for our confidence interval in part (b):

	$n = 10$	$n = 40$	$n = 100$
$p = 0.1$	0.93053	0.94406	0.9375
$p = 0.3$	0.92463	0.94432	0.93659
$p = 0.5$	0.97861	0.961	0.94316

We can clearly see from comparing these two tables that our interval construction in part (b) leads to generally much higher simulated coverage probabilities.

```

1      # %%
2  import numpy as np
3  from scipy.stats import norm
4

```



```

5 z_alpha = np.abs(norm.ppf(0.05/2))
6 z_alpha_squared = z_alpha ** 2
7
8 NUM_SIMULATIONS = 100 * 1000
9 SAMPLE_SIZES = [10, 40, 100]
10 TRUE_PARAMS = [0.1, 0.3, 0.5]
11
12 def generate_data(n,p):
13     return np.random.binomial(n=n, p=p) # number of successes
14
15 def part_a_interval(num_successes, n):
16     p_hat = num_successes / n
17     delta = z_alpha * np.sqrt(p_hat * (1 - p_hat) / n)
18     return p_hat - delta, p_hat + delta
19
20 def part_b_interval(num_successes, n):
21     p_hat = num_successes / n
22     denominator = 2 * (n + z_alpha_squared)
23     center = (2 * n * p_hat + z_alpha_squared) / denominator
24     delta = np.sqrt((2 * n * p_hat + z_alpha_squared) ** 2 - 4 * n *
25     ↪ (p_hat ** 2) * (n + z_alpha_squared))
26     delta /= denominator
27     return center - delta, center + delta
28
29 # %%
30
31 for n in SAMPLE_SIZES:
32     for p in TRUE_PARAMS:
33         NUM_COVERAGE_A = 0
34         for _ in range(NUM_SIMULATIONS):
35             n_successes = generate_data(n, p)
36             lower, upper = part_a_interval(n_successes, n)
37
38             if p >= lower and p <= upper: NUM_COVERAGE_A += 1
39
40     print(f"[PART (A)] For sample size n={n} and p={p}, simulated
41     ↪ coverage probability: {NUM_COVERAGE_A / NUM_SIMULATIONS}")
42
43
44 for n in SAMPLE_SIZES:
45     for p in TRUE_PARAMS:
46         NUM_COVERAGE_B = 0
47         for _ in range(NUM_SIMULATIONS):
48             n_successes = generate_data(n, p)
49             lower, upper = part_b_interval(n_successes, n)

```

```
50         if p >= lower and p <= upper: NUM_COVERAGE_B += 1
51
52
53     print(f"[PART (B)] For sample size n={n} and p={p}, simulated
54     ↪ coverage probability: {NUM_COVERAGE_B / NUM_SIMULATIONS}")
55
56 # %%
```