

# PSETs Landing Page\*

Anish Krishna Lakkapragada

This is the documentation for using my PSET PDFs responsibly. I post these LaTeX'd PSETs (1) as an education resource for friends at other universities, fellow Yalies, and all those interested and (2) for quick reference. These PSETs are not to be used irresponsibly; only look at the solution after giving each problem an honest attempt. **If YOU USE THESE PSETS TO CHEAT, YOU ARE NOT ONLY STUPID BUT YOU ARE CHEATING YOURSELF OUT OF THE ABILITY TO FALL IN LOVE WITH MATH.** Furthermore, I am not smarter than you and my solutions did not always get a perfect score.

The general format for accessing the (one-indexed) `N`th assigned PSET PDF of a Yale course with course number `CODE` is:

`https://anish.lakkapragada.com/notes/TYPE-CODE/psets/N.pdf`

where `TYPE` is `stats` or `math`. Similarly, to access my solution for this PSET you can go to:

`https://anish.lakkapragada.com/notes/TYPE-CODE/sols/N.pdf`

These PSETs and associated solution PDFs are synchronized daily at 4:20AM with my computer files through a Cronjob Shell Script. If you want to contribute any corrections, please email `anish.lakkapragada@yale.edu`.

---

\*Note that PDF here is referring to Portable Document Format, not to be confused with the veritable Probability Density Function.

---

## Math 226: HW 2, Due Sep 12 by Midnight

---

1. (9 points) Part *a*) requires familiarity with the vector space  $\mathcal{F}(S, \mathbb{F})$ . Check out Example 3 of Section 1.2 which is discussed in ULA section.

- a) (6 points) We define the set of  $L^2(\mathbb{R})$  functions as

$$L^2(\mathbb{R}) := \{f \in \mathcal{F}(\mathbb{R}, \mathbb{R}) : \int_{-\infty}^{\infty} |f(x)|^2 dx < \infty\}$$

Prove that the set of  $L^2(\mathbb{R})$  functions defined on the real line with the operations of addition, scalar multiplication (as in the Example 3) is a vector space.

- b) (3 points) (Problem 13 in the book) Let  $V := \{(a_1, a_2) : a_1, a_2 \in \mathbb{R}\}$ . If  $(a_1, a_2)$  and  $(b_1, b_2)$  are elements on  $V$  and  $c \in \mathbb{R}$ , define

$$(a_1, a_2) + (b_1, b_2) = (a_1 + b_1, a_2 + b_2) \quad c(a_1, a_2) = (ca_1, ca_2)$$

Is  $V$  a vector space over  $\mathbb{R}$  with these operations? Justify your answer.

### Practice Problems : Sec 1.2 : 10 - 22

2. (26 points)

- a) (7 points) (P.10, Sec 1.3) Prove that  $W_1 = \{(a_1, a_2, \dots, a_n) \in \mathbb{F}^n : a_1 + a_2 + \dots + a_n = 0\}$  is a subspace of  $\mathbb{F}^n$ , but  $W_2 = \{(a_1, a_2, \dots, a_n) \in \mathbb{F}^n : a_1 + a_2 + \dots + a_n = 1\}$  is not.

- **Remark:** This problem tells you that if  $V$  is any vector space, then any hyper-plane of  $V$  is a vector space if it passes through the origin. Take for example  $W_1$  as  $x + y + z = 0$  and  $W_2$  as  $x + y + z = 1$  as a subspace of  $\mathbb{R}^3$ .  $x + y + z = 0$ ,  $x + y + z = 1$  are the hyper-planes of  $\mathbb{R}^3$ . One is a subspace the other one is not. (Check out the Problems 11, 13, 14 in Section 1.3)

- b) (7 points) We define

$$\mathbb{Z}_2^n := \{v = (a_1, a_2, \dots, a_n) : a_i \in \mathbb{Z}_2\}$$

Recall that  $\mathbb{Z}_2^n$  defines a vector space over  $\mathbb{Z}_2$  (with addition and multiplication we discussed in class). Let  $E_n(v)$  define the number of nonzero components of  $v$ . For example if  $v = (1, 0, 1, 0, 0, 1)$  then  $E_6(v) = 3$ . Prove that the subset of  $\mathbb{Z}_2^n$  with even  $E_n$  defines a subspace of  $\mathbb{Z}_2^n$ .

Hint:  $E_n(u + v)$  can be formulated in terms of  $E_n(u)$ ,  $E_n(v)$ , and the common non-zero components of  $u, v$ .

- c) (6 points) Let  $W := \{f \in P_3(\mathbb{R}) : f(0) = f'(0), f(1) = 0\}$ . How does the elements of  $W$  look like? Show that  $W$  is a subspace of  $P_3(\mathbb{R})$ . Is it a subspace of  $P_4(\mathbb{R})$ ?
- d) (6 points) (P.18, Sec 1.3) Prove that a subset  $W$  of a vector space  $V$  is a subspace of  $V$  if and only if  $W \neq \emptyset$ , and whenever  $a \in \mathbb{F}$  and  $x, y \in W$ , then  $ax + y \in W$ .
3. (16 points) Let  $V$  be a vector space over a field  $\mathbb{F}$ , and suppose  $U \leq V$  and  $W \leq V$ . That is suppose that  $U$  and  $W$  are subspaces of  $V$ .
- a) (6 points) Prove that  $U \cap W \leq V$ .
- b) (6 points) We define  $U + W = \{u + w : u \in U, w \in W\}$ . Prove that  $U + W \leq V$ .

c) (4 points) Find two subspaces of  $\mathbb{R}^2$  whose union is *not* a subspace of  $\mathbb{R}^2$ . Make sure to validate your answer.

- **Remark:** This shows some important things. The intersection of subspaces is a subspace, but the union of subspaces (usually) isn't. The latter one is a subspace if and only if one of the subspaces is the subset of the other one. You can prove it as an exercise. ( It is problem 19 of Sec 1.2)

**Practice Problems :** Sec 1.3 : 1, 2, 3, 8, 9, 11,12, 13, 14, 17, 19

4. (13 points)

Let  $S := \{1 + x, 1 - x^2, x^3, x^4\}$  be a subset of  $P_4(\mathbb{R})$ .

- (2 points) How does the elements of  $\text{Span}(S)$  look like. Write down a clear definition for  $\text{Span}(S)$ .
- (5 points) Find a polynomial in  $P_4(\mathbb{R})$  that can not be written as a linear combination of the elements of  $S$ . Validate your answer. Deduce that  $S$  does not generate  $P_4(\mathbb{R})$ .
- (6 points) Prove that  $\text{Span}(S) = \{f \in P_4(\mathbb{R}) : 2f(0) = 2f'(0) - f''(0)\}$ . Hint: Let  $A, B$  are subsets of the same set. Then to show  $A = B$ , one needs to show  $A \subseteq B$ , and  $B \subseteq A$ .