PHARMACOVIGILANCE INTERNSHIP

ASSIGNMENT 01: ADR IDENTIFICATION BASICS

Submitted by: Anisha Hussain
Date: 06-11-2026

1. INTRODUCTION

Pharmacovigilance is the science and activities related to the detection, assessment, understanding and prevention of adverse effects or any other drug-related problems. An Adverse Drug Reaction (ADR) is a harmful and unintended response associated with the use of a medicinal product at normally used doses.

ADR identification is an important component of pharmacovigilance because it helps healthcare professionals recognise potential medication-related risks, improve patient safety and prevent further harm.

2. OBJECTIVES

The objectives of this assignment are:

- To understand the concept of Adverse Drug Reactions.
- To identify suspected ADRs from clinical case scenarios.
- To analyse patient symptoms and medication history.
- To establish the temporal relationship between drug administration and the onset of symptoms.
- To identify the suspected medication responsible for the reaction.
- To understand the importance of ADR monitoring and reporting.

3. BASIC CONCEPTS

a. Adverse Drug Reaction (ADR)

An ADR is a harmful and unintended response associated with a medicinal product when used at normally recommended doses.

b. Suspected Drug

A medication that may be responsible for producing an adverse reaction based on the available clinical information and timing.

c. Temporal Relationship

The time interval between the administration of a drug and the appearance of an adverse reaction. A reasonable temporal relationship supports a possible association.

d. Dechallenge

The observation of a suspected reaction after withdrawal of the suspected drug or reduction in its dose.

e. Rechallenge

The recurrence of an adverse reaction following the reintroduction of a suspected drug. Rechallenge should not be deliberately performed when it could expose a patient to harm.

4. METHODOLOGY

The following approach was used to analyse the clinical case scenarios:

- Review the patient's demographic and medical information.
- Identify all medications being taken by the patient.
- Identify the reported signs and symptoms.
- Establish the time interval between drug administration and symptom onset.
- Consider alternative explanations, including underlying diseases and concomitant medications.
- Identify the suspected medication.
- Document the findings and provide a preliminary conclusion.

5. CASE SCENARIO ANALYSIS

CASE STUDY 1: SKIN RASH ASSOCIATED WITH AMOXICILLIN

- **Patient Information**

- **Age:** 35 years

- **Gender:** Female

- **Medical condition:** Bacterial respiratory tract infection

- **Suspected medication:** Amoxicillin

- **Dose:** 500 mg three times daily (fictional case detail)

- **Route:** Oral

- **Clinical History:** A 35-year-old female patient was prescribed amoxicillin for a bacterial respiratory tract infection. After three days of treatment, the patient developed a skin rash accompanied by itching.
  The symptoms were not reported before starting the medication.

- **ADR Identification**

PARAMETER	FINDINGS
Suspected drug	Amoxicillin
Adverse reaction	Skin rash and itching
Time of onset	3 days after treatment initiation
Nature of reaction	Possible hypersensitivity reaction
Temporal relationship	Present
Alternative causes	Infection or other exposures need evaluation

- **Analysis**

The appearance of a skin rash and itching after the initiation of amoxicillin suggests a possible drug-related hypersensitivity reaction.

The temporal relationship between medication administration and symptom onset supports a suspected association. However, other possible causes should also be investigated.

- **Conclusion**

Amoxicillin is identified as the suspected drug associated with the reported skin rash and itching. The case represents a suspected ADR requiring appropriate clinical evaluation and documentation.

CASE STUDY 2: GASTROINTESTINAL DISTURBANCES ASSOCIATED WITH METFORMIN

- **Patient Information**

- Age: 52 years

- Gender: Male

- Medical condition: Type 2 Diabetes Mellitus

- Suspected medication: Metformin

- Dose: 500 mg twice daily (fictional case detail)

- Route: Oral

- **Clinical History:** A 52-year-old male patient with Type 2 Diabetes Mellitus was started on metformin for glycaemic control.
  Within four days of initiating treatment, the patient reported nausea, abdominal discomfort and loose stools.
  The patient did not report similar symptoms before starting the medication.

- **ADR Identification**

PARAMETER	FINDINGS
Suspected drug	Metformin
Adverse reaction	Nausea, abdominal discomfort and diarrhoea
Time of onset	4 days after treatment initiation
Nature of reaction	Gastrointestinal adverse reaction
Temporal relationship	Present
Alternative causes	Dietary factors, infection and other medications need evaluation

- Analysis

Metformin is commonly associated with gastrointestinal adverse effects, particularly during the initial phase of treatment.

The reported symptoms appeared shortly after starting metformin, which supports a possible relationship between the medication and the adverse reaction.

However, other gastrointestinal conditions and contributing factors should be considered.

- **Conclusion**

Metformin is identified as the suspected medication associated with nausea, abdominal discomfort and diarrhoea in this case.

CASE STUDY 3: MUSCLE PAIN ASSOCIATED WITH ATORVASTATIN

- **Patient Information**

- Age: 60 years

- Gender: Male

- Medical condition: Hypercholesterolaemia

- Suspected medication: Atorvastatin

- Dose: 20 mg once daily (fictional case detail)

- Route: Oral

- **Clinical History:** A 60-year-old male patient was prescribed atorvastatin for the management of elevated cholesterol levels.
  After approximately four weeks of treatment, the patient reported unexplained muscle pain and tenderness, particularly in the thighs.
  The patient had not experienced similar symptoms before starting the medication.

- **ADR Identification**

PARAMETER	FINDINGS
Suspected drug	Atorvastatin
Adverse reaction	Muscle pain and tenderness
Time of onset	Approximately 4 weeks
Nature of reaction	Possible statin-associated muscle symptoms
Temporal relationship	Present
Alternative causes	Physical activity, other medications and underlying muscle disorders need evaluation

- **Analysis**

Atorvastatin belongs to the statin class of lipid-lowering medications. Muscle-related adverse effects have been associated with statin therapy.

The development of muscle pain following the initiation of atorvastatin suggests a possible medication-related reaction.

Further clinical evaluation, including assessment of creatine kinase levels when indicated, may help investigate the reported symptoms.

- **Conclusion**

Atorvastatin is identified as the suspected medication associated with muscle pain and tenderness. Further clinical assessment is required to evaluate the relationship.

6. COMPARATIVE SUMMARY OF CASE STUDIES

PARAMETER	CASE 1	CASE 2	CASE 3
Suspected drug	Amoxicillin	Metformin	Atorvastatin
ADR	Skin rash, itching	Nausea, diarrhoea	Muscle pain
Onset	3 days	4 days	4 weeks
System affected	Skin	Gastrointestinal	Musculoskeletal
Suspected association	Possible	Possible	Possible

7. OBSERVATIONS AND FINDINGS

Based on the analysis of the three fictional clinical scenarios, the following observations were made:

- ADRs may affect different organ systems, including the skin, gastrointestinal tract and musculoskeletal system.
- The onset of an adverse reaction may vary depending on the medication and the individual patient.
- Reviewing the medication history is essential for identifying suspected drug-related reactions.
- Establishing the temporal relationship helps identify potential associations between medications and reported symptoms.
- The presence of a temporal relationship alone does not establish definitive causality.
- Alternative causes and concomitant medications must be considered during ADR identification.

8. IMPORTANCE OF ADR IDENTIFICATION

Early identification of suspected ADRs is important because it:

- Improves patient safety.

- Helps prevent medication-related complications.

- Supports appropriate clinical decision-making.

- Contributes to pharmacovigilance databases.

- Helps identify potential drug safety signals.

- Supports the safe and rational use of medicines.

9. CONCLUSION

This assignment provided an understanding of the basic principles of Adverse Drug Reaction identification through the analysis of three clinical case scenarios.

The cases demonstrated how patient symptoms, medication history and the temporal relationship between drug administration and symptom onset can help identify suspected ADRs.

The exercise also highlighted the importance of considering alternative causes before attributing a reaction to a particular medication.

ADR identification and reporting play an important role in pharmacovigilance and contribute to improving medication safety and patient care.

10. REFERENCES

1. Indian Pharmacopoeia Commission. Pharmacovigilance Programme of India (PvPI). ADR Reporting and Pharmacovigilance Information. https://ipc.gov.in/PvPI/adr.html

2. Indian Pharmacopoeia Commission. Pharmacovigilance Programme of India – Toolkit. https://ipc.gov.in/PvPI/toolkit.html

3. World Health Organization. Programme for International Drug Monitoring. https://www.who.int/teams/regulation-prequalification/regulation-and-safety/pharmacovigilance/networks/pidm