

Gwinnett School of Math, Science, and Technology

Multivariable Calculus Yearlong Notes

Anish Goyal
1st Period

Donny Thurston
Educator

2023-2024




Table of Contents

1 Chapter 1: Systems of Linear Equations and Matrices	5
1.1 Matrix Operations	5
1.1.1 Addition & Subtraction	5
1.1.2 Scalar Multiplication	5
1.1.3 Matrix Multiplication	5
1.1.4 Properties of Matrix Arithmetic	6
1.1.5 Examples	6
1.2 Transpose of a Matrix	7
1.2.1 Transpose Matrix Properties	7
1.3 Homework — “Matrix Stuff” (08/03/2023)	8
1.3.1 Suppose that A, B, C, D and E are matrices with the following sizes:	8
1.3.2 Consider the matrices	8
2 Intro to Systems	10
2.1 Review: Solve the following systems	11
2.1.1 Consistent	11
2.1.2 Inconsistent	11
2.2 The Augmented Matrix	12
2.3 Elementary Row Operations	12
2.3.1 Example 1... again	12
2.4 Connection to Matrices	12
2.4.1 Example 2: again	13
2.4.2 Example 3: again	13
2.4.3 Example 4: Solve the following system	13
2.4.4 Elementary Row Operations & REF Homework Problem (08/08/2023)	14
2.5 Gaussian Elimination	14
2.5.1 Examples	15
2.6 Gaussian Elimination With Back-Substitution	15
2.6.1 Goal:	15
2.6.2 Gaussian Elimination Homework Problem (08/09/2023)	16
2.7 Gauss-Jordan Elimination	17
2.7.1 Goal:	17
2.8 Matrix Properties, Equations, and Inverses	17
2.8.1 With Real Numbers	17
2.8.2 With Matrices	17
2.8.2.1 Multiply:	17
2.8.3 Matrix Inverses	18
3 Chapter 2: Determinants	19
3.1 Prior Knowledge:	19

3.2	Minors & Cofactors	19
3.2.1	Example	19
3.3	Cofactor Expansion	20
3.3.1	Example	20
3.3.2	Does the method generalize to 2×2 matrices?	21
3.3.3	Find the determinant of a 4×4	21
3.4	Theorem	21
3.4.1	Example	22
3.5	Triangular Matrices	22
3.6	An Important Definition	22
3.7	A Pair of Theorems	23
3.7.1	Theorem: If a square matrix A has a row of column of zeros, then $\det(A) = 0$	23
3.7.2	Theorem: If A is a square matrix, then $\det(A) = \det(A^T)$	23
3.8	Unit 1 & 2 Homework Problems	24
3.8.1	"Gaussian Elimination" (08/11/2023)	24
3.8.1.1	Solve this system using Gaussian Elimination	24
3.8.1.2	Solve this system using Gaussian Elimination	24
3.8.2	"Inverses and Determinants" (08/14)	25
3.8.2.1	Find the determinants of the following:	25
3.8.2.2	Find the INVERSES of those matrices:	25
3.8.3	Inverses and Determinants (08/15)	26
3.8.3.1	Use a matrix equation to solve the following problems:	26
3.8.4	Consistent Systems (08/21)	27
3.8.4.1	Solve the linear systems together by reducing the appropriate augmented matrix.	27
3.8.4.2	Determine the conditions on b , if any, in order to guarantee that the linear system is consistent.	28
3.8.5	Another "determining the conditions" problem:	28
3.8.6	Triangular and Diagonal Matrices	29
3.8.6.1	Find A^2	29
3.8.6.2	Find A^{-k} , such that k is some nonzero constant	30
3.8.6.3	Find a diagonal matrix A that satisfies the given condition	32
3.8.7	Determinants and Triangular Matrices (08/29)	33
3.8.7.1	What is C_{32}	33
3.8.7.2	Find all values of λ such that $ A = 0$	33
3.8.7.3	For the matrix $\begin{bmatrix} 3 & 0 & 0 \\ 2 & -1 & 5 \\ 1 & 9 & -4 \end{bmatrix}$ find the determinant 3 different ways with cofactor expansion. Pick different rows and columns each time.	34

3.8.7.4	Evaluate $\det(A)$ by a cofactor expansion along a row or column of your choice	35
3.8.7.5	Evaluate the determinant of the following matrices by just looking at them.	35
3.8.7.6	Show that the value of the determinant is independent of θ	35
3.8.8	Row operations and Determinants (08/31)	36
3.8.8.1	Find the determinant of $\begin{bmatrix} 1 & -3 & 0 \\ -2 & 4 & 1 \\ 5 & -2 & 2 \end{bmatrix}$ WITHOUT using cofactor expansion	36
3.8.8.2	Find the determinant of $\begin{bmatrix} 2 & 1 & 3 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 2 & 1 & 0 \\ 0 & 1 & 2 & 3 \end{bmatrix}$	37
3.8.9	Adjoins and Cramer's Rule (09/05)	38
3.8.9.1	Find the inverse of $A = \begin{bmatrix} 2 & 5 & 5 \\ -1 & -1 & 0 \\ 2 & 4 & 3 \end{bmatrix}$ using the adjoint method	38
3.8.9.2	Solve the following system of equations using Cramer's Rule	39
4	Chapter 5: Eigenvectors and Eigenvalues	40
4.0.1	Eigenvalues and Eigenvectors (11/06)	40
4.1	Examples	40
4.1.1	$\vec{x} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ is an eigenvector of $A = \begin{bmatrix} 3 & -2 \\ 1 & 0 \end{bmatrix}$ because	40
4.1.2	Let $A = \begin{bmatrix} 1 & 6 \\ 5 & 2 \end{bmatrix}$, $\vec{u} = \begin{bmatrix} 6 \\ -5 \end{bmatrix}$, $\vec{v} = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$. Are \vec{u} and \vec{v} eigenvectors of A ?	41
4.2	Eigenvector Homework Problem (11/06)	41
4.2.1	$A = \begin{bmatrix} 4 & 0 & 1 \\ 2 & 3 & 2 \\ 1 & 0 & 4 \end{bmatrix}$; $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$	41

1 Chapter 1: Systems of Linear Equations and Matrices

1.1 Matrix Operations

- Matrix operations are given as: rows x columns
- Two matrices are equal \Leftrightarrow they have the same dimensions and values

1.1.1 Addition & Subtraction

Two matrices can be added/subtracted \Leftrightarrow they have the same dimensions.

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} + \begin{bmatrix} 7 & 0 & 1 \\ -1 & 0 & 2 \end{bmatrix} = \begin{bmatrix} 8 & 2 & 4 \\ 3 & 5 & 8 \end{bmatrix}$$

1.1.2 Scalar Multiplication

- Scalar multiplication is defined as multiplying each element of a matrix by a number

$$3 \begin{bmatrix} 2 & 1 \\ 5 & 2 \end{bmatrix} = \begin{bmatrix} 6 & 3 \\ 15 & 6 \end{bmatrix}$$

1.1.3 Matrix Multiplication

- We can **only** multiply an $(m \times n)$ by $(n \times p)$ matrix.
- The resulting matrix will be $(m \times p)$

1.1.4 Properties of Matrix Arithmetic

- (a) $A + B = B + A$ (**Commutative law for addition**)
- (b) $A + (B + C) = (A + B) + C$ (**Associative law for addition**)
- (c) $A(BC) = (AB)C$ (**Associative law for multiplication**)
- (d) $A(B + C) = AB + AC$ (**Left distributive law**)
- (e) $(B + C)A = BA + CA$ (**Right distributive law**)
- (f) $A(B - C) = AB - AC$
- (g) $(B - C)A = BA - CA$
- (h) $a(B+C) = aB + aC$
- (i) $a(B-C) = aB - aC$
- (j) $(a+b)C = aC + bC$
- (k) $(a-b)C = aC - bC$
- (l) $a(bC) = (ab)C$
- (m) $a(BC) = (aB)C = B(aC)$

1.1.5 Examples

1.

$$\begin{aligned} & \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \\ &= \begin{bmatrix} 1 \cdot 1 + 2 \cdot 3 & 1 \cdot 2 + 2 \cdot 4 \\ 3 \cdot 1 + 4 \cdot 3 & 3 \cdot 2 + 4 \cdot 4 \end{bmatrix} \\ &= \begin{bmatrix} 7 & 10 \\ 15 & 22 \end{bmatrix} \end{aligned}$$

2.

$$\begin{aligned} & \begin{bmatrix} 2 & -3 \\ 5 & 0 \\ -2 & 4 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} -1 \\ 3 \end{bmatrix} \\ &= \begin{bmatrix} 2 \cdot (-1) + (-3) \cdot 3 \\ 5 \cdot (-1) + 0 \cdot 3 \\ -2 \cdot (-1) + 4 \cdot 3 \\ 1 \cdot (-1) + 2 \cdot 3 \end{bmatrix} \\ &= \begin{bmatrix} -11 \\ -5 \\ 14 \\ 5 \end{bmatrix} \end{aligned}$$

3.

$$\begin{aligned} & \begin{bmatrix} 4 & 5 & -1 \end{bmatrix} \begin{bmatrix} 8 \\ 0 \\ 2 \end{bmatrix} \\ &= [4 \cdot 8 + 5 \cdot 0 + (-1) \cdot 2] \\ &= [30] \end{aligned}$$

1.2 Transpose of a Matrix

The transpose of an $(m \times n)$ matrix is the $(n \times m)$ matrix where the rows and columns are swapped.

$$\text{If } B = \begin{bmatrix} 4 & 2 \\ -1 & 0 \\ 3 & 5 \end{bmatrix}, B^T = \begin{bmatrix} 4 & -1 & 3 \\ 2 & 0 & 5 \end{bmatrix}$$

$$\begin{aligned} B \cdot B^T &= \begin{bmatrix} 4 & 2 \\ -1 & 0 \\ 3 & 5 \end{bmatrix} \begin{bmatrix} 4 & -1 & 3 \\ 2 & 0 & 5 \end{bmatrix} \\ &= \begin{bmatrix} 4 \cdot 4 + 2 \cdot 2 & 4 \cdot (-1) + 2 \cdot 0 & 4 \cdot 3 + 2 \cdot 5 \\ (-1) \cdot 4 + 0 \cdot 2 & (-1) \cdot (-1) + 0 \cdot 0 & (-1) \cdot 3 + 0 \cdot 5 \\ 3 \cdot 4 + 5 \cdot 2 & 3 \cdot (-1) + 5 \cdot 0 & 3 \cdot 3 + 5 \cdot 5 \end{bmatrix} \\ &= \begin{bmatrix} 20 & -4 & 22 \\ -4 & 1 & -3 \\ 22 & -3 & 34 \end{bmatrix} \end{aligned}$$

- The transpose of a matrix is **always** multiplicative with the original.
- There is also a **main diagonal** that is the diagonal from the top left to the bottom right, but only square matrices have these.
- The **trace** of a square matrix A is equal to the sum of all the elements on the main diagonal: $\text{tr}(A)$

1.2.1 Transpose Matrix Properties

- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(A - B)^T = A^T - B^T$
- $(kA)^T = kA^T$
- $(AB)^T = B^T A^T$

1.3 Homework — “Matrix Stuff” (08/03/2023)

1.3.1 Suppose that A, B, C, D and E are matrices with the following sizes:

A	B	C	D	E
(3×2)	(2×3)	(3×3)	(3×2)	(2×3)

For each matrix operation, sort them into undefined if the operation can't be done, or defined if it can along with the correct dimensions of the outcome.

Undefined	Defined; (4×2)	Defined; (5×5)	Defined; (5×2)
BA	$AC + D$	$E(A + B)$	$(A^T + E)D$
$AB + B$			$E(AC)$
$E^T A$			
$AE + B$			

1.3.2 Consider the matrices

$$A = \begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix}, B = \begin{bmatrix} 4 & -1 \\ 0 & 2 \end{bmatrix}, C = \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix}, D = \begin{bmatrix} 1 & 5 & 2 \\ -1 & 0 & 1 \\ 3 & 2 & 4 \end{bmatrix}, E = \begin{bmatrix} 6 & 1 & 3 \\ -1 & 1 & 2 \\ 4 & 1 & 3 \end{bmatrix}$$

In each part, compute the given expression (where possible).

2. $2A^T + C$

$$\begin{aligned} 2A^T + C &= 2 \begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix}^T + \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix} \\ &= 2 \begin{bmatrix} 3 & -1 & 1 \\ 0 & 2 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix} \\ &= \begin{bmatrix} 6 & -2 & 2 \\ 0 & 4 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix} \\ &= \begin{bmatrix} 7 & 2 & 4 \\ 3 & 5 & 7 \end{bmatrix} \end{aligned}$$

3. $B^T + 5C^T$

$$\begin{aligned}
 B^T + 5C^T &= \begin{bmatrix} 4 & -1 \\ 0 & 2 \end{bmatrix}^T + 5 \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix}^T \\
 &= \begin{bmatrix} 4 & 0 \\ -1 & 2 \end{bmatrix} + 5 \begin{bmatrix} 1 & 3 \\ 4 & 1 \\ 2 & 5 \end{bmatrix} \\
 &= \begin{bmatrix} 4 & 0 \\ -1 & 2 \end{bmatrix} + \begin{bmatrix} 5 & 15 \\ 20 & 5 \\ 10 & 25 \end{bmatrix} \\
 &= \text{Undefined}
 \end{aligned}$$

4. $2E^T - 3D^T$

$$\begin{aligned}
 2E^T - 3D^T &= 2 \begin{bmatrix} 6 & 1 & 3 \\ -1 & 1 & 2 \\ 4 & 1 & 3 \end{bmatrix}^T - 3 \begin{bmatrix} 1 & 5 & 2 \\ -1 & 0 & 1 \\ 3 & 2 & 4 \end{bmatrix}^T \\
 &= 2 \begin{bmatrix} 6 & -1 & 4 \\ 1 & 1 & 1 \\ 3 & 2 & 3 \end{bmatrix} - 3 \begin{bmatrix} 1 & -1 & 3 \\ 5 & 0 & 2 \\ 2 & 1 & 4 \end{bmatrix} \\
 &= \begin{bmatrix} 12 & -2 & 8 \\ 2 & 2 & 2 \\ 6 & 4 & 6 \end{bmatrix} - \begin{bmatrix} 3 & -3 & 9 \\ 15 & 0 & 6 \\ 6 & 3 & 12 \end{bmatrix} \\
 &= \begin{bmatrix} 9 & -5 & -1 \\ -13 & 2 & -4 \\ 0 & 1 & -6 \end{bmatrix}
 \end{aligned}$$

5. $\text{tr}(DE)$

$$\begin{aligned}
 \text{tr}(DE) &= \text{tr} \left(\begin{bmatrix} 1 & 5 & 2 \\ -1 & 0 & 1 \\ 3 & 2 & 4 \end{bmatrix} \begin{bmatrix} 6 & 1 & 3 \\ -1 & 1 & 2 \\ 4 & 1 & 3 \end{bmatrix} \right) \\
 &= \text{tr} \left(\begin{bmatrix} 1 \cdot 6 + 5 \cdot (-1) + 2 \cdot 4 & 1 \cdot 1 + 5 \cdot 1 + 2 \cdot 1 & 1 \cdot 3 + 5 \cdot 2 + 2 \cdot 3 \\ (-1) \cdot 6 + 0 \cdot (-1) + 1 \cdot 4 & (-1) \cdot 1 + 0 \cdot 1 + 1 \cdot 1 & (-1) \cdot 3 + 0 \cdot 2 + 1 \cdot 3 \\ 3 \cdot 6 + 2 \cdot (-1) + 4 \cdot 4 & 3 \cdot 1 + 2 \cdot 1 + 4 \cdot 1 & 3 \cdot 3 + 2 \cdot 2 + 4 \cdot 3 \end{bmatrix} \right) \\
 &= \text{tr} \left(\begin{bmatrix} 9 & 8 & 19 \\ -2 & 0 & 0 \\ 32 & 9 & 25 \end{bmatrix} \right) \\
 &= 34
 \end{aligned}$$

2 Intro to Systems

What are we looking for?

Lines: How many possible solutions?

- Infinite solutions
- One solution
- No solutions


Planes: How many possible solutions?

- Infinite solutions
- No solutions

What does linear actually mean?

- The word linear really means that you've got equations with variables and **all** of the variables are degree one.
- This means that there is no limit to the number of dimensions in a linear system.

Linear Systems in Three Unknowns



2.1 Review: Solve the following systems

1.
$$\begin{cases} 2x + y = 10 \\ 3x - y = 5 \end{cases}$$

$$5x = 15$$

$$x = 3$$

$$2(3) + y = 10$$

$$6 + y = 10$$

$$y = 4$$

2.
$$\begin{cases} 2x + y = 10 \\ 6x + 3y = 10 \end{cases}$$

$$y = 10 - 2x$$

$$6x + 3(10 - 2x) = 10$$

$$6x + 30 - 6x = 10$$

$$30 = 10 \therefore \text{no solution}$$

3.
$$\begin{cases} 5x - 2y = 4 \\ 15x - 6y = 12 \end{cases}$$

$$0 = 0$$

$$12 = 12 \therefore \text{infinite solutions}$$

2.1.1 Consistent

- A system of equations is **consistent** if it has at least one solution.

2.1.2 Inconsistent

- A system of equations is **inconsistent** if it has no solutions.

2.2 The Augmented Matrix

$$\begin{cases} x - y + 2z = 5 \\ 2x - 2y + 4z = 10 \\ 3x - 3y + 6z = 15 \end{cases} \longrightarrow \left[\begin{array}{ccc|c} 1 & -1 & 2 & 5 \\ 2 & -2 & 4 & 10 \\ 3 & -3 & 6 & 15 \end{array} \right]$$

2.3 Elementary Row Operations

1. Interchange 2 rows
2. Multiply a row by a non-zero constant
3. Add/subtract a multiple of one row to/from another row

Doing these things changes the matrix, but it's the same system!

2.3.1 Example 1... again

$$\begin{cases} 2x + y = 10 \\ 3x - y = 5 \end{cases}$$

$$\begin{aligned} \left[\begin{array}{cc|c} 2 & 1 & 10 \\ 3 & -1 & 5 \end{array} \right] &\xrightarrow{\frac{1}{2}R_1} \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 5 \\ 3 & -1 & 5 \end{array} \right] \xrightarrow{R_2 - 3R_1} \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 5 \\ 0 & -\frac{5}{2} & -10 \end{array} \right] \\ &\xrightarrow{-\frac{2}{5}R_2} \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 5 \\ 0 & 1 & 4 \end{array} \right] \xrightarrow{R_1 - \frac{1}{2}R_2} \left[\begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 4 \end{array} \right] \end{aligned}$$

And so... $x = 3$ and $y = 4$!

2.4 Connection to Matrices

If we can make a system's matrix look like

$$\left[\begin{array}{ccc|c} 1 & 0 & 0 & c_1 \\ 0 & 1 & 0 & c_2 \\ 0 & 0 & 1 & c_3 \end{array} \right],$$

then the solution to the system will be the ordered triple (c_1, c_2, c_3) .

2.4.1 Example 2: again

$$\begin{cases} 2x + y = 10 \\ 6x + 3y = 10 \end{cases}$$

$$\left[\begin{array}{cc|c} 2 & 1 & 10 \\ 6 & 3 & 10 \end{array} \right] \xrightarrow{\frac{1}{2}R_1} \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 5 \\ 6 & 3 & 10 \end{array} \right] \xrightarrow{R_2-6R_1} \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 5 \\ 0 & 0 & -20 \end{array} \right]$$

This is inconsistent, so there is no solution.

2.4.2 Example 3: again

$$\begin{cases} 5x - 2y = 4 \\ 15x - 6y = 12 \end{cases}$$

$$\left[\begin{array}{cc|c} 5 & -2 & 4 \\ 15 & -6 & 12 \end{array} \right] \xrightarrow{\frac{1}{5}R_1} \left[\begin{array}{cc|c} 1 & -\frac{2}{5} & \frac{4}{5} \\ 15 & -6 & 12 \end{array} \right] \xrightarrow{R_2-15R_1} \left[\begin{array}{cc|c} 1 & -\frac{2}{5} & \frac{4}{5} \\ 0 & 0 & 0 \end{array} \right]$$

Since $0 = 0$, there are infinitely many solutions.

2.4.3 Example 4: Solve the following system

$$\begin{cases} x_1 - 2x_2 + x_3 = 0 \\ 2x_2 - 8x_3 = 8 \\ -4x_1 + 5x_2 + 9x_3 = -9 \end{cases}$$

$$\begin{aligned} & \left[\begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 2 & -8 & 8 \\ -4 & 5 & 9 & -9 \end{array} \right] \xrightarrow{R_3+4R_1} \left[\begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 2 & -8 & 8 \\ 0 & -3 & 13 & -9 \end{array} \right] \xrightarrow{R_3+\frac{3}{2}R_2} \left[\begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 2 & -8 & 8 \\ 0 & 0 & -1 & 3 \end{array} \right] \\ & \xrightarrow{\frac{1}{2}R_2} \left[\begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 1 & -4 & 4 \\ 0 & -3 & 13 & -9 \end{array} \right] \xrightarrow{R_1+2R_2} \left[\begin{array}{ccc|c} 1 & 0 & -7 & 8 \\ 0 & 1 & -4 & 4 \\ 0 & 0 & 1 & 3 \end{array} \right] \xrightarrow{\begin{matrix} R_1+7R_3 \\ R_2+4R_3 \end{matrix}} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 29 \\ 0 & 1 & 0 & 16 \\ 0 & 0 & 1 & 3 \end{array} \right] \end{aligned}$$

Therefore the solution to (x_1, x_2, x_3) is $(29, 16, 3)$.

2.4.4 Elementary Row Operations & REF Homework Problem (08/08/2023)

$$\begin{cases} x + y + 2z = 8 \\ -x - 2y + 3z = 1 \\ 3x - 7y + 4z = 10 \end{cases}$$

$$\begin{aligned} & \left[\begin{array}{ccc|c} 1 & 1 & 2 & 8 \\ -1 & -2 & 3 & 1 \\ 3 & -7 & 4 & 10 \end{array} \right] \xrightarrow[\substack{R_2+R_1 \\ R_3-3R_1}]{\substack{R_2+R_1 \\ R_3-3R_1}} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 8 \\ 0 & -1 & 5 & 9 \\ 0 & -10 & -2 & -14 \end{array} \right] \xrightarrow[\substack{-R_2 \\ -R_3}]{\substack{-R_2 \\ -R_3}} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 8 \\ 0 & 1 & -5 & -9 \\ 0 & 10 & 2 & 14 \end{array} \right] \\ & \xrightarrow[\substack{R_1-R_2 \\ R_3-10R_2}]{\substack{R_1-R_2 \\ R_3-10R_2}} \left[\begin{array}{ccc|c} 1 & 0 & 7 & 17 \\ 0 & 1 & -5 & -9 \\ 0 & 0 & 52 & 104 \end{array} \right] \xrightarrow[\substack{1/52 R_3}]{\substack{1/52 R_3}} \left[\begin{array}{ccc|c} 1 & 0 & 7 & 17 \\ 0 & 1 & -5 & -9 \\ 0 & 0 & 1 & 2 \end{array} \right] \xrightarrow[\substack{R_1-7R_3 \\ R_2+5R_3}]{\substack{R_1-7R_3 \\ R_2+5R_3}} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 2 \end{array} \right] \end{aligned}$$

Therefore, the solution to (x, y, z) is $(3, 1, 2)$.

2.5 Gaussian Elimination

Vocabulary: A matrix is in Row Echelon Form (REF) if:

- (a) Any rows of all zeroes are placed at the bottom of the matrix
- (b) All other rows have a leading 1 ("pivot")
- (c) As we move down the matrix, each leading 1 is further to the right than the 1 above it

A matrix is in Row Reduced Echelon Form if the three above conditions are met in addition to:

- (d) Each column with a leading 1 has all other entries in the column as a 0. ("pivot column")

2.5.1 Examples

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 8 \\ 0 & 1 & 0 & 6 & -3 \\ 0 & 0 & 1 & 7 & 10 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

REF? ✓
RREF? ✓

$$\begin{bmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

REF? ✓
RREF? ✗

$$\begin{bmatrix} 1 & 2 & -3 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & -4 \end{bmatrix}$$

REF? ✗
RREF? ✗

2.6 Gaussian Elimination With Back-Substitution

2.6.1 Goal:

To get the augmented matrix in REF

Solve:
$$\begin{cases} x_1 - 2x_2 + 3x_3 = 9 \\ -x_1 + 3x_2 = -4 \\ 2x_1 - 5x_2 + 5x_3 = 17 \end{cases}$$

$$\begin{aligned} & \left[\begin{array}{ccc|c} 1 & -2 & 3 & 9 \\ -1 & 3 & 0 & -4 \\ 2 & -5 & 5 & 17 \end{array} \right] \xrightarrow[R_3-2R_1]{R_2+R_1} \left[\begin{array}{ccc|c} 1 & -2 & 3 & 9 \\ 0 & 1 & 3 & 5 \\ 0 & -1 & -1 & -1 \end{array} \right] \xrightarrow[R_3+R_2]{R_1+2R_2} \left[\begin{array}{ccc|c} 1 & 0 & 9 & 19 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 2 & 4 \end{array} \right] \\ & \xrightarrow{\frac{1}{2}R_3} \left[\begin{array}{ccc|c} 1 & 0 & 9 & 19 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 1 & 2 \end{array} \right] \end{aligned}$$

$$x + 9z = 19$$

$$y + 3z = 5$$

$$z = 2$$

$$\therefore z = 2, y = 5 - 3z, x = 19 - 9z$$

$$z = 2, y = 5 - 3(2), x = 19 - 9(2)$$

$$z = 2, y = -1, x = 1$$

Therefore, the solution (x_1, x_2, x_3) is $(1, -1, 2)$.

2.6.2 Gaussian Elimination Homework Problem (08/09/2023)

$$\begin{cases} -2w + y + z = -3 \\ x + 2y - z = 2 \\ -3w + 2x + 4y + z = -2 \\ -w + x - 4y - 7z = -19 \end{cases}$$

$$\begin{aligned} & \left[\begin{array}{cccc|c} -2 & 0 & 1 & 1 & -3 \\ 0 & 1 & 2 & -1 & 2 \\ -3 & 2 & 4 & 1 & -2 \\ -1 & 1 & -4 & -7 & -19 \end{array} \right] \xrightarrow{R_4} \left[\begin{array}{cccc|c} -1 & 1 & -4 & -7 & -19 \\ 0 & 1 & 2 & -1 & 2 \\ -3 & 2 & 4 & 1 & -2 \\ -2 & 0 & 1 & 1 & -3 \end{array} \right] \xrightarrow{-R_1} \\ & \left[\begin{array}{cccc|c} 1 & -1 & 4 & 7 & 19 \\ 0 & 1 & 2 & -1 & 2 \\ -3 & 2 & 4 & 1 & -2 \\ -2 & 0 & 1 & 1 & -3 \end{array} \right] \xrightarrow{\begin{array}{l} R_3+3R_1 \\ R_4+2R_1 \end{array}} \left[\begin{array}{cccc|c} 1 & -1 & 4 & 7 & 19 \\ 0 & 1 & 2 & -1 & 2 \\ 0 & -1 & 16 & 22 & 55 \\ 0 & -2 & 9 & 15 & 35 \end{array} \right] \xrightarrow{\begin{array}{l} R_1+R_2 \\ R_3+R_2 \\ R_4+2R_2 \end{array}} \\ & \left[\begin{array}{cccc|c} 1 & 0 & 6 & 6 & 21 \\ 0 & 1 & 2 & -1 & 2 \\ 0 & 0 & 18 & 21 & 57 \\ 0 & 0 & 13 & 13 & 39 \end{array} \right] \xrightarrow{\frac{1}{18}R_3} \left[\begin{array}{cccc|c} 1 & 0 & 6 & 6 & 21 \\ 0 & 1 & 2 & -1 & 2 \\ 0 & 0 & 1 & \frac{7}{6} & \frac{19}{6} \\ 0 & 0 & 13 & 13 & 39 \end{array} \right] \xrightarrow{\begin{array}{l} R_1-6R_3 \\ R_2-2R_3 \\ R_4-13R_3 \end{array}} \\ & \left[\begin{array}{cccc|c} 1 & 0 & 0 & -1 & 2 \\ 0 & 1 & 0 & -\frac{10}{3} & -\frac{13}{3} \\ 0 & 0 & 1 & \frac{7}{6} & \frac{19}{6} \\ 0 & 0 & 0 & -\frac{13}{6} & -\frac{13}{6} \end{array} \right] \xrightarrow{-\frac{6}{13}R_4} \left[\begin{array}{cccc|c} 1 & 0 & 0 & -1 & 2 \\ 0 & 1 & 0 & -\frac{10}{3} & -\frac{13}{3} \\ 0 & 0 & 1 & \frac{7}{6} & \frac{19}{6} \\ 0 & 0 & 0 & 1 & 1 \end{array} \right] \xrightarrow{\begin{array}{l} R_1+R_4 \\ R_2+\frac{10}{3}R_4 \\ R_3-\frac{7}{6}R_4 \end{array}} \\ & \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 3 \\ 0 & 1 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right] \Rightarrow \begin{cases} w = 3 \\ x = -1 \\ y = 2 \\ z = 1 \end{cases} \end{aligned}$$

2.7 Gauss-Jordan Elimination

2.7.1 Goal:

To get the matrix into RREF

$$\text{Solve: } \begin{cases} x_1 - 3x_3 = -2 \\ 3x_1 + x_2 - 2x_3 = 5 \\ 2x_1 + 2x_2 + x_3 = 4 \end{cases}$$

$$\begin{aligned} & \left[\begin{array}{ccc|c} 1 & 0 & -3 & -2 \\ 3 & 1 & -2 & 5 \\ 2 & 2 & 1 & 4 \end{array} \right] \xrightarrow[R_3-2R_1]{R_2-3R_1} \left[\begin{array}{ccc|c} 1 & 0 & -3 & -2 \\ 0 & 1 & 7 & 11 \\ 0 & 2 & 7 & 8 \end{array} \right] \xrightarrow{R_3-2R_2} \left[\begin{array}{ccc|c} 1 & 0 & -3 & -2 \\ 0 & 1 & 7 & 11 \\ 0 & 0 & -7 & -14 \end{array} \right] \\ & \xrightarrow{\frac{-1}{7}R_3} \left[\begin{array}{ccc|c} 1 & 0 & -3 & -2 \\ 0 & 1 & 7 & 11 \\ 0 & 0 & 1 & 2 \end{array} \right] \xrightarrow[R_2-7R_3]{R_1+3R_3} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 4 \\ 0 & 1 & 0 & -3 \\ 0 & 0 & 1 & 2 \end{array} \right] \Rightarrow \begin{cases} x_1 = 4 \\ x_2 = -3 \\ x_3 = 2 \end{cases} \end{aligned}$$

2.8 Matrix Properties, Equations, and Inverses

2.8.1 With Real Numbers

- If $ab = bc$, then $a = c$, if $b \neq 0$
- If $ab = 0$, then $a = 0$ or $b = 0$, or both

2.8.2 With Matrices

- If $AB = AC$, then $B = C$, if A is invertible
- If $AB = [0]$, then $A = [0]$ or $B = [0]$, or both

2.8.2.1 Multiply:

$$\begin{aligned} & \begin{bmatrix} 2 & 3 \\ 3 & 5 \end{bmatrix} \begin{bmatrix} 5 & -3 \\ -3 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 2(5) + 3(-3) & 2(-3) + 3(2) \\ 3(5) + 5(-3) & 3(-3) + 5(2) \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

2.8.3 Matrix Inverses

- If a matrix has an inverse, it is said to be invertible or non-singular.
- If a matrix does not have an inverse, it is said to be singular.
- Every square matrix has a “special number” associated with it called the **determinant**.
- For the 2×2 matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, the determinant is $ad - bc$
- $A^{-1} = \frac{1}{\det A} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$
- When $\det A = 0$, the matrix is singular and has no inverse (since you cannot divide by zero)

Find the inverse of $A = \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}$

$$\begin{aligned} \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}^{-1} &= \frac{1}{\det A} \begin{bmatrix} 2 & -3 \\ -1 & 4 \end{bmatrix} \\ &= \frac{1}{(4)(2) - (3)(1)} \begin{bmatrix} 2 & -3 \\ -1 & 4 \end{bmatrix} \\ &= \frac{1}{5} \begin{bmatrix} 2 & -3 \\ -1 & 4 \end{bmatrix} \\ &= \begin{bmatrix} \frac{2}{5} & -\frac{3}{5} \\ -\frac{1}{5} & \frac{4}{5} \end{bmatrix} \end{aligned}$$

3 Chapter 2: Determinants

3.1 Prior Knowledge:

$$\begin{bmatrix} 10 & -4 \\ -3 & -5 \end{bmatrix} = -50 - = -62$$

$$\begin{aligned} & \begin{bmatrix} 2 & 4 & 3 \\ -1 & 2 & 3 \\ 3 & 0 & -2 \end{bmatrix} \\ & = ((2 \cdot 2 \cdot -2) + (4 \cdot 3 \cdot 3) + (3 \cdot -1 \cdot 0)) - ((3 \cdot 2 \cdot 3) + (0 \cdot 3 \cdot 2) + (-2 \cdot -1 \cdot 4)) \\ & = (-8 + 36 + 0) - (18 + 0 + 8) \\ & = 28 - 26 \\ & = 2 \end{aligned}$$

3.2 Minors & Cofactors

Given a square matrix A, the minor of matrix element a_{ij} , (M_{ij}) is the determinant of the matrix formed by removing the i^{th} row and j^{th} column from matrix A.

The cofactor of matrix element a_{ij} , $C_{ij} = (-1)^{i+j} \cdot M_{ij}$

3.2.1 Example

$$\text{Let } \det \begin{bmatrix} 2 & 4 & 3 \\ -1 & 2 & 3 \\ 3 & 0 & -2 \end{bmatrix}. \text{ What is the cofactor of element } (1, 1)?$$

Cofactor checkerboard:

$$\begin{bmatrix} + & - & + \\ - & + & - \\ + & - & + \end{bmatrix}$$

$$M_{11} = \begin{vmatrix} 2 & 3 \\ 0 & -2 \end{vmatrix} = -4$$

$$C_{11} = 1 \cdot -4 = -4$$

Find the minor and cofactor of: \ a) $a_{21} = -1$

$$M_{21} = \begin{vmatrix} 4 & 3 \\ 0 & -2 \end{vmatrix} = -8$$

$$C_{21} = 8$$

b) $a_{33} = -2$

$$M_{33} = \begin{vmatrix} 2 & 4 \\ -1 & 2 \end{vmatrix} = 8$$

$$C_{33} = 8$$

3.3 Cofactor Expansion

- 1) Pick a row or column
- 2) Multiply every entry in that row or column by it's corresponding cofactor
- 3) Add those together. That's it

$$A = \begin{bmatrix} 6 & 7 & -1 \\ 0 & 4 & 1 \\ 2 & 5 & -3 \end{bmatrix}$$

$$\begin{aligned} \det(A) &= 6 \begin{vmatrix} 4 & 1 \\ 5 & -3 \end{vmatrix} + 7 \begin{vmatrix} 0 & 1 \\ 2 & -3 \end{vmatrix} + -1 \begin{vmatrix} 0 & 4 \\ 2 & 5 \end{vmatrix} \\ &= 6(-17) + 7(2) + (-1(-8)) \\ &= -102 + 14 + 8 \\ &= -80 \end{aligned}$$

3.3.1 Example

$$A = \begin{bmatrix} 6 & 4 & 2 \\ 5 & -6 & 1 \\ 0 & 3 & 0 \end{bmatrix}$$

$$\begin{aligned} &6 \begin{vmatrix} -6 & 1 \\ 3 & 0 \end{vmatrix} + 4 \begin{vmatrix} 5 & 1 \\ 0 & 0 \end{vmatrix} + 2 \begin{vmatrix} 5 & -6 \\ 0 & 3 \end{vmatrix} \\ &= 6(-3) + 0 + 2(15) \\ &= -18 + 30 \\ &= 12 \end{aligned}$$

3.3.2 Does the method generalize to 2×2 matrices?

$$\begin{aligned} & \begin{vmatrix} 3 & 5 \\ 7 & 2 \end{vmatrix} \\ &= 3|2| - 5|7| \\ &= 6 - 35 \\ &= -29 \end{aligned}$$

The determinant of a 1×1 matrix is... **itself!**

3.3.3 Find the determinant of a 4×4

$$A = \begin{bmatrix} -3 & 2 & 0 & 8 \\ 2 & 1 & 0 & -4 \\ 5 & -2 & 1 & 5 \\ 2 & 3 & 0 & 6 \end{bmatrix}$$

$$\begin{aligned} &= 0 + 0 + \begin{vmatrix} -3 & 2 & 8 \\ 2 & 1 & -4 \\ 2 & 3 & 6 \end{vmatrix} + 0 \\ &= -2 \begin{vmatrix} 2 & 8 \\ 3 & 6 \end{vmatrix} + \begin{vmatrix} -3 & 8 \\ 2 & 6 \end{vmatrix} - \left(-4 \begin{vmatrix} -3 & 2 \\ 2 & 3 \end{vmatrix} \right) \\ &= 24 - 34 - 52 \\ &= -62 \end{aligned}$$

3.4 Theorem

If A is an $n \times n$ matrix, then regardless of which row or column of A is chosen, the number obtained by multiplying the elements in that row or column by their corresponding cofactors is **always the same** and is called the determinant of A .

3.4.1 Example

Find the determinant of $A = \begin{bmatrix} 1 & 0 & 0 & -1 \\ 3 & 1 & 2 & 2 \\ 1 & 0 & -2 & 1 \\ 2 & 0 & 0 & 1 \end{bmatrix}$

$$\begin{aligned} & 1 \cdot \begin{vmatrix} 1 & 0 & -1 \\ 1 & -2 & 1 \\ 2 & 0 & 1 \end{vmatrix} \\ &= \left(-2 \begin{vmatrix} 1 & -1 \\ 2 & 1 \end{vmatrix} \right) \\ &= -6 \end{aligned}$$

3.5 Triangular Matrices

Find the determinant of $A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 2 & 2 \\ 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 4 \end{bmatrix}$

$$\begin{aligned} & \begin{vmatrix} 2 & 2 & 2 \\ 0 & 3 & 3 \\ 0 & 0 & 4 \end{vmatrix} \\ &= 2 \begin{vmatrix} 3 & 3 \\ 0 & 4 \end{vmatrix} \\ &= 2(3 \cdot 4) \\ &= 2 \cdot 12 \\ &= 24 \end{aligned}$$

If A is an $n \times n$ triangular matrix, then $\det(A)$ is equal to the product of the elements along the main diagonal.

3.6 An Important Definition

Elementary Matrix a matrix that can be obtained from the $n \times n$ identity matrix by performing a single row operation. \

Are the following matrices elementary? 1) $\begin{bmatrix} 1 & 0 \\ -5 & 1 \end{bmatrix} + (R_3 + 5R_1)$ yes 2) $\begin{bmatrix} -5 & 1 \\ 1 & 0 \end{bmatrix} + (R_1 + 5R_2)$...
no

3.7 A Pair of Theorems

3.7.1 Theorem: If a square matrix A has a row of column of zeros, then $\det(A) = 0$

3.7.2 Theorem: If A is a square matrix, then $\det(A) = \det(A^T)$

3.8 Unit 1 & 2 Homework Problems

3.8.1 "Gaussian Elimination" (08/11/2023)

3.8.1.1 Solve this system using Gaussian Elimination

$$\begin{cases} x_1 + x_2 + 2x_3 = 8 \\ -x_1 - 2x_2 + 3x_3 = 1 \\ 3x_1 - 7x_2 + 4x_3 = 10 \end{cases}$$

$$\Rightarrow \left[\begin{array}{ccc|c} 1 & 1 & 2 & 8 \\ -1 & -2 & 3 & 1 \\ 3 & -7 & 4 & 10 \end{array} \right] \xrightarrow[\substack{R_2+R_1 \\ R_3-3R_1}]{\substack{R_2+R_1 \\ R_3-3R_1}} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 8 \\ 0 & -1 & 5 & 9 \\ 0 & -10 & -2 & -14 \end{array} \right] \xrightarrow{-R_2} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 8 \\ 0 & 1 & -5 & -9 \\ 0 & -10 & -2 & -14 \end{array} \right]$$

$$\xrightarrow{R_3+10R_2} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 8 \\ 0 & 1 & -5 & -9 \\ 0 & 0 & -52 & -104 \end{array} \right] \xrightarrow{-\frac{1}{52}R_3} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 8 \\ 0 & 1 & -5 & -9 \\ 0 & 0 & 1 & 2 \end{array} \right]$$

$$\therefore \begin{cases} x_1 + x_2 + 2x_3 = 8 \\ x_2 - 5x_3 = -9 \\ x_3 = 2 \end{cases} \Rightarrow \begin{cases} x_1 = 3 \\ x_2 = 1 \\ x_3 = 2 \end{cases}$$

3.8.1.2 Solve this system using Gaussian Elimination

$$\begin{cases} x_1 - 2x_2 + 3x_3 = 0 \\ -2x_1 - 3x_2 - 4x_3 = 0 \\ 2x_1 - 4x_2 + 4x_3 = 0 \end{cases}$$

$$\Rightarrow \left[\begin{array}{ccc|c} 1 & -2 & 3 & 0 \\ -2 & -3 & -4 & 0 \\ 2 & -4 & 4 & 0 \end{array} \right] \xrightarrow[\substack{R_2+2R_1 \\ R_3-2R_1}]{\substack{R_2+2R_1 \\ R_3-2R_1}} \left[\begin{array}{ccc|c} 1 & -2 & 3 & 0 \\ 0 & -7 & 2 & 0 \\ 0 & 0 & -2 & 0 \end{array} \right] \xrightarrow[\substack{-\frac{1}{7}R_2 \\ -\frac{1}{2}R_3}]{\substack{-\frac{1}{7}R_2 \\ -\frac{1}{2}R_3}} \left[\begin{array}{ccc|c} 1 & -2 & 3 & 0 \\ 0 & 1 & \frac{2}{7} & 0 \\ 0 & 0 & 1 & 0 \end{array} \right]$$

$$\therefore \begin{cases} x_1 - 2x_2 + 3x_3 = 0 \\ x_2 + \frac{2}{7}x_3 = 0 \\ x_3 = 0 \end{cases} \Rightarrow 1 \neq 0 \therefore \text{no solution}$$

3.8.2 "Inverses and Determinants" (08/14)

3.8.2.1 Find the determinants of the following:

$$1) \begin{bmatrix} 2 & -3 \\ 4 & 4 \end{bmatrix}$$

$$\begin{vmatrix} 2 & -3 \\ 4 & 4 \end{vmatrix} = 2(4) - (-3)(4) = 8 + 12 = 20$$

$$2) \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$$

$$\begin{vmatrix} 2 & 0 \\ 0 & 3 \end{vmatrix} = 2(3) - 0(0) = 6$$

$$3) \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

$$\begin{vmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{vmatrix} = \cos^2 \theta + \sin^2 \theta = 1$$

3.8.2.2 Find the INVERSES of those matrices:

$$1) \begin{bmatrix} 2 & -3 \\ 4 & 4 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -3 \\ 4 & 4 \end{bmatrix}^{-1} = \frac{1}{20} \begin{bmatrix} 4 & 3 \\ -4 & 2 \end{bmatrix} = \begin{bmatrix} \frac{1}{5} & \frac{3}{20} \\ -\frac{1}{5} & \frac{1}{10} \end{bmatrix}$$

$$2) \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}^{-1} = \frac{1}{6} \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{3} \end{bmatrix}$$

$$3) \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

$$\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}^{-1} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

3.8.3 Inverses and Determinants (08/15)

3.8.3.1 Use a matrix equation to solve the following problems:

$$1) \begin{cases} 3x_1 - 2x_2 = 1 \\ 4x_1 + 5x_2 = 3 \end{cases}$$

$$\Rightarrow \begin{bmatrix} 3 & -2 \\ 4 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} -1 \\ 3 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 3 & -2 \\ 4 & 5 \end{bmatrix}^{-1} \begin{bmatrix} -1 \\ 3 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \frac{1}{23} \begin{bmatrix} 5 & 2 \\ -4 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ 3 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \frac{1}{23} \begin{bmatrix} -1 \\ 9 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} -\frac{1}{23} \\ \frac{9}{23} \end{bmatrix}$$

$$2) \begin{cases} 6x_1 + x_2 = 0 \\ 4x_1 - 3x_2 = -2 \end{cases}$$

$$\Rightarrow \begin{bmatrix} 6 & 1 \\ 4 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ -2 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 6 & 1 \\ 4 & -3 \end{bmatrix}^{-1} \begin{bmatrix} 0 \\ -2 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \frac{1}{-22} \begin{bmatrix} -3 & -1 \\ -4 & 6 \end{bmatrix} \begin{bmatrix} 0 \\ -2 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \frac{1}{-22} \begin{bmatrix} 2 \\ 8 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} -\frac{1}{11} \\ \frac{4}{11} \end{bmatrix}$$

3.8.4 Consistent Systems (08/21)

3.8.4.1 Solve the linear systems together by reducing the appropriate augmented matrix.

$$\begin{cases} x_1 - 5x_2 = b_1 \\ 3x_1 + 2x_2 = b_2 \end{cases}$$

- 1) $b_1 = 1, b_2 = 4$
- 2) $b_1 = -2, b_2 = 5$

First, let's solve it for the general case:

$$\left[\begin{array}{cc|c} 1 & -5 & b_1 \\ 3 & 2 & b_2 \end{array} \right] \xrightarrow{R_2 - 3R_1} \left[\begin{array}{cc|c} 1 & -5 & b_1 \\ 0 & 17 & b_2 - 3b_1 \end{array} \right] \xrightarrow{\frac{1}{17}R_2} \left[\begin{array}{cc|c} 1 & -5 & b_1 \\ 0 & 1 & \frac{b_2 - 3b_1}{17} \end{array} \right] \xrightarrow{R_1 + 5R_2} \left[\begin{array}{cc|c} 1 & 0 & \frac{2b_1 + 5b_2}{17} \\ 0 & 1 & \frac{-3b_1 + b_2}{17} \end{array} \right]$$

Therefore, the solution to the general case is $(x_1, x_2) = \left(\frac{2b_1 + 5b_2}{17}, \frac{-3b_1 + b_2}{17} \right)$

And so, for the specific cases:

- 1) $(x_1, x_2) = \left(\frac{2(1) + 5(4)}{17}, \frac{-3(1) + 4}{17} \right) = \left(\frac{13}{17}, \frac{1}{17} \right)$
- 2) $(x_1, x_2) = \left(\frac{2(-2) + 5(5)}{17}, \frac{-3(-2) + 5}{17} \right) = \left(\frac{16}{17}, \frac{11}{17} \right)$

3.8.4.2 Determine the conditions on b , if any, in order to guarantee that the linear system is consistent.

$$\begin{cases} x_1 + 3x_2 = b_1 \\ -2x_1 + x_2 = b_2 \end{cases}$$

$$\left[\begin{array}{cc|c} 1 & 3 & b_1 \\ -2 & 1 & b_2 \end{array} \right] \xrightarrow{R_2+2R_1} \left[\begin{array}{cc|c} 1 & 3 & b_1 \\ 0 & 7 & b_2+2b_1 \end{array} \right] \xrightarrow{\frac{1}{7}R_2} \left[\begin{array}{cc|c} 1 & 3 & b_1 \\ 0 & 1 & \frac{b_2+2b_1}{7} \end{array} \right] \xrightarrow{R_1-3R_2} \left[\begin{array}{cc|c} 1 & 0 & \frac{b_1-3b_2}{7} \\ 0 & 1 & \frac{b_2+2b_1}{7} \end{array} \right]$$

There are no conditions. The system is consistent for all values of b_1 and b_2 .

3.8.5 Another “determining the conditions” problem:

$$\begin{cases} x_1 - 2x_2 - x_3 = b_1 \\ -4x_1 + 5x_2 + 2x_3 = b_2 \\ -4x_1 + 7x_2 + 4x_3 = b_3 \end{cases}$$

$$\left[\begin{array}{ccc|c} 1 & -2 & -1 & b_1 \\ -4 & 5 & 2 & b_2 \\ -4 & 7 & 4 & b_3 \end{array} \right] \xrightarrow[R_3+4R_1]{R_2+4R_1} \left[\begin{array}{ccc|c} 1 & -2 & -1 & b_1 \\ 0 & -3 & -2 & b_2+4b_1 \\ 0 & -1 & 0 & b_3+4b_1 \end{array} \right] \xrightarrow{-\frac{1}{3}R_2} \left[\begin{array}{ccc|c} 1 & -2 & -1 & b_1 \\ 0 & 1 & \frac{2}{3} & \frac{-b_2-4b_1}{3} \\ 0 & 0 & -\frac{2}{3} & \frac{b_3+4b_1}{3} \end{array} \right]$$

$$\xrightarrow{-\frac{3}{2}R_3} \left[\begin{array}{ccc|c} 1 & -2 & -1 & b_1 \\ 0 & 1 & \frac{2}{3} & \frac{-b_2-4b_1}{3} \\ 0 & 0 & 1 & \frac{-b_3-4b_1}{2} \end{array} \right]$$

Therefore, the system is consistent for all values of b_1 , b_2 , and b_3 .

3.8.6 Triangular and Diagonal Matrices

3.8.6.1 Find A^2

$$1) A = \begin{bmatrix} 1 & 0 \\ 0 & -2 \end{bmatrix}$$

$$\begin{aligned} A^2 &= \begin{bmatrix} 1 & 0 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -2 \end{bmatrix} \\ &= \begin{bmatrix} 1(1) + 0(0) & 1(0) + 0(-2) \\ 0(1) + (-2)(0) & 0(0) + (-2)(-2) \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 4 \end{bmatrix} \end{aligned}$$

$$2) A = \begin{bmatrix} -6 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$

$$\begin{aligned} A^2 &= \begin{bmatrix} -6 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 5 \end{bmatrix} \begin{bmatrix} -6 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 5 \end{bmatrix} \\ &= \begin{bmatrix} (-6)(-6) + (0)(0) + (0)(0) & (-6)(0) + (0)(3) + (0)(0) & (-6)(0) + (0)(0) + (0)(5) \\ (0)(-6) + (3)(0) + (0)(0) & (0)(0) + (3)(3) + (0)(0) & (0)(0) + (3)(0) + (0)(5) \\ (0)(-6) + (0)(0) + (5)(0) & (0)(0) + (0)(3) + (5)(0) & (0)(0) + (0)(0) + (5)(5) \end{bmatrix} \\ &= \begin{bmatrix} 36 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 25 \end{bmatrix} \end{aligned}$$

3.8.6.2 Find A^{-k} , such that k is some nonzero constant

$$A = \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & -4 & 0 & 0 \\ 0 & 0 & -3 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}$$

$$\begin{aligned} A^{-k} &= \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & -4 & 0 & 0 \\ 0 & 0 & -3 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}^{-k} \\ &= \begin{bmatrix} 2^{-k} & 0 & 0 & 0 \\ 0 & (-4)^{-k} & 0 & 0 \\ 0 & 0 & (-3)^{-k} & 0 \\ 0 & 0 & 0 & 2^{-k} \end{bmatrix} \end{aligned}$$

4. Determine whether each matrix is symmetric or not.

$$\begin{bmatrix} -8 & -8 \\ 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -1 \\ 1 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 0 & -7 \\ -7 & 7 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 4 \\ 4 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 & 2 \\ 1 & 5 & -6 \\ 2 & 6 & 6 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -1 & 3 \\ -1 & 5 & 1 \\ 3 & 1 & 7 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 & 1 \\ 0 & 2 & 0 \\ 3 & 0 & 0 \end{bmatrix}$$

Symmetric

$$\begin{bmatrix} 0 & -7 \\ -7 & 7 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 4 \\ 4 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -1 & 3 \\ -1 & 5 & 1 \\ 3 & 1 & 7 \end{bmatrix}$$

Not symmetric

$$\begin{bmatrix} 2 & -1 \\ 1 & 2 \end{bmatrix}$$

$$\begin{bmatrix} -8 & -8 \\ 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 & 2 \\ 1 & 5 & -6 \\ 2 & 6 & 6 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 & 1 \\ 0 & 2 & 0 \\ 3 & 0 & 0 \end{bmatrix}$$

3.8.6.3 Find a diagonal matrix A that satisfies the given condition

$$1) A^5 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

$$\begin{aligned} A &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}^{\frac{1}{5}} \\ &= \begin{bmatrix} 1^{\frac{1}{5}} & 0 & 0 \\ 0 & (-1)^{\frac{1}{5}} & 0 \\ 0 & 0 & (-1)^{\frac{1}{5}} \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix} \end{aligned}$$

$$2) A^{-2} = \begin{bmatrix} 9 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{aligned} A &= \begin{bmatrix} 9 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-\frac{1}{2}} \\ &= \begin{bmatrix} 9^{-\frac{1}{2}} & 0 & 0 \\ 0 & 4^{-\frac{1}{2}} & 0 \\ 0 & 0 & 1^{-\frac{1}{2}} \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{3} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

3.8.7 Determinants and Triangular Matrices (08/29)

3.8.7.1 What is C_{32}

$$A = \begin{bmatrix} 2 & 3 & -1 & 1 \\ -3 & 2 & 0 & 3 \\ 3 & -2 & 1 & 0 \\ 3 & -2 & 1 & 4 \end{bmatrix}$$

$$\begin{aligned} C_{32} &= (-1)^{3+2} \begin{vmatrix} 2 & -1 & 1 \\ -3 & 0 & 3 \\ 3 & 1 & 0 \end{vmatrix} \\ &= - \begin{vmatrix} 2 & -1 & 1 \\ -3 & 0 & 3 \\ 3 & 1 & 0 \end{vmatrix} \\ &= - \left(2 \begin{vmatrix} 0 & 3 \\ 1 & 0 \end{vmatrix} - (-1) \begin{vmatrix} -3 & 3 \\ 3 & 0 \end{vmatrix} + 1 \begin{vmatrix} -3 & 0 \\ 3 & 1 \end{vmatrix} \right) \\ &= - (2(-3) - (-1)(-9) + 1(-3)) \\ &= -(-6 + 9 - 3) \\ &= 0 \end{aligned}$$

3.8.7.2 Find all values of λ such that $|A| = 0$

$$A = \begin{bmatrix} \lambda - 2 & 1 \\ -5 & \lambda + 4 \end{bmatrix}$$

$$\begin{aligned} \det(A) &= (\lambda - 2)(\lambda + 4) - (-5)(1) \\ &= \lambda^2 + 2\lambda - 8 + 5 \\ &= \lambda^2 + 2\lambda - 3 \\ &= (\lambda + 3)(\lambda - 1) \\ &= 0 \end{aligned}$$

Therefore, $\lambda = -3, 1$

3.8.7.3 For the matrix $\begin{bmatrix} 3 & 0 & 0 \\ 2 & -1 & 5 \\ 1 & 9 & -4 \end{bmatrix}$ find the determinant 3 different ways with cofactor expansion. Pick different rows and columns each time.

$$\begin{aligned} \det(A) &= 3 \begin{vmatrix} -1 & 5 \\ 9 & -4 \end{vmatrix} - 0 \begin{vmatrix} 2 & 5 \\ 1 & -4 \end{vmatrix} + 0 \begin{vmatrix} 2 & -1 \\ 1 & 9 \end{vmatrix} \\ &= 3(-1(-4) - 5(9)) - 0(2(-4) - 5(1)) + 0(2(9) - (-1)(1)) \\ &= 3(4 - 45) - 0(-8 - 5) + 0(18 + 1) \\ &= 3(-41) - 0(-13) + 0(19) \\ &= 36 \end{aligned}$$

$$\begin{aligned} \det(A) &= 0 \begin{vmatrix} 2 & 5 \\ 9 & -4 \end{vmatrix} - 3 \begin{vmatrix} 3 & 0 \\ 1 & -4 \end{vmatrix} + 0 \begin{vmatrix} 3 & 0 \\ 2 & 5 \end{vmatrix} \\ &= 0(2(-4) - 5(9)) - 3(3(-4) - 0(1)) + 0(3(5) - 0(2)) \\ &= 0(-8 - 45) - 3(-12 - 0) + 0(15 - 0) \\ &= 0(-53) - 3(-12) \\ &= 36 \end{aligned}$$

$$\begin{aligned} \det(A) &= 0 \begin{vmatrix} 2 & -1 \\ 9 & -4 \end{vmatrix} - 0 \begin{vmatrix} 3 & 0 \\ 1 & -4 \end{vmatrix} + 3 \begin{vmatrix} 3 & 0 \\ 2 & -1 \end{vmatrix} \\ &= 0(2(-4) - (-1)(9)) - 0(3(-4) - 0(1)) + 3(3(-1) - 0(2)) \\ &= 0(-8 + 9) - 0(-12 - 0) + 3(-3 - 0) \\ &= 0(1) - 0(-12) + 3(-3) \\ &= 0 + 0 - 9 \\ &= 36 \end{aligned}$$

3.8.7.4 Evaluate $\det(A)$ by a cofactor expansion along a row or column of your choice

$$A = \begin{bmatrix} 1 & k & k^2 \\ 1 & k & k^2 \\ 1 & k & k^2 \end{bmatrix}$$

$$\begin{aligned} \det(A) &= 1 \begin{vmatrix} k & k^2 \\ k & k^2 \end{vmatrix} - k \begin{vmatrix} 1 & k^2 \\ 1 & k^2 \end{vmatrix} + k^2 \begin{vmatrix} 1 & k \\ 1 & k \end{vmatrix} \\ &= 1(k^2 - k^2) - k(1(k^2) - k^2(1)) + k^2(1(k) - k(1)) \\ &= 0 \end{aligned}$$

3.8.7.5 Evaluate the determinant of the following matrices by just looking at them.

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\det(A) = 1(-1)(1) = -1$$

$$A = \begin{bmatrix} 1 & 2 & 7 & -3 \\ 0 & 1 & -4 & 1 \\ 0 & 0 & 2 & 7 \\ 0 & 0 & 0 & 3 \end{bmatrix}$$

$$\det(A) = 1(1)(2)(3) = 6$$

3.8.7.6 Show that the value of the determinant is independent of θ

$$A = \begin{vmatrix} \sin \theta & \cos \theta & 0 \\ -\cos \theta & \sin \theta & 0 \\ \sin \theta - \cos \theta & \sin \theta + \cos \theta & 1 \end{vmatrix}$$

$$\begin{aligned} \det(A) &= \sin \theta \begin{vmatrix} \sin \theta & 0 \\ \sin \theta + \cos \theta & 1 \end{vmatrix} - \cos \theta \begin{vmatrix} \cos \theta & 0 \\ \sin \theta + \cos \theta & 1 \end{vmatrix} \\ &\quad + 0 \begin{vmatrix} \cos \theta & \sin \theta \\ \sin \theta + \cos \theta & \sin \theta \end{vmatrix} \\ &= \sin \theta (\sin \theta(1) - 0(\sin \theta + \cos \theta)) - \cos \theta (\cos \theta(1) - 0(\sin \theta + \cos \theta)) \\ &\quad + 0 (\cos \theta(\sin \theta) - \sin \theta(\sin \theta + \cos \theta)) \\ &= \sin^2 \theta - \cos^2 \theta \\ &= 1 \end{aligned}$$

3.8.8 Row operations and Determinants (08/31)

3.8.8.1 Find the determinant of $\begin{bmatrix} 1 & -3 & 0 \\ -2 & 4 & 1 \\ 5 & -2 & 2 \end{bmatrix}$ WITHOUT using cofactor expansion

$$\begin{aligned}\det(A) &= \begin{vmatrix} 1 & -3 & 0 \\ -2 & 4 & 1 \\ 5 & -2 & 2 \end{vmatrix} \\ &= \begin{vmatrix} 1 & -3 & 0 \\ 0 & -2 & 1 \\ 0 & 13 & 2 \end{vmatrix} \\ &= \begin{vmatrix} 1 & -3 & 0 \\ 0 & -2 & 1 \\ 0 & 0 & \frac{28}{2} \end{vmatrix} \\ &= 1(-2)\left(\frac{28}{2}\right) \\ &= -28\end{aligned}$$

3.8.8.2 Find the determinant of $\begin{bmatrix} 2 & 1 & 3 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 2 & 1 & 0 \\ 0 & 1 & 2 & 3 \end{bmatrix}$

$$\begin{aligned} \det(A) &= \begin{vmatrix} 2 & 1 & 3 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 2 & 1 & 0 \\ 0 & 1 & 2 & 3 \end{vmatrix} \\ &= \begin{vmatrix} 2 & 1 & 3 & 1 \\ 0 & -2 & -5 & -1 \\ 0 & 2 & 1 & 0 \\ 0 & 1 & 2 & 3 \end{vmatrix} \\ &= \begin{vmatrix} 2 & 1 & 3 & 1 \\ 0 & -2 & -5 & -1 \\ 0 & 0 & -4 & -1 \\ 0 & 0 & -3 & 2 \end{vmatrix} \\ &= 2(-2)(-4)(2) \\ &= 64 \end{aligned}$$

3.8.9 Adjoints and Cramer's Rule (09/05)

3.8.9.1 Find the inverse of $A = \begin{bmatrix} 2 & 5 & 5 \\ -1 & -1 & 0 \\ 2 & 4 & 3 \end{bmatrix}$ using the adjoint method

$$\begin{aligned}\det(A) &= 2 \begin{vmatrix} -1 & 0 \\ 4 & 3 \end{vmatrix} - 5 \begin{vmatrix} -1 & 0 \\ 2 & 3 \end{vmatrix} + 5 \begin{vmatrix} -1 & -1 \\ 2 & 4 \end{vmatrix} \\ &= 2(-3) - 5(-3) + 5(-2) \\ &= -6 + 15 - 10 \\ &= -1\end{aligned}$$

$$\begin{aligned}\text{adj}(A) &= \begin{bmatrix} (-1)^{1+1} \begin{vmatrix} -1 & 0 \\ 4 & 3 \end{vmatrix} & (-1)^{1+2} \begin{vmatrix} -1 & 0 \\ 2 & 3 \end{vmatrix} & (-1)^{1+3} \begin{vmatrix} -1 & -1 \\ 2 & 4 \end{vmatrix} \\ (-1)^{2+1} \begin{vmatrix} 5 & 5 \\ 4 & 3 \end{vmatrix} & (-1)^{2+2} \begin{vmatrix} 2 & 5 \\ 2 & 3 \end{vmatrix} & (-1)^{2+3} \begin{vmatrix} 2 & 5 \\ 2 & 4 \end{vmatrix} \\ (-1)^{3+1} \begin{vmatrix} 5 & 5 \\ -1 & 0 \end{vmatrix} & (-1)^{3+2} \begin{vmatrix} 2 & 5 \\ -1 & 0 \end{vmatrix} & (-1)^{3+3} \begin{vmatrix} 2 & 5 \\ -1 & -1 \end{vmatrix} \end{bmatrix} \\ &= \begin{bmatrix} (-1)(3) & -(-1)(3) & -4 + 2 \\ -(15 - 20) & 6 - 10 & -(8 - 10) \\ 5 & -5 & -2 + 5 \end{bmatrix}^T \\ &= \begin{bmatrix} -3 & 3 & -2 \\ 5 & -4 & 2 \\ 5 & -5 & 3 \end{bmatrix}^T \\ &= \begin{bmatrix} -3 & 5 & 5 \\ 3 & -4 & -5 \\ -2 & 2 & 3 \end{bmatrix} \\ \therefore A^{-1} &= - \begin{bmatrix} -3 & 5 & 5 \\ 3 & -4 & -5 \\ -2 & 2 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 3 & -5 & -5 \\ -3 & 4 & 5 \\ 2 & -2 & -3 \end{bmatrix}\end{aligned}$$

3.8.9.2 Solve the following system of equations using Cramer's Rule

$$\begin{cases} 4x + 5y = 2 \\ 11x + y + 2z = 3 \\ x + 5y + 2z = 1 \end{cases} \rightarrow \begin{vmatrix} 4 & 5 & 0 \\ 11 & 1 & 2 \\ 1 & 5 & 2 \end{vmatrix} \rightarrow 4 \begin{vmatrix} 1 & 2 \\ 5 & 2 \end{vmatrix} - 5 \begin{vmatrix} 11 & 2 \\ 1 & 2 \end{vmatrix} = -132$$

$$\begin{aligned} \det(x) &= \begin{vmatrix} 2 & 5 & 0 \\ 3 & 1 & 2 \\ 1 & 5 & 2 \end{vmatrix} \\ &= 2 \begin{vmatrix} 1 & 2 \\ 5 & 2 \end{vmatrix} - 5 \begin{vmatrix} 3 & 2 \\ 1 & 2 \end{vmatrix} \\ &= 2(2 - 10) - 5(6 - 2) \\ &= -16 - 20 \\ &= -36 \end{aligned}$$

$$\begin{aligned} \det(y) &= \begin{vmatrix} 4 & 2 & 0 \\ 11 & 3 & 2 \\ 1 & 1 & 2 \end{vmatrix} \\ &= 4 \begin{vmatrix} 3 & 2 \\ 1 & 2 \end{vmatrix} - 2 \begin{vmatrix} 11 & 2 \\ 1 & 2 \end{vmatrix} \\ &= 4(6 - 2) - 2(22 - 2) \\ &= 16 - 40 \\ &= -24 \end{aligned}$$

$$\begin{aligned} \det(z) &= \begin{vmatrix} 4 & 5 & 2 \\ 11 & 1 & 3 \\ 1 & 5 & 1 \end{vmatrix} \\ &= 4 \begin{vmatrix} 1 & 3 \\ 5 & 1 \end{vmatrix} - 5 \begin{vmatrix} 11 & 3 \\ 1 & 3 \end{vmatrix} + 2 \begin{vmatrix} 11 & 1 \\ 1 & 5 \end{vmatrix} \\ &= 4(1 - 15) - 5(33 - 3) + 2(55 - 1) \\ &= -56 - 150 + 108 \\ &= -98 \end{aligned}$$

Therefore, the solution $(x, y, z) = \left(\frac{3}{11}, \frac{2}{11}, -\frac{49}{66}\right)$

4 Chapter 5: Eigenvectors and Eigenvalues

4.0.1 Eigenvalues and Eigenvectors (11/06)

If A is an $n \times n$ matrix, then a non-zero vector \mathbf{x} , in R^n , is called an eigenvector of A if $A\mathbf{x}$ is a scalar multiple of \mathbf{x} ; that is $A\mathbf{x} = \lambda\mathbf{x}$ for some scalar λ . This scalar λ is called an eigenvalue of A and \mathbf{x} is said to be an eigenvector corresponding to λ .

See, normally, multiplying a vector by a square matrix changes both the magnitude and the direction of the vector. Really screws it up.

Some examples:

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}$$

$$\begin{bmatrix} 5 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 5 \\ 4 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \end{bmatrix}$$

$$\begin{bmatrix} 7 & 8 \\ -2 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 23 \\ 4 \end{bmatrix}$$

However, there are some ways to get consistent results.

4.1 Examples

4.1.1 $\vec{x} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ is an eigenvector of $A = \begin{bmatrix} 3 & -2 \\ 1 & 0 \end{bmatrix}$ because

$$A\vec{x} = \begin{bmatrix} 3 & -2 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 2 \end{bmatrix} = 2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = 2\vec{x} \therefore \lambda = 2$$

4.1.2 Let $A = \begin{bmatrix} 1 & 6 \\ 5 & 2 \end{bmatrix}$, $\vec{u} = \begin{bmatrix} 6 \\ -5 \end{bmatrix}$, $\vec{v} = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$. Are \vec{u} and \vec{v} eigenvectors of A ?

$$A\vec{u} = \begin{bmatrix} 1 & 6 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} 6 \\ -5 \end{bmatrix} = \begin{bmatrix} 1(6) + 6(-5) \\ 5(6) + 2(-5) \end{bmatrix} = \begin{bmatrix} -24 \\ 20 \end{bmatrix} = -4 \begin{bmatrix} 6 \\ -5 \end{bmatrix} \therefore \lambda = -4$$

$$A\vec{v} = \begin{bmatrix} 1 & 6 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} 3 \\ -2 \end{bmatrix} = \begin{bmatrix} 1(3) + 6(-2) \\ 5(3) + 2(-2) \end{bmatrix} = \begin{bmatrix} -9 \\ 11 \end{bmatrix} \neq \lambda \vec{v}$$

4.2 Eigenvector Homework Problem (11/06)

Confirm by multiplication that \mathbf{x} is an eigenvector of A , and find the corresponding eigenvalue.

4.2.1 $A = \begin{bmatrix} 4 & 0 & 1 \\ 2 & 3 & 2 \\ 1 & 0 & 4 \end{bmatrix}; \mathbf{x} = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$

$$A\mathbf{x} = \begin{bmatrix} 4 & 0 & 1 \\ 2 & 3 & 2 \\ 1 & 0 & 4 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 4(1) + 0(2) + 1(1) \\ 2(1) + 3(2) + 2(1) \\ 1(1) + 0(2) + 4(1) \end{bmatrix} = \begin{bmatrix} 5 \\ 10 \\ 5 \end{bmatrix} = 5 \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \therefore \lambda = 5$$