

Gwinnett School of Math, Science, and Technology

Multivariable Calculus Yearlong Notes

Anish Goyal
1st Period

Donny Thurston
Educator

2023-2024



Table of Contents

1 Chapter 1: Systems of Linear Equations and Matrices	4
1.1 Matrix Operations	4
1.1.1 Addition & Subtraction	4
1.1.2 Scalar Multiplication	4
1.1.3 Matrix Multiplication	4
1.1.4 Properties of Matrix Arithmetic	5
1.1.5 Examples	5
1.2 Transpose of a Matrix	6
1.2.1 Transpose Matrix Properties	6
1.3 Homework — “Matrix Stuff” (08/03/2023)	7
1.3.1 Suppose that A, B, C, D and E are matrices with the following sizes:	7
1.3.2 Consider the matrices	7
2 Intro to Systems	9
2.1 Review: Solve the following systems	10
2.1.1 Consistent	10
2.1.2 Inconsistent	10
2.2 The Augmented Matrix	11
2.3 Elementary Row Operations	11
2.3.1 Example 1... again	11
2.4 Connection to Matrices	11
2.4.1 Example 2: again	12
2.4.2 Example 3: again	12
2.4.3 Example 4: Solve the following system	12
2.4.4 Elementary Row Operations & REF Homework Problem (08/08/2023)	13
2.5 Gaussian Elimination	13
2.5.1 Examples	14
2.6 Gaussian Elimination With Back-Substitution	14
2.6.1 Goal:	14
2.6.2 Gaussian Elimination Homework Problem (08/09/2023)	15
2.7 Gauss-Jordan Elimination	16
2.7.1 Goal:	16
2.8 Matrix Properties, Equations, and Inverses	16
2.8.1 With Real Numbers	16
2.8.2 With Matrices	16
2.8.2.1 Multiply:	16
2.8.3 Matrix Inverses	17
3 Chapter 2: Determinants	18
3.1 Prior Knowledge:	18

3.2	Minors & Cofactors	18
3.2.1	Example	18
3.3	Cofactor Expansion	19
3.3.1	Example	19
3.3.2	Does the method generalize to 2×2 matrices?	20
3.3.3	Find the determinant of a 4×4	20
3.4	Theorem	20
3.4.1	Example	21
3.5	Triangular Matrices	21
3.5.1	An important definitions	21

1 Chapter 1: Systems of Linear Equations and Matrices

1.1 Matrix Operations

- Matrix operations are given as: rows x columns
- Two matrices are equal \Leftrightarrow they have the same dimensions and values

1.1.1 Addition & Subtraction

Two matrices can be added/subtracted \Leftrightarrow they have the same dimensions.

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} + \begin{bmatrix} 7 & 0 & 1 \\ -1 & 0 & 2 \end{bmatrix} = \begin{bmatrix} 8 & 2 & 4 \\ 3 & 5 & 8 \end{bmatrix}$$

1.1.2 Scalar Multiplication

- Scalar multiplication is defined as multiplying each element of a matrix by a number

$$3 \begin{bmatrix} 2 & 1 \\ 5 & 2 \end{bmatrix} = \begin{bmatrix} 6 & 3 \\ 15 & 6 \end{bmatrix}$$

1.1.3 Matrix Multiplication

- We can **only** multiply an $(m \times n)$ by $(n \times p)$ matrix.
- The resulting matrix will be $(m \times p)$

1.1.4 Properties of Matrix Arithmetic

- (a) $A + B = B + A$ (**Commutative law for addition**)
- (b) $A + (B + C) = (A + B) + C$ (**Associative law for addition**)
- (c) $A(BC) = (AB)C$ (**Associative law for multiplication**)
- (d) $A(B + C) = AB + AC$ (**Left distributive law**)
- (e) $(B + C)A = BA + CA$ (**Right distributive law**)
- (f) $A(B - C) = AB - AC$
- (g) $(B - C)A = BA - CA$
- (h) $a(B+C) = aB + aC$
- (i) $a(B-C) = aB - aC$
- (j) $(a+b)C = aC + bC$
- (k) $(a-b)C = aC - bC$
- (l) $a(bC) = (ab)C$
- (m) $a(BC) = (aB)C = B(aC)$

1.1.5 Examples

1.

$$\begin{aligned} & \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \\ &= \begin{bmatrix} 1 \cdot 1 + 2 \cdot 3 & 1 \cdot 2 + 2 \cdot 4 \\ 3 \cdot 1 + 4 \cdot 3 & 3 \cdot 2 + 4 \cdot 4 \end{bmatrix} \\ &= \begin{bmatrix} 7 & 10 \\ 15 & 22 \end{bmatrix} \end{aligned}$$

2.

$$\begin{aligned} & \begin{bmatrix} 2 & -3 \\ 5 & 0 \\ -2 & 4 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} -1 \\ 3 \end{bmatrix} \\ &= \begin{bmatrix} 2 \cdot (-1) + (-3) \cdot 3 \\ 5 \cdot (-1) + 0 \cdot 3 \\ -2 \cdot (-1) + 4 \cdot 3 \\ 1 \cdot (-1) + 2 \cdot 3 \end{bmatrix} \\ &= \begin{bmatrix} -11 \\ -5 \\ 14 \\ 5 \end{bmatrix} \end{aligned}$$

3.

$$\begin{aligned} & \begin{bmatrix} 4 & 5 & -1 \end{bmatrix} \begin{bmatrix} 8 \\ 0 \\ 2 \end{bmatrix} \\ &= [4 \cdot 8 + 5 \cdot 0 + (-1) \cdot 2] \\ &= [30] \end{aligned}$$

1.2 Transpose of a Matrix

The transpose of an $(m \times n)$ matrix is the $(n \times m)$ matrix where the rows and columns are swapped.

$$\text{If } B = \begin{bmatrix} 4 & 2 \\ -1 & 0 \\ 3 & 5 \end{bmatrix}, B^T = \begin{bmatrix} 4 & -1 & 3 \\ 2 & 0 & 5 \end{bmatrix}$$

$$\begin{aligned} B \cdot B^T &= \begin{bmatrix} 4 & 2 \\ -1 & 0 \\ 3 & 5 \end{bmatrix} \begin{bmatrix} 4 & -1 & 3 \\ 2 & 0 & 5 \end{bmatrix} \\ &= \begin{bmatrix} 4 \cdot 4 + 2 \cdot 2 & 4 \cdot (-1) + 2 \cdot 0 & 4 \cdot 3 + 2 \cdot 5 \\ (-1) \cdot 4 + 0 \cdot 2 & (-1) \cdot (-1) + 0 \cdot 0 & (-1) \cdot 3 + 0 \cdot 5 \\ 3 \cdot 4 + 5 \cdot 2 & 3 \cdot (-1) + 5 \cdot 0 & 3 \cdot 3 + 5 \cdot 5 \end{bmatrix} \\ &= \begin{bmatrix} 20 & -4 & 22 \\ -4 & 1 & -3 \\ 22 & -3 & 34 \end{bmatrix} \end{aligned}$$

- The transpose of a matrix is **always** multiplicative with the original.
- There is also a **main diagonal** that is the diagonal from the top left to the bottom right, but only square matrices have these.
- The **trace** of a square matrix A is equal to the sum of all the elements on the main diagonal: $\text{tr}(A)$

1.2.1 Transpose Matrix Properties

- $(A^T)^T = A$
- $(A + B)^T = A^T + B^T$
- $(A - B)^T = A^T - B^T$
- $(kA)^T = kA^T$
- $(AB)^T = B^T A^T$

1.3 Homework — “Matrix Stuff” (08/03/2023)

1.3.1 Suppose that A, B, C, D and E are matrices with the following sizes:

A	B	C	D	E
(3×2)	(2×3)	(3×3)	(3×2)	(2×3)

For each matrix operation, sort them into undefined if the operation can't be done, or defined if it can along with the correct dimensions of the outcome.

Undefined	Defined; (4×2)	Defined; (5×5)	Defined; (5×2)
BA	$AC + D$	$E(A + B)$	$(A^T + E)D$
$AB + B$			$E(AC)$
$E^T A$			
$AE + B$			

1.3.2 Consider the matrices

$$A = \begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix}, B = \begin{bmatrix} 4 & -1 \\ 0 & 2 \end{bmatrix}, C = \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix}, D = \begin{bmatrix} 1 & 5 & 2 \\ -1 & 0 & 1 \\ 3 & 2 & 4 \end{bmatrix}, E = \begin{bmatrix} 6 & 1 & 3 \\ -1 & 1 & 2 \\ 4 & 1 & 3 \end{bmatrix}$$

In each part, compute the given expression (where possible).

2. $2A^T + C$

$$\begin{aligned} 2A^T + C &= 2 \begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix}^T + \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix} \\ &= 2 \begin{bmatrix} 3 & -1 & 1 \\ 0 & 2 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix} \\ &= \begin{bmatrix} 6 & -2 & 2 \\ 0 & 4 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix} \\ &= \begin{bmatrix} 7 & 2 & 4 \\ 3 & 5 & 7 \end{bmatrix} \end{aligned}$$

3. $B^T + 5C^T$

$$\begin{aligned}
 B^T + 5C^T &= \begin{bmatrix} 4 & -1 \\ 0 & 2 \end{bmatrix}^T + 5 \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix}^T \\
 &= \begin{bmatrix} 4 & 0 \\ -1 & 2 \end{bmatrix} + 5 \begin{bmatrix} 1 & 3 \\ 4 & 1 \\ 2 & 5 \end{bmatrix} \\
 &= \begin{bmatrix} 4 & 0 \\ -1 & 2 \end{bmatrix} + \begin{bmatrix} 5 & 15 \\ 20 & 5 \\ 10 & 25 \end{bmatrix} \\
 &= \text{Undefined}
 \end{aligned}$$

4. $2E^T - 3D^T$

$$\begin{aligned}
 2E^T - 3D^T &= 2 \begin{bmatrix} 6 & 1 & 3 \\ -1 & 1 & 2 \\ 4 & 1 & 3 \end{bmatrix}^T - 3 \begin{bmatrix} 1 & 5 & 2 \\ -1 & 0 & 1 \\ 3 & 2 & 4 \end{bmatrix}^T \\
 &= 2 \begin{bmatrix} 6 & -1 & 4 \\ 1 & 1 & 1 \\ 3 & 2 & 3 \end{bmatrix} - 3 \begin{bmatrix} 1 & -1 & 3 \\ 5 & 0 & 2 \\ 2 & 1 & 4 \end{bmatrix} \\
 &= \begin{bmatrix} 12 & -2 & 8 \\ 2 & 2 & 2 \\ 6 & 4 & 6 \end{bmatrix} - \begin{bmatrix} 3 & -3 & 9 \\ 15 & 0 & 6 \\ 6 & 3 & 12 \end{bmatrix} \\
 &= \begin{bmatrix} 9 & -5 & -1 \\ -13 & 2 & -4 \\ 0 & 1 & -6 \end{bmatrix}
 \end{aligned}$$

5. $\text{tr}(DE)$

$$\begin{aligned}
 \text{tr}(DE) &= \text{tr} \left(\begin{bmatrix} 1 & 5 & 2 \\ -1 & 0 & 1 \\ 3 & 2 & 4 \end{bmatrix} \begin{bmatrix} 6 & 1 & 3 \\ -1 & 1 & 2 \\ 4 & 1 & 3 \end{bmatrix} \right) \\
 &= \text{tr} \left(\begin{bmatrix} 1 \cdot 6 + 5 \cdot (-1) + 2 \cdot 4 & 1 \cdot 1 + 5 \cdot 1 + 2 \cdot 1 & 1 \cdot 3 + 5 \cdot 2 + 2 \cdot 3 \\ (-1) \cdot 6 + 0 \cdot (-1) + 1 \cdot 4 & (-1) \cdot 1 + 0 \cdot 1 + 1 \cdot 1 & (-1) \cdot 3 + 0 \cdot 2 + 1 \cdot 3 \\ 3 \cdot 6 + 2 \cdot (-1) + 4 \cdot 4 & 3 \cdot 1 + 2 \cdot 1 + 4 \cdot 1 & 3 \cdot 3 + 2 \cdot 2 + 4 \cdot 3 \end{bmatrix} \right) \\
 &= \text{tr} \left(\begin{bmatrix} 9 & 8 & 19 \\ -2 & 0 & 0 \\ 32 & 9 & 25 \end{bmatrix} \right) \\
 &= 34
 \end{aligned}$$

2 Intro to Systems

What are we looking for?

Lines: How many possible solutions?

- Infinite solutions
- One solution
- No solutions


Planes: How many possible solutions?

- Infinite solutions
- No solutions

What does linear actually mean?

- The word linear *really* means that you've got equations with variables and **all** of the variables are degree one.
- This means that there is no limit to the number of dimensions in a linear system.

Linear Systems in Three Unknowns



2.1 Review: Solve the following systems

1.
$$\begin{cases} 2x + y = 10 \\ 3x - y = 5 \end{cases}$$

$$5x = 15$$

$$x = 3$$

$$2(3) + y = 10$$

$$6 + y = 10$$

$$y = 4$$

2.
$$\begin{cases} 2x + y = 10 \\ 6x + 3y = 10 \end{cases}$$

$$y = 10 - 2x$$

$$6x + 3(10 - 2x) = 10$$

$$6x + 30 - 6x = 10$$

$$30 = 10. \therefore \text{no solution}$$

3.
$$\begin{cases} 5x - 2y = 4 \\ 15x - 6y = 12 \end{cases}$$

$$0 = 0$$

$$12 = 12. \therefore \text{infinite solutions}$$

2.1.1 Consistent

- A system of equations is **consistent** if it has at least one solution.

2.1.2 Inconsistent

- A system of equations is **inconsistent** if it has no solutions.

2.2 The Augmented Matrix

$$\begin{cases} x - y + 2z = 5 \\ 2x - 2y + 4z = 10 \\ 3x - 3y + 6z = 15 \end{cases} \longrightarrow \left[\begin{array}{ccc|c} 1 & -1 & 2 & 5 \\ 2 & -2 & 4 & 10 \\ 3 & -3 & 6 & 15 \end{array} \right]$$

2.3 Elementary Row Operations

1. Interchange 2 rows
2. Multiply a row by a non-zero constant
3. Add/subtract a multiple of one row to/from another row

Doing these things changes the matrix, but it's the same system!

2.3.1 Example 1... again

$$\begin{cases} 2x + y = 10 \\ 3x - y = 5 \end{cases}$$

$$\begin{aligned} \left[\begin{array}{cc|c} 2 & 1 & 10 \\ 3 & -1 & 5 \end{array} \right] &\xrightarrow{\frac{1}{2}R_1} \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 5 \\ 3 & -1 & 5 \end{array} \right] \xrightarrow{R_2 - 3R_1} \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 5 \\ 0 & -\frac{5}{2} & -10 \end{array} \right] \\ &\xrightarrow{-\frac{2}{5}R_2} \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 5 \\ 0 & 1 & 4 \end{array} \right] \xrightarrow{R_1 - \frac{1}{2}R_2} \left[\begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 4 \end{array} \right] \end{aligned}$$

And so... $x = 3$ and $y = 4$!

2.4 Connection to Matrices

If we can make a system's matrix look like

$$\left[\begin{array}{ccc|c} 1 & 0 & 0 & c_1 \\ 0 & 1 & 0 & c_2 \\ 0 & 0 & 1 & c_3 \end{array} \right],$$

then the solution to the system will be the ordered triple (c_1, c_2, c_3) .

2.4.1 Example 2: again

$$\begin{cases} 2x + y = 10 \\ 6x + 3y = 10 \end{cases}$$

$$\left[\begin{array}{cc|c} 2 & 1 & 10 \\ 6 & 3 & 10 \end{array} \right] \xrightarrow{\frac{1}{2}R_1} \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 5 \\ 6 & 3 & 10 \end{array} \right] \xrightarrow{R_2-6R_1} \left[\begin{array}{cc|c} 1 & \frac{1}{2} & 5 \\ 0 & 0 & -20 \end{array} \right]$$

This is inconsistent, so there is no solution.

2.4.2 Example 3: again

$$\begin{cases} 5x - 2y = 4 \\ 15x - 6y = 12 \end{cases}$$

$$\left[\begin{array}{cc|c} 5 & -2 & 4 \\ 15 & -6 & 12 \end{array} \right] \xrightarrow{\frac{1}{5}R_1} \left[\begin{array}{cc|c} 1 & -\frac{2}{5} & \frac{4}{5} \\ 15 & -6 & 12 \end{array} \right] \xrightarrow{R_2-15R_1} \left[\begin{array}{cc|c} 1 & -\frac{2}{5} & \frac{4}{5} \\ 0 & 0 & 0 \end{array} \right]$$

Since $0 = 0$, there are infinitely many solutions.

2.4.3 Example 4: Solve the following system

$$\begin{cases} x_1 - 2x_2 + x_3 = 0 \\ 2x_2 - 8x_3 = 8 \\ -4x_1 + 5x_2 + 9x_3 = -9 \end{cases}$$

$$\begin{aligned} & \left[\begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 2 & -8 & 8 \\ -4 & 5 & 9 & -9 \end{array} \right] \xrightarrow{R_3+4R_1} \left[\begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 2 & -8 & 8 \\ 0 & -3 & 13 & -9 \end{array} \right] \xrightarrow{R_3+\frac{3}{2}R_2} \left[\begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 2 & -8 & 8 \\ 0 & 0 & -1 & 3 \end{array} \right] \\ & \xrightarrow{\frac{1}{2}R_2} \left[\begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 1 & -4 & 4 \\ 0 & -3 & 13 & -9 \end{array} \right] \xrightarrow{R_1+2R_2} \left[\begin{array}{ccc|c} 1 & 0 & -7 & 8 \\ 0 & 1 & -4 & 4 \\ 0 & 0 & 1 & 3 \end{array} \right] \xrightarrow{\begin{matrix} R_1+7R_3 \\ R_2+4R_3 \end{matrix}} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 29 \\ 0 & 1 & 0 & 16 \\ 0 & 0 & 1 & 3 \end{array} \right] \end{aligned}$$

Therefore the solution to (x_1, x_2, x_3) is $(29, 16, 3)$.

2.4.4 Elementary Row Operations & REF Homework Problem (08/08/2023)

$$\begin{cases} x + y + 2z = 8 \\ -x - 2y + 3z = 1 \\ 3x - 7y + 4z = 10 \end{cases}$$

$$\begin{aligned} & \left[\begin{array}{ccc|c} 1 & 1 & 2 & 8 \\ -1 & -2 & 3 & 1 \\ 3 & -7 & 4 & 10 \end{array} \right] \xrightarrow[\substack{R_2+R_1 \\ R_3-3R_1}]{\substack{R_2+R_1 \\ R_3-3R_1}} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 8 \\ 0 & -1 & 5 & 9 \\ 0 & -10 & -2 & -14 \end{array} \right] \xrightarrow[\substack{-R_2 \\ -R_3}]{\substack{-R_2 \\ -R_3}} \left[\begin{array}{ccc|c} 1 & 1 & 2 & 8 \\ 0 & 1 & -5 & -9 \\ 0 & 10 & 2 & 14 \end{array} \right] \\ & \xrightarrow[\substack{R_1-R_2 \\ R_3-10R_2}]{\substack{R_1-R_2 \\ R_3-10R_2}} \left[\begin{array}{ccc|c} 1 & 0 & 7 & 17 \\ 0 & 1 & -5 & -9 \\ 0 & 0 & 52 & 104 \end{array} \right] \xrightarrow[\substack{1/52 R_3}]{\substack{1/52 R_3}} \left[\begin{array}{ccc|c} 1 & 0 & 7 & 17 \\ 0 & 1 & -5 & -9 \\ 0 & 0 & 1 & 2 \end{array} \right] \xrightarrow[\substack{R_1-7R_3 \\ R_2+5R_3}]{\substack{R_1-7R_3 \\ R_2+5R_3}} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 2 \end{array} \right] \end{aligned}$$

Therefore, the solution to (x, y, z) is $(3, 1, 2)$.

2.5 Gaussian Elimination

Vocabulary: A matrix is in Row Echelon Form (REF) if:

- (a) Any rows of all zeroes are placed at the bottom of the matrix
- (b) All other rows have a leading 1 ("pivot")
- (c) As we move down the matrix, each leading 1 is further to the right than the 1 above it

A matrix is in Row Reduced Echelon Form if the three above conditions are met in addition to:

- (d) Each column with a leading 1 has all other entries in the column as a 0. ("pivot column")

2.5.1 Examples

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 8 \\ 0 & 1 & 0 & 6 & -3 \\ 0 & 0 & 1 & 7 & 10 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

REF? ✓
RREF? ✓

$$\begin{bmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

REF? ✓
RREF? ✗

$$\begin{bmatrix} 1 & 2 & -3 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & -4 \end{bmatrix}$$

REF? ✗
RREF? ✗

2.6 Gaussian Elimination With Back-Substitution

2.6.1 Goal:

To get the augmented matrix in REF

Solve:
$$\begin{cases} x_1 - 2x_2 + 3x_3 = 9 \\ -x_1 + 3x_2 = -4 \\ 2x_1 - 5x_2 + 5x_3 = 17 \end{cases}$$

$$\begin{aligned} & \left[\begin{array}{ccc|c} 1 & -2 & 3 & 9 \\ -1 & 3 & 0 & -4 \\ 2 & -5 & 5 & 17 \end{array} \right] \xrightarrow[R_3-2R_1]{R_2+R_1} \left[\begin{array}{ccc|c} 1 & -2 & 3 & 9 \\ 0 & 1 & 3 & 5 \\ 0 & -1 & -1 & -1 \end{array} \right] \xrightarrow[R_3+R_2]{R_1+2R_2} \left[\begin{array}{ccc|c} 1 & 0 & 9 & 19 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 2 & 4 \end{array} \right] \\ & \xrightarrow{\frac{1}{2}R_3} \left[\begin{array}{ccc|c} 1 & 0 & 9 & 19 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 1 & 2 \end{array} \right] \end{aligned}$$

$$x + 9z = 19$$

$$y + 3z = 5$$

$$z = 2$$

$$\therefore z = 2, y = 5 - 3z, x = 19 - 9z$$

$$z = 2, y = 5 - 3(2), x = 19 - 9(2)$$

$$z = 2, y = -1, x = 1$$

Therefore, the solution (x_1, x_2, x_3) is $(1, -1, 2)$.

2.6.2 Gaussian Elimination Homework Problem (08/09/2023)

$$\begin{cases} -2w + y + z = -3 \\ x + 2y - z = 2 \\ -3w + 2x + 4y + z = -2 \\ -w + x - 4y - 7z = -19 \end{cases}$$

$$\begin{aligned} & \left[\begin{array}{cccc|c} -2 & 0 & 1 & 1 & -3 \\ 0 & 1 & 2 & -1 & 2 \\ -3 & 2 & 4 & 1 & -2 \\ -1 & 1 & -4 & -7 & -19 \end{array} \right] \xrightarrow{R_4} \left[\begin{array}{cccc|c} -1 & 1 & -4 & -7 & -19 \\ 0 & 1 & 2 & -1 & 2 \\ -3 & 2 & 4 & 1 & -2 \\ -2 & 0 & 1 & 1 & -3 \end{array} \right] \xrightarrow{-R_1} \\ & \left[\begin{array}{cccc|c} 1 & -1 & 4 & 7 & 19 \\ 0 & 1 & 2 & -1 & 2 \\ -3 & 2 & 4 & 1 & -2 \\ -2 & 0 & 1 & 1 & -3 \end{array} \right] \xrightarrow{\begin{matrix} R_3+3R_1 \\ R_4+2R_1 \end{matrix}} \left[\begin{array}{cccc|c} 1 & -1 & 4 & 7 & 19 \\ 0 & 1 & 2 & -1 & 2 \\ 0 & -1 & 16 & 22 & 55 \\ 0 & -2 & 9 & 15 & 35 \end{array} \right] \xrightarrow{\begin{matrix} R_1+R_2 \\ R_3+R_2 \\ R_4+2R_2 \end{matrix}} \\ & \left[\begin{array}{cccc|c} 1 & 0 & 6 & 6 & 21 \\ 0 & 1 & 2 & -1 & 2 \\ 0 & 0 & 18 & 21 & 57 \\ 0 & 0 & 13 & 13 & 39 \end{array} \right] \xrightarrow{\frac{1}{18}R_3} \left[\begin{array}{cccc|c} 1 & 0 & 6 & 6 & 21 \\ 0 & 1 & 2 & -1 & 2 \\ 0 & 0 & 1 & \frac{7}{6} & \frac{19}{6} \\ 0 & 0 & 13 & 13 & 39 \end{array} \right] \xrightarrow{\begin{matrix} R_1-6R_3 \\ R_2-2R_3 \\ R_4-13R_3 \end{matrix}} \\ & \left[\begin{array}{cccc|c} 1 & 0 & 0 & -1 & 2 \\ 0 & 1 & 0 & -\frac{10}{3} & -\frac{13}{3} \\ 0 & 0 & 1 & \frac{7}{6} & \frac{19}{6} \\ 0 & 0 & 0 & -\frac{13}{6} & -\frac{13}{6} \end{array} \right] \xrightarrow{-\frac{6}{13}R_4} \left[\begin{array}{cccc|c} 1 & 0 & 0 & -1 & 2 \\ 0 & 1 & 0 & -\frac{10}{3} & -\frac{13}{3} \\ 0 & 0 & 1 & \frac{7}{6} & \frac{19}{6} \\ 0 & 0 & 0 & 1 & 1 \end{array} \right] \xrightarrow{\begin{matrix} R_1+R_4 \\ R_2+\frac{10}{3}R_4 \\ R_3-\frac{7}{6}R_4 \end{matrix}} \\ & \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 3 \\ 0 & 1 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 1 \end{array} \right] \Rightarrow \begin{cases} w = 3 \\ x = -1 \\ y = 2 \\ z = 1 \end{cases} \end{aligned}$$

2.7 Gauss-Jordan Elimination

2.7.1 Goal:

To get the matrix into RREF

$$\text{Solve: } \begin{cases} x_1 - 3x_3 = -2 \\ 3x_1 + x_2 - 2x_3 = 5 \\ 2x_1 + 2x_2 + x_3 = 4 \end{cases}$$

$$\begin{aligned} & \left[\begin{array}{ccc|c} 1 & 0 & -3 & -2 \\ 3 & 1 & -2 & 5 \\ 2 & 2 & 1 & 4 \end{array} \right] \xrightarrow[R_3-2R_1]{R_2-3R_1} \left[\begin{array}{ccc|c} 1 & 0 & -3 & -2 \\ 0 & 1 & 7 & 11 \\ 0 & 2 & 7 & 8 \end{array} \right] \xrightarrow{R_3-2R_2} \left[\begin{array}{ccc|c} 1 & 0 & -3 & -2 \\ 0 & 1 & 7 & 11 \\ 0 & 0 & -7 & -14 \end{array} \right] \\ & \xrightarrow{\frac{-1}{7}R_3} \left[\begin{array}{ccc|c} 1 & 0 & -3 & -2 \\ 0 & 1 & 7 & 11 \\ 0 & 0 & 1 & 2 \end{array} \right] \xrightarrow[R_2-7R_3]{R_1+3R_3} \left[\begin{array}{ccc|c} 1 & 0 & 0 & 4 \\ 0 & 1 & 0 & -3 \\ 0 & 0 & 1 & 2 \end{array} \right] \Rightarrow \begin{cases} x_1 = 4 \\ x_2 = -3 \\ x_3 = 2 \end{cases} \end{aligned}$$

2.8 Matrix Properties, Equations, and Inverses

2.8.1 With Real Numbers

- If $ab = bc$, then $a = c$, if $b \neq 0$
- If $ab = 0$, then $a = 0$ or $b = 0$, or both

2.8.2 With Matrices

- If $AB = AC$, then $B = C$, if A is invertible
- If $AB = [0]$, then $A = [0]$ or $B = [0]$, or both

2.8.2.1 Multiply:

$$\begin{aligned} & \begin{bmatrix} 2 & 3 \\ 3 & 5 \end{bmatrix} \begin{bmatrix} 5 & -3 \\ -3 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 2(5) + 3(-3) & 2(-3) + 3(2) \\ 3(5) + 5(-3) & 3(-3) + 5(2) \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

2.8.3 Matrix Inverses

- If a matrix has an inverse, it is said to be invertible or non-singular.
- If a matrix does not have an inverse, it is said to be singular.
- Every square matrix has a “special number” associated with it called the **determinant**.
- For the 2×2 matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, the determinant is $ad - bc$
- $A^{-1} = \frac{1}{\det A} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$
- When $\det A = 0$, the matrix is singular and has no inverse (since you cannot divide by zero)

Find the inverse of $A = \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}$

$$\begin{aligned} \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}^{-1} &= \frac{1}{\det A} \begin{bmatrix} 2 & -3 \\ -1 & 4 \end{bmatrix} \\ &= \frac{1}{(4)(2) - (3)(1)} \begin{bmatrix} 2 & -3 \\ -1 & 4 \end{bmatrix} \\ &= \frac{1}{5} \begin{bmatrix} 2 & -3 \\ -1 & 4 \end{bmatrix} \\ &= \begin{bmatrix} \frac{2}{5} & -\frac{3}{5} \\ -\frac{1}{5} & \frac{4}{5} \end{bmatrix} \end{aligned}$$

3 Chapter 2: Determinants

3.1 Prior Knowledge:

$$\begin{bmatrix} 10 & -4 \\ -3 & -5 \end{bmatrix} = -50 - = -62$$

$$\begin{aligned} & \begin{bmatrix} 2 & 4 & 3 \\ -1 & 2 & 3 \\ 3 & 0 & -2 \end{bmatrix} \\ & = ((2 \cdot 2 \cdot -2) + (4 \cdot 3 \cdot 3) + (3 \cdot -1 \cdot 0)) - ((3 \cdot 2 \cdot 3) + (0 \cdot 3 \cdot 2) + (-2 \cdot -1 \cdot 4)) \\ & = (-8 + 36 + 0) - (18 + 0 + 8) \\ & = 28 - 26 \\ & = 2 \end{aligned}$$

3.2 Minors & Cofactors

Given a square matrix A, the minor of matrix element a_{ij} , (M_{ij}) is the determinant of the matrix formed by removing the i^{th} row and j^{th} column from matrix A.

The cofactor of matrix element a_{ij} , $C_{ij} = (-1)^{i+j} \cdot M_{ij}$

3.2.1 Example

Let $\det \begin{bmatrix} 2 & 4 & 3 \\ -1 & 2 & 3 \\ 3 & 0 & -2 \end{bmatrix}$. What is the cofactor of element (1, 1)?

Cofactor checkerboard:

$$\begin{bmatrix} + & - & + \\ - & + & - \\ + & - & + \end{bmatrix}$$

$$M_{11} = \begin{vmatrix} 2 & 3 \\ 0 & -2 \end{vmatrix} = -4$$

$$C_{11} = 1 \cdot -4 = -4$$

Find the minor and cofactor of: \ a) $a_{21} = -1$

$$M_{21} = \begin{vmatrix} 4 & 3 \\ 0 & -2 \end{vmatrix} = -8$$

$$C_{21} = 8$$

b) $a_{33} = -2$

$$M_{33} = \begin{vmatrix} 2 & 4 \\ -1 & 2 \end{vmatrix} = 8$$

$$C_{33} = 8$$

3.3 Cofactor Expansion

- 1) Pick a row or column
- 2) Multiply every entry in that row or column by it's corresponding cofactor
- 3) Add those together. That's it

$$A = \begin{bmatrix} 6 & 7 & -1 \\ 0 & 4 & 1 \\ 2 & 5 & -3 \end{bmatrix}$$

$$\begin{aligned} \det(A) &= 6 \begin{vmatrix} 4 & 1 \\ 5 & -3 \end{vmatrix} + 7 \begin{vmatrix} 0 & 1 \\ 2 & -3 \end{vmatrix} + -1 \begin{vmatrix} 0 & 4 \\ 2 & 5 \end{vmatrix} \\ &= 6(-17) + 7(2) + (-1(-8)) \\ &= -102 + 14 + 8 \\ &= -80 \end{aligned}$$

3.3.1 Example

$$A = \begin{bmatrix} 6 & 4 & 2 \\ 5 & -6 & 1 \\ 0 & 3 & 0 \end{bmatrix}$$

$$\begin{aligned} &6 \begin{vmatrix} -6 & 1 \\ 3 & 0 \end{vmatrix} + 4 \begin{vmatrix} 5 & 1 \\ 0 & 0 \end{vmatrix} + 2 \begin{vmatrix} 5 & -6 \\ 0 & 3 \end{vmatrix} \\ &= 6(-3) + 0 + 2(15) \\ &= -18 + 30 \\ &= 12 \end{aligned}$$

3.3.2 Does the method generalize to 2×2 matrices?

$$\begin{aligned} & \begin{vmatrix} 3 & 5 \\ 7 & 2 \end{vmatrix} \\ &= 3|2| - 5|7| \\ &= 6 - 35 \\ &= -29 \end{aligned}$$

The determinant of a 1×1 matrix is... **itself!**

3.3.3 Find the determinant of a 4×4

$$A = \begin{bmatrix} -3 & 2 & 0 & 8 \\ 2 & 1 & 0 & -4 \\ 5 & -2 & 1 & 5 \\ 2 & 3 & 0 & 6 \end{bmatrix}$$

$$\begin{aligned} &= 0 + 0 + \begin{vmatrix} -3 & 2 & 8 \\ 2 & 1 & -4 \\ 2 & 3 & 6 \end{vmatrix} + 0 \\ &= -2 \begin{vmatrix} 2 & 8 \\ 3 & 6 \end{vmatrix} + \begin{vmatrix} -3 & 8 \\ 2 & 6 \end{vmatrix} - \left(-4 \begin{vmatrix} -3 & 2 \\ 2 & 3 \end{vmatrix} \right) \\ &= 24 - 34 - 52 \\ &= -62 \end{aligned}$$

3.4 Theorem

If A is an $n \times n$ matrix, then regardless of which row or column of A is chosen, the number obtained by multiplying the elements in that row or column by their corresponding cofactors is **always the same** and is called the determinant of A .

3.4.1 Example

Find the determinant of $A = \begin{bmatrix} 1 & 0 & 0 & -1 \\ 3 & 1 & 2 & 2 \\ 1 & 0 & -2 & 1 \\ 2 & 0 & 0 & 1 \end{bmatrix}$

$$\begin{aligned} & 1 \cdot \begin{vmatrix} 1 & 0 & -1 \\ 1 & -2 & 1 \\ 2 & 0 & 1 \end{vmatrix} \\ &= \left(-2 \begin{vmatrix} 1 & -1 \\ 2 & 1 \end{vmatrix} \right) \\ &= -6 \end{aligned}$$

3.5 Triangular Matrices

Find the determinant of $A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 2 & 2 \\ 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 4 \end{bmatrix}$

$$\begin{aligned} & \begin{vmatrix} 2 & 2 & 2 \\ 0 & 3 & 3 \\ 0 & 0 & 4 \end{vmatrix} \\ &= 2 \begin{vmatrix} 3 & 3 \\ 0 & 4 \end{vmatrix} \\ &= 2(3 \cdot 4) \\ &= 2 \cdot 12 \\ &= 24 \end{aligned}$$

If A is an $n \times n$ triangular matrix, then $\det(A)$ is equal to the product of the elements along the main diagonal.

3.5.1 An important definitions

Elementary Matrix a matrix that can be obtained from the $n \times n$ identity matrix by performing a single row operation. \

Are the following matrices elementary? 1) $\begin{bmatrix} 1 & 0 \\ -5 & 1 \end{bmatrix} + (R_3 + 5R_1)$ yes 2) $\begin{bmatrix} -5 & 1 \\ 1 & 0 \end{bmatrix} + (R_1 + 5R_2)$...
no