

Pingala Series

Anita Dash

CONTENTS

1	JEE 2019	1
2	Pingala Series	1
3	Power of the Z transform	3

Abstract—This manual provides a simple introduction to Transforms

1 JEE 2019

Let

$$a_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \quad n \geq 1 \quad (1.1)$$

$$b_n = a_{n-1} + a_{n+1}, \quad n \geq 2, \quad b_1 = 1 \quad (1.2)$$

Verify the following using a python code.

1.1

$$\sum_{k=1}^n a_k = a_{n+2} - 1, \quad n \geq 1 \quad (1.3)$$

1.2

$$\sum_{k=1}^{\infty} \frac{a_k}{10^k} = \frac{10}{89} \quad (1.4)$$

1.3

$$b_n = \alpha^n + \beta^n, \quad n \geq 1 \quad (1.5)$$

1.4

$$\sum_{k=1}^{\infty} \frac{b_k}{10^k} = \frac{8}{89} \quad (1.6)$$

Solution: Below is the python code for the question.

```
wget https://github.com/anitadash/EE3900/
blob/main/Pingala/Codes/Section_1.py
```

2 PINGALA SERIES

2.1 The *one sided* Z-transform of $x(n)$ is defined as

$$X^+(z) = \sum_{n=0}^{\infty} x(n)z^{-n}, \quad z \in \mathbb{C} \quad (2.1)$$

2.2 The *Pingala* series is generated using the difference equation

$$x(n+2) = x(n+1) + x(n), \quad x(0) = x(1) = 1, n \geq 0 \quad (2.2)$$

Generate a stem plot for $x(n)$.

Solution: Below is the python code that generates the stem plot of $x(n)$ in fig-2.2.

```
wget https://github.com/anitadash/EE3900/
blob/main/Pingala/Codes/Section_2_2.
py
```




Fig. 2.2

2.3 Find $X^+(z)$.

Solution: Taking the one-sided Z-transform on

both sides of (2.2),

$$\mathcal{Z}^+[x(n+2)] = \mathcal{Z}^+[x(n+1)] + \mathcal{Z}^+[x(n)] \quad (2.3)$$

$$\mathcal{Z}^+[x(n+2)] = z^2 X^+(z) - z^2 \cdot 1 - z \cdot 1 \quad (2.4)$$

$$\mathcal{Z}^+[x(n+1)] = z X^+(z) - z \cdot 1 \quad (2.5)$$

$$z^2 X^+(z) - z^2 \cdot 1 - z \cdot 1 = z X^+(z) - z \cdot 1 + z X^+(z) \quad (2.6)$$

$$(z^2 - z - 1) X^+(z) = z^2 \quad (2.7)$$

$$X^+(z) = \frac{1}{1 - z^{-1} - z^{-2}} \quad (2.8)$$

$$= \frac{1}{(1 - \alpha z^{-1})(1 - \beta z^{-1})}, \quad |z| > \alpha \quad (2.9)$$

2.4 Find $x(n)$.

Solution: Expanding $X^+(z)$ in (2.9) using partial fractions, we get

$$X^+(z) = \frac{1}{(\alpha - \beta) z^{-1}} \left[\frac{1}{1 - \alpha z^{-1}} - \frac{1}{1 - \beta z^{-1}} \right] \quad (2.10)$$

$$= \frac{1}{(\alpha - \beta)} \sum_{n=0}^{\infty} (\alpha^n - \beta^n) z^{-n+1} \quad (2.11)$$

$$= \sum_{n=1}^{\infty} \frac{\alpha^n - \beta^n}{\alpha - \beta} z^{-n+1} \quad (2.12)$$

$$= \sum_{k=0}^{\infty} \frac{\alpha^{k+1} - \beta^{k+1}}{\alpha - \beta} z^{-k} \quad (2.13)$$

where $k := n + 1$. Thus,

$$x(n) = \frac{\alpha^{n+1} - \beta^{n+1}}{\alpha - \beta} u(n) = a_{n+1} u(n) \quad (2.14)$$

2.5 Sketch

$$y(n) = x(n-1) + x(n+1), \quad n \geq 0 \quad (2.15)$$

Solution: Below is the python code that generates the plot of $y(n)$ in fig-2.5.

```
wget https://github.com/anitadash/EE3900/
blob/main/Pingala/Codes/Section_2_5.
py
```

2.6 Find $Y^+(z)$.

Solution: Taking the one-sided Z-transform on




Fig. 2.5

both sides of (2.15),

$$\mathcal{Z}^+[y(n)] = \mathcal{Z}^+[x(n+1)] + \mathcal{Z}^+[x(n-1)] \quad (2.16)$$

$$Y^+(z) = z X^+(z) - z x(0) + z^{-1} X^+(z) + z x(-1) \quad (2.17)$$

$$= \frac{z + z^{-1}}{1 - z^{-1} - z^{-2}} - z \quad (2.18)$$

$$= \frac{1 + 2z^{-1}}{1 - z^{-1} - z^{-2}}, \quad |z| > \alpha \quad (2.19)$$

since $x(n) = 0 \quad \forall n < 0$.

2.7 Find $y(n)$.

Solution: Using (2.9),

$$Y^+(z) = (1 + 2z^{-1}) \sum_{n=0}^{\infty} x(n) z^{-n} \quad (2.20)$$

$$= \sum_{n=0}^{\infty} x(n) z^{-n} + \sum_{n=1}^{\infty} 2x(n-1) z^{-n} \quad (2.21)$$

$$= x(0) + \sum_{n=1}^{\infty} (x(n) + 2x(n-1)) z^{-n} \quad (2.22)$$

α, β are the roots of the equation $z^2 - z - 1$

$$y(n) = \frac{(\alpha^{n+1} - \beta^{n+1}) + (2\alpha^n + 2\beta^n)}{\alpha - \beta} \quad (2.23)$$

$$= \frac{(\alpha^{n+2} - \beta^{n+2}) - \alpha\beta(\alpha^n + \beta^n)}{\alpha - \beta} \quad (2.24)$$

$$= \frac{(\alpha - \beta)(\alpha^{n+1} + \beta^{n+1})}{\alpha - \beta} \quad (2.25)$$

$$= \alpha^{n+1} + \beta^{n+1} \quad (2.26)$$

Thus, $y(n) = \alpha^{n+1} + \beta^{n+1}$ for $n \geq 0$ as $\alpha + \beta = 1$. with the definition of b_n , we see that $y(n) = b_{n+1}$. Hence, $b_n = \alpha^n + \beta^n$.

3 POWER OF THE Z TRANSFORM

3.1 Show that

$$\sum_{k=1}^n a_k = \sum_{k=0}^{n-1} x(n) = x(n) * u(n-1) \quad (3.1)$$

Solution: From (2.14), and noting that $x(n) = 0 \forall n < 0$,

$$\sum_{k=1}^n a_k = \sum_{k=0}^{n-1} x(k) \quad (3.2)$$

$$= \sum_{k=-\infty}^{n-1} x(k) \quad (3.3)$$

$$= \sum_{k=-\infty}^{\infty} x(k) u(n-1-k) \quad (3.4)$$

$$= x(n) * u(n-1) \quad (3.5)$$

3.2 Show that

$$a_{n+2} - 1, \quad n \geq 1 \quad (3.6)$$

can be expressed as

$$[x(n+1) - 1] u(n) \quad (3.7)$$

Solution: From (2.14),

$$a_{n+2} - 1 = [x(n+1) - 1], \quad n \geq 0 \quad (3.8)$$

and so, using the definition of $u(n)$,

$$a_{n+2} - 1 = [x(n+1) - 1] u(n) \quad (3.9)$$

3.3 Show that

$$\sum_{k=1}^{\infty} \frac{a_k}{10^k} = \frac{1}{10} \sum_{k=0}^{\infty} \frac{x(k)}{10^k} = \frac{1}{10} X^+ (10) \quad (3.10)$$

Solution:

$$\sum_{k=1}^{\infty} \frac{a_k}{10^k} = \frac{1}{10} \sum_{k=0}^{\infty} \frac{a_{k+1}}{10^k} \quad (3.11)$$

$$= \frac{1}{10} \sum_{k=0}^{\infty} \frac{x(k)}{10^k} \quad (3.12)$$

$$= \frac{1}{10} X^+(z) \quad (3.13)$$

$$= \frac{1}{10} \times \frac{100}{89} = \frac{10}{89} \quad (3.14)$$

3.4 Show that

$$\alpha^n + \beta^n, \quad n \geq 1 \quad (3.15)$$

can be expressed as

$$w(n) = (\alpha^{n+1} + \beta^{n+1}) u(n) \quad (3.16)$$

and find $W(z)$.

Solution: Putting $n = k + 1$ and using the definition of $u(n)$,

$$\alpha^n + \beta^n = (\alpha^{k+1} + \beta^{k+1}) u(k) \quad (3.17)$$

Hence, (??) can be expressed as

$$w(n) = (\alpha^{n+1} + \beta^{n+1}) u(n) = y(n) \quad (3.18)$$

Therefore,

$$W(z) = Y(z) = \frac{1 + 2z^{-1}}{1 - z^{-1} - z^{-2}} \quad (3.19)$$

3.5 Show that

$$\sum_{k=1}^{\infty} \frac{b_k}{10^k} = \frac{1}{10} \sum_{k=0}^{\infty} \frac{y(k)}{10^k} = \frac{1}{10} Y^+ (10) \quad (3.20)$$

Solution:

$$\sum_{k=1}^{\infty} \frac{b_k}{10^k} = \frac{1}{10} \sum_{k=0}^{\infty} \frac{b_{k+1}}{10^k} \quad (3.21)$$

$$= \frac{1}{10} \sum_{k=0}^{\infty} \frac{y(k)}{10^k} \quad (3.22)$$

$$= \frac{1}{10} Y^+(z) \quad (3.23)$$

$$= \frac{1}{10} \times \frac{120}{89} = \frac{12}{89} \quad (3.24)$$

3.6 Solve the JEE 2019 problem.

Solution: We know that

$$\sum_{k=1}^n a_k = x(n) * u(n-1) \quad (3.25)$$

But

$$x(n) * u(n-1) \stackrel{Z}{\rightleftharpoons} X(z) z^{-1} U(z) \quad (3.26)$$

$$= \frac{z^{-1}}{(1 - z^{-1} - z^{-2})(1 - z^{-1})} \quad (3.27)$$

$$= z \left[\frac{1}{1 - z^{-1} - z^{-2}} - \frac{1}{1 - z^{-1}} \right] \quad (3.28)$$

$$\stackrel{z}{=} z \sum_{n=0}^{\infty} (x(n) - 1) z^{-n} \quad (3.29)$$

$$= \sum_{n=0}^{\infty} (x(n) - 1) z^{-n+1} \quad (3.30)$$

$$= \sum_{n=0}^{\infty} (x(n+1) - 1) z^{-n} \quad (3.31)$$

$$(3.32)$$

From (2.14), we get

$$\sum_{k=1}^n a_k = a_{n+2} - 1 \quad (3.33)$$

We have solved the remaining options in the problems earlier. Therefore, options 1, 2, and 3 are correct and option 4 is incorrect.