



The new global lithological map database GLiM: A representation of rock properties at the Earth surface

Jens Hartmann and Nils Moosdorf

*Institute for Biogeochemistry and Marine Chemistry, KlimaCampus, Universität Hamburg,
Bundesstrasse 55, DE-20146 Hamburg, Germany (geo@hattes.de)*

[1] Lithology describes the geochemical, mineralogical, and physical properties of rocks. It plays a key role in many processes at the Earth surface, especially the fluxes of matter to soils, ecosystems, rivers, and oceans. Understanding these processes at the global scale requires a high resolution description of lithology. A new high resolution global lithological map (GLiM) was assembled from existing regional geological maps translated into lithological information with the help of regional literature. The GLiM represents the rock types of the Earth surface with 1,235,400 polygons. The lithological classification consists of three levels. The first level contains 16 lithological classes comparable to previously applied definitions in global lithological maps. The additional two levels contain 12 and 14 subclasses, respectively, which describe more specific rock attributes. According to the GLiM, the Earth is covered by 64% sediments (a third of which are carbonates), 13% metamorphics, 7% plutonics, and 6% volcanics, and 10% are covered by water or ice. The high resolution of the GLiM allows observation of regional lithological distributions which often vary from the global average. The GLiM enables regional analysis of Earth surface processes at global scales. A gridded version of the GLiM is available at the PANGAEA Database (<http://dx.doi.org/10.1594/PANGAEA.788537>).

Components: 24,000 words, 4 figures, 3 tables.

Keywords: data set; global map; lithology.

Index Terms: 1030 Geochemistry: Geochemical cycles (0330); 3610 Mineralogy and Petrology: Geochemical modeling (1009, 8410); 3617 Mineralogy and Petrology: Alteration and weathering processes (1039).

Received 2 August 2012; **Revised** 8 October 2012; **Accepted** 8 October 2012; **Published** 7 December 2012.

Hartmann, J., and N. Moosdorf (2012), The new global lithological map database GLiM: A representation of rock properties at the Earth surface, *Geochem. Geophys. Geosyst.*, 13, Q12004, doi:10.1029/2012GC004370.

1. Introduction

[2] Lithological information, describing the geochemical and physical properties of rocks, is increasingly in demand in different disciplines of Earth system science. Lithology is a key variable in many fields, including landscape evolution [de Caritat et al., 2012; Coulthard, 2001; Dürr et al., 2011; McAuliffe, 1994], pathways of water fluxes

[Gleeson et al., 2011], river chemical composition [Gibbs, 1967; Hartmann, 2009; Hartmann et al., 2010; Meybeck, 1987], isotope provenance [Bataille and Bowen, 2012; Godfrey et al., 2009; Lacan and Jeandel, 2005], matter supply to ecosystems [Porder et al., 2007] and lateral land ocean matter fluxes [Bernier, 1994; Godderis et al., 2009; Hartmann, 2009; Hartmann and Moosdorf, 2011; Meybeck, 1987; Moosdorf et al., 2011].

[3] Lithology is often highly heterogeneous at the regional and local scale, and even small occurrences of certain rock types can provide important information for matter fluxes at the Earth's surface (e.g., carbonate or evaporite occurrence for river chemical compositions) [e.g., *Mast et al.*, 1990]. Lithological maps should therefore sufficiently represent this heterogeneity.

[4] There are three existing digital global lithological maps, developed by *Bluth and Kump* [1991] (revised by *Gibbs and Kump* [1994]), *Amiotte Suchet et al.* [2003], and *Dürr et al.* [2005]. The first two maps are grid-based representations, based on works of the Ronov group [e.g., *Khain et al.*, 1981], with a resolution of 2° and 1°, respectively, and distinguish 13 and 6 lithological classes. The latter map [*Dürr et al.*, 2005] is a vector-map translated from the second edition of a geological world map at a scale of 1:25,000,000 [*Dottin*, 1990], using additional literature. *Dürr et al.* [2005] distinguished 15 lithological classes (including water and ice) for application in weathering and hydrological studies ("hydrolithology").

[5] The development of regional to global scale lateral/land-ocean matter flux models [*Hartmann et al.*, 2010; *Jansen et al.*, 2010] has highlighted that the existing maps did not sufficiently represent the regional lithological variability in North America [*Moosdorf et al.*, 2010] or highly active weathering areas with small occurrences of relevant rock types [*Hartmann and Moosdorf*, 2011]. Thus, we developed a new global lithological map data set (GLiM) to represent rock properties at a sufficient resolution for the development and calibration of regional to global scale matter flux models. This publication describes the major characteristics of the new map. The map is designed in a way that can be easily enhanced, e.g., by age, tectonic or geochemical information, and can be applied to a wide range of studies.

2. Methods

[6] In this section, we briefly describe the methods applied to translate and assemble the global lithological map (GLiM). A detailed description of the preparation processes, including all individual map sources, is given in Appendix A.

[7] The GLiM was assembled from 92 regional lithological maps of the highest available resolution (target scale: 1:1,000,000). The selection of the geological source maps represents a compromise

between detail (which would be best at a higher resolution), feasibility (high resolution data sets are difficult to handle at global scale) and consistency (all maps at one scale would be elegant). Most of the maps used were national or state geological maps, acquired from their individual developers, converted to a vector-based GIS usable format, and assembled into continental data sets using ESRI ArcGIS 10 software. The amount of processing depended on the original data format: paper maps were scanned and digitized by hand (i.e., each geological border was manually redrawn) whereas less conversion was necessary for maps already available in digital vector format. An accurate description of the lithological distribution requires a topologically sound data set, without holes or overlaps of individual polygons. We resolved more than 120,000 topological errors manually and automatically to correct for overlaps and gaps existing within the source maps and to harmonize borders of neighboring maps (see Appendix A for details).

[8] The descriptions of rocks – for simplicity, the term "rock" encompasses here all types of rocks including unconsolidated sediments – of the individual units were derived either from the geological maps (if available), or from additional literature. In total, 318 literature sources were used to describe the rocks, resulting in ~22,000 unique descriptors in the geodatabase. The geological boundaries between the source maps were not harmonized and rock descriptions were generalized into 16 lithological classes, based on the classification system by *Dürr et al.* [2005] (Table 1). To improve the representation of the regional heterogeneity, two additional levels of information were added as subclasses following *Moosdorf et al.* [2010] (Table 1). These subclasses add information to the first level lithological class (e.g., "siliciclastic sediment"), to describe additional lithological properties (e.g., the dominating grain size) or local special attributes (e.g., mapped coal occurrences or mentioned pyroclastics), which may be useful for analyses of a wide range of processes.

[9] The second level contains 12 and the third level 14 subclasses (Table 1). In the GLiM, a full lithological classification consists therefore of three two-digit codes ("xxyyzz"). The mandatory first level class (xx) represents the dominant lithology. The second (yy) and third (zz) optional level subclasses closer define the rocks and highlight local special attributes. However, yy and zz attributes

Table 1. The Lithological Classes, Subclasses and Literature Values of Previous Lithological Maps for Comparison^a

Code	Description	Global – Literature Values											
		Africa	Antarctica	Asia	Austasi	Europe	North America	South America	Global	Dürr et al. [2005]	Amiotte Suchet et al. [2003]	Gibbs and Kump [1994]	
ev ig mt nd pa pb pi py psc ism ss su va vb vi wb pr Cl	Evaporites	0.6%	0.0%	0.3%	0.5%	0.0%	0.1%	0.3%	0.3%	0.1%			
	Ice and Glaciers	0.0%	88.3%	0.1%	0.0%	0.0%	7.5%	0.1%	8.7%	0.2%			
	Metamorphics	27.6%	6.4%	6.9%	3.0%	13.6%	13.2%	12.6%	13.0%	4.1%	27.5% ^b	20.0% ^b	
	No Data	0.0%	0.0%	0.2%	0.0%	0.0%	0.1%	0.0%	0.1%				
	Acid plutonic rocks	1.1%	0.7%	7.2%	3.6%	10.0%	8.5%	9.7%	5.7%	7.2%			
	Basic plutonic rocks	0.2%	0.5%	1.1%	1.0%	0.8%	0.8%	0.2%	0.7%	0.2%			
	Intermediate plutonic rocks	0.1%	0.0%	0.3%	0.3%	0.5%	0.9%	0.7%	0.4%				
	Pyroclastics	0.0%	0.0%	0.6%	0.2%	0.1%	1.6%	1.4%	0.6%				
	Carbonate sedimentary rocks	9.4%	0.0%	10.0%	5.6%	17.6%	9.0%	1.5%	7.8%	10.4%	13.4%	9.3%	
	Mixed sedimentary rocks	4.6%	1.0%	25.2%	14.2%	22.3%	15.3%	12.3%	14.6%	7.8%			
	Siliclastic sedimentary rocks	16.4%	1.7%	14.9%	13.0%	14.7%	21.6%	25.7%	16.3%	16.3%	51.6% ^c	36.5% ^c	
	Unconsolidated sediments	35.1%	0.0%	24.0%	52.1%	15.3%	12.9%	26.4%	24.6%	29.7%			
	Acid volcanic rocks	0.1%	0.0%	1.4%	1.1%	1.5%	1.6%	1.6%	1.0%	1.0%	2.3%		
	Basic volcanic rocks	3.3%	0.9%	4.2%	2.2%	2.8%	4.0%	4.2%	3.5%	5.8%	5.2%	6.8%	
	Intermediate volcanic rocks	0.6%	0.5%	2.4%	3.1%	0.4%	1.7%	2.5%	1.7%				
	Water Bodies	0.9%	0.0%	1.3%	0.1%	0.4%	1.4%	0.8%	0.9%	0.6%			
	Precambrian rocks									11.5%		27.5%	
	Complex lithology									5.5%			
	First Level												
ad am ds gr la lo mx or pu py sh ss	Alluvial deposits	3.8%	0.0%	2.3%	12.4%	0.0%	3.1%	9.9%	4.1%	15.5%			
	Mafic metamorphics mentioned	0.4%	0.0%	2.3%	0.7%	4.7%	0.7%	4.3%	1.7%				
	Dune sands	17.6%	0.0%	1.9%	12.4%	0.0%	0.0%	0.0%	5.3%	1.5%			
	Greenstone mentioned	0.0%	0.0%	0.1%	0.1%	0.3%	0.1%	0.3%	0.1%				
	Laterites	0.0%	0.0%	0.4%	1.9%	0.1%	0.0%	3.0%	0.6%				
	Loess	0.0%	0.0%	3.1%	0.1%	0.0%	0.6%	0.2%	1.1%	2.6%			
	Mixed grain size	30.8%	2.4%	44.5%	28.7%	52.5%	29.5%	33.9%	33.5%				
	Organic sediment	0.1%	0.0%	0.2%	0.6%	0.6%	0.1%	0.0%	0.2%				
	(Pure) carbonate	11.0%	0.0%	6.2%	6.4%	4.9%	4.5%	1.8%	5.9%				
	Pyroclastics mentioned	1.5%	0.3%	6.4%	6.6%	3.6%	2.8%	5.3%	4.0%				
	Fine grained	1.7%	0.3%	9.1%	7.4%	5.8%	12.8%	5.2%	6.6%		25.4% ^d	12.6% ^d	
	Coarse grained	1.0%	0.0%	4.1%	10.3%	4.7%	6.7%	10.2%	4.7%		26.2% ^d	23.9% ^d	
	Second Level: yy												
	bs ch cl ev fe	Alluvial deposits	3.8%	0.0%	2.3%	12.4%	0.0%	3.1%	9.9%	4.1%	15.5%		
		Mafic metamorphics mentioned	0.4%	0.0%	2.3%	0.7%	4.7%	0.7%	4.3%	1.7%			
		Dune sands	17.6%	0.0%	1.9%	12.4%	0.0%	0.0%	0.0%	5.3%	1.5%		
		Greenstone mentioned	0.0%	0.0%	0.1%	0.1%	0.3%	0.1%	0.3%	0.1%			
		Laterites	0.0%	0.0%	0.4%	1.9%	0.1%	0.0%	3.0%	0.6%			
		Loess	0.0%	0.0%	3.1%	0.1%	0.0%	0.6%	0.2%	1.1%	2.6%		
Mixed grain size		30.8%	2.4%	44.5%	28.7%	52.5%	29.5%	33.9%	33.5%				
Organic sediment		0.1%	0.0%	0.2%	0.6%	0.6%	0.1%	0.0%	0.2%				
(Pure) carbonate		11.0%	0.0%	6.2%	6.4%	4.9%	4.5%	1.8%	5.9%				
Pyroclastics mentioned		1.5%	0.3%	6.4%	6.6%	3.6%	2.8%	5.3%	4.0%				
Fine grained		1.7%	0.3%	9.1%	7.4%	5.8%	12.8%	5.2%	6.6%		25.4% ^d	12.6% ^d	
Coarse grained		1.0%	0.0%	4.1%	10.3%	4.7%	6.7%	10.2%	4.7%		26.2% ^d	23.9% ^d	
Third Level: zz													
bs ch cl ev fe		Black shale mentioned	0.0%	0.0%	0.5%	0.3%	0.6%	0.4%	0.5%	0.3%			
		Chert mentioned	0.1%	0.0%	0.2%	0.9%	1.1%	1.3%	0.2%	0.4%			
		Fossil plant organic material mentioned	0.2%	0.3%	5.1%	3.7%	2.1%	1.8%	4.1%	2.8%			
		Subordinate evaporites mentioned	0.0%	0.0%	8.5%	6.6%	2.8%	1.0%	3.8%	3.8%			
		Reduced-Iron minerals mentioned	0.1%	0.0%	0.1%	0.4%	0.1%	0.4%	0.5%	0.2%			

Table 1. (continued)

Code	Description	Global – Literature Values						
		Africa	Antarctica	Asia	Austasi	Europe	North America	South America
gl	Glacial influence mentioned	0.1%	0.0%	1.7%	0.6%	5.2%	0.9%	2.9%
mt	Metamorphic influence mentioned	0.9%	1.2%	3.8%	3.3%	8.8%	9.0%	4.6%
ph	Phosphorous-rich minerals mentioned	0.0%	0.0%	0.7%	0.0%	0.1%	0.0%	0.0%
pr	Subordinate plutonics mentioned	0.1%	0.0%	1.1%	0.5%	0.0%	0.4%	1.5%
pt	Pyrite mentioned	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%
sr	Subordinate sedimentary rocks mentioned	0.4%	1.2%	1.2%	1.2%	3.2%	0.9%	1.8%
su	Subordinate unconsolidated sediments mentioned	0.0%	0.0%	4.0%	1.0%	2.9%	0.1%	2.6%
vr	Subordinate volcanics mentioned	0.2%	0.8%	4.5%	1.7%	2.6%	2.5%	1.6%
we	Intensive weathering	0.0%	0.0%	1.9%	0.6%	0.0%	0.0%	0.0%

^aThe original classification was adapted to match the current classes where necessary and possible. Values represent the proportion of areal extent.

^bOriginal class: “Shield rocks”, including metamorphics and plutonics.

^cOriginal classes: “Sand and Sandstones” and “Shale” [Amiotte Suchet *et al.*, 2003] as well as “Sandstones” and “Shales” [Gibbs and Kump, 1994]. Thus, the value for Amiotte Suchet *et al.* [2003] includes also unconsolidated sediments.

^dInterpreted for comparison from the original classes “Sand and Sandstone”, “Sandstone”, and “Shale”.

were not set if no special attributes were described in the used source of the lithological information.

3. Results and Discussion

[10] The new global map represents the global distribution of the different lithological classes in an unprecedented high resolution (Figure 1; see auxiliary material for a higher-resolution version of the map).¹ The map scale is on average 1:3,750,000 (area weighted). The assembled map consists in a total of 1,235,400 polygons.

[11] The terrestrial Earth surface is covered by 64% sediments (a third of which are carbonates), 13% metamorphics, 7% plutonics, and 6% volcanics (Table 1). No rock classes were assigned to areas of ice and inland water bodies, which cover 10% of the map area (including Antarctica). Significant regional differences in lithology distribution are observable: e.g., carbonate containing rock units represent 40% of the area in Europe while they represent only 14% in South America (Table 1).

[12] Not all mapped units are provided with second and third level information: 68% and 20% have an yy-subclass or a zz-subclass assigned, respectively. These subclasses allow additional analyses: e.g., 15% of the mapped sedimentary lithological classes (siliciclastic sedimentary rocks (SS), unconsolidated sediments (SU), mixed sedimentary rocks (SM), including carbonate sedimentary rocks (SC) are dominated by fine grained sediments and 8% are dominated by coarse grain sizes. About 9% of these classes represent pure carbonate sedimentary rock without mapped clastic sediments, and 62% are sediments of mixed or not determinable grain size. This example highlights the difficulties in representing the local heterogeneity of lithological properties, despite the high detail for the global data set. Another example for the use of the subclasses are evaporites in Asia, which are rarely dominant and are therefore mapped only in a few areas as lithological class (xx = ev: 0.28%). However, they occur subordinately in other lithological units covering a far larger area (zz = ev: 8.52%).

[13] In total, 437 different combinations of lithological classes (xx) and subclasses (yy, zz) occur. Development of a categorized lithological map is a continuous compromise between accuracy and simplicity. On the one hand, the diversity of the described variables demands as many classes as

¹Auxiliary materials are available in the HTML. doi:10.1029/2012GC004370.




Figure 1. Representation of the global lithological map database (GLiM) showing the basic lithological classes (first level of information). Please note that the map resolution is finer than the print resolution (simplified gridded data available for download at <http://dx.doi.org/10.1594/PANGAEA.788537>).

possible to sufficiently represent individual features, but on the other hand, the number of classes should be minimized to improve usability. To optimize its usage, the GLiM features three class levels, which can be recombined as needed to represent the desired attributes for individual applications. However, the map is still subject to a significant uncertainty considering rock properties of some lithological classes, which is highlighted by the large amount of mixed sediments (xx = sm) at the Earth's surface (14.6%). Despite of that carbonate rocks and siliciclastic rocks are different, considering a large range of physical or chemical properties (e.g., dissolution rates or aquifer characteristics) they occur often “mixed” in one geological unit, undistinguishable at the scale of the here used maps.

[14] The third level subclasses (zz), including metamorphic influence mentioned (mt), subordinate plutonics mentioned (pr), subordinate sedimentary rocks mentioned (sr), subordinate volcanics mentioned (vr), and subordinate unconsolidated sediments mentioned (su), indicate the presence of rock types other than those defining the first level

lithological class. The fact that these subclasses occur in 10% of the global area highlights the heterogeneity of the mapped lithological units and that they should not be considered as being homogeneous. For example, the first level class “basic volcanics (vb),” although dominated by basalt can also contain layers of sediments. Moreover, such mixing effects do not only occur where they can be documented.

[15] This internal heterogeneity, considering needed local simplifications due to mapping at the given scales of source data, is also highlighted by a cross-check with 164,953 point data with information on lithology (see section A5 for details). These show a direct match to the mapped lithological classes of the GLiM in 40% of the cases. However, if further lithological classes of the GLiM, including information of subclasses, are considered to match the point data, 64% of the point information is represented by the GLiM.

[16] Although the general rock composition of the Earth's surface is remarkably similar between the existing digital lithological maps (Table 1), the




Figure 2. Comparison of the GLiM with previous lithological world maps for an example region in the eastern Himalayas. The scale of the detail lithological maps is 1:3,000,000.

representation of the rock distribution varies largely between them. Particularly the complex geological settings have been greatly improved in the GLiM compared to the previous maps (Figure 2), as it was previously shown for North America [Moosdorf *et al.*, 2010]. The improvement, visually observable in Figure 2, is also documented by a comparison with point information on lithology from geochemical databases (see section A5 for details). The ‘ground-truth’ evaluation of the GLiM results in 40% accurate data points (64% including ‘similar’ lithologies), whereas the lithological map of Dürr *et al.* [2005] accurately represents 30% of the data points, and the lithological maps of Amiotte Suchet *et al.* [2003] and Bluth and Kump [1991] correspond to 22% and 23% of the data, respectively. However, a direct comparison of the maps is difficult due to the broader classification system

of the older maps, as for example the class “complicated lithology,” which is resolved now to a more detailed classification in the GLiM. The design of the GLiM allows future inclusion of more detailed regional maps as well as additional information, e.g., age information of the lithological units or their tectonic history in future versions.

4. Summary

[17] The distribution of rock-attributes and rocktypes at a very high resolution (“average” scale 1:3,750,000) can represent the heterogeneity of the rocks at the Earth’s surface. Strong differences in the lithological cover of the different continents are observable in the new global lithological map (GLiM) (Table 1).

[18] The GLiM allows the assessment of global scale research questions at high resolution and thus helps to advance our knowledge of Earth surface processes. The resolution facilitates comparison of results, e.g., of different regional matter flux studies. The regional differences between the properties of lithology could be used to better parameterize globally comparable regional models. In addition, the application of GLiM-derivates of distinct coarser resolution can be used to analyze scaling effects, which might be useful if such derivates are used in Earth system models. In addition, the architecture of GLiM allows more details or levels of information to be added and could thus be developed further in accordance with specific scientific questions. A gridded version of the GLiM for global scale studies is provided in the PANGAEA database: <http://dx.doi.org/10.1594/PANGAEA.788537>.

Appendix A: Sources, Methods, Definitions

A1. Sources of the GLiM

[19] The Global Lithological Map (GLiM) was assembled from geological maps with a target resolution of 1:1 000 000 and ideally with a national extent or larger. However, if no suitable national geological map of the required quality was available, either coarser maps were used (e.g., in Africa), or finer maps assembled (e.g., the combined state geological maps of the preliminary geological map database of the conterminous U.S.A.). Table A1 lists all sources of the GLiM.

A2. Geographical Combination Methods

[20] To translate the geological maps into lithological maps and aggregate them, they needed to be in a standardized form. The preparation of each map differed according to the original format (original formats are provided in Table A1). The workflow is described in the following and in Figure A1:

[21] 1. Paper maps were scanned as pixel images using an A0-capable scanner at a resolution of 300 DPI.

[22] 2. Pixel images were transformed into shapefiles by manually clicking all the lines representing geological features and creating a shapefile in ArcEditor. Usually, the geological boundaries were cut from a mantling polygon to reduce topology errors. If possible, the image was converted to a black/white image before digitalization, to enable

the use of the ArcScan tools (included in the ESRI ArcGIS package), which increase digitalization speed.

[23] 3. Vector PDF-files were converted into files of '.dxf'-format, and imported as line-features into ArcGIS, georeferenced, and converted to polygons using the ArcGIS tool "Line to Polygon." Those polygons were attributed by hand, and checked for errors due to the import procedure. In Somalia, the georeferencing of the original vector-pdf was not feasible with geographical transformations offered in ArcGIS. That map was rubber sheeted to the country borders after georeferencing, to reduce the spatial error.

[24] 4. MapInfo files were converted into shapefiles using the EVTools GIS and imported in ArcGIS.

[25] 5. Shapefiles were imported in an ESRI-file geodatabase and projected to the ECKERT IV projection (WGS84 ellipsoid as predefined in ArcGIS 10). The geological map of the Canaries needed to be manually spatially adjusted to its correct position, which was done based on the global shoreline data set [Wessel and Smith, 1996].

[26] The geometry of the shapefiles of some of the maps was corrupted, e.g., with "self intersections" (as well as some original data sets). Before they could be further processed, the geometry of the shape was repaired with the "repair geometry" tool (ArcGIS) and the results cross-checked with the original data. The areas that were particularly affected were Bolivia, Brazil, Chile, Indonesia, and Somalia.

[27] In some maps (e.g., in Bangladesh, Venezuela), large polygons represented the coastal ocean and larger rivers. The river part of these polygons was retained but the ocean part was erased. The border between retained river data and the erased ocean data was defined manually using satellite imagery (MS Bing® maps implemented in ArcGIS Online). Some of the original maps contained a large number of very small topological errors, which were first solved automatically by clustering polygon vertexes closer than a certain threshold apart. Affected areas and the thresholds were: Brazil (50 m), Somalia (500 m), Soviet Union (20 m).

[28] Some of the source maps contained holes for larger water bodies. The larger of these holes were filled with polygons and attributed as water body by hand; smaller holes were filled and merged automatically to the neighboring polygon sharing the longest border. Areas affected by that procedure are e.g., Brazil, Denmark, Indonesia, and Ireland.



Table A1. Sources of the GLiM as Well as Formats and Scale of the Map Source

Country (Region)	Map Sources	Lithology Sources	Original Format	Scale (1:X mil)
Africa	<i>Persits et al.</i> [2002]	<i>Africa</i> <i>Alberti et al.</i> [1999], <i>Auchapt et al.</i> [1987], <i>Bow et al.</i> [2009], <i>Egyptian Geological Survey</i> [1981], <i>Furman and Graham</i> [1994], <i>Geological and Mineral Resources Department</i> [1981], <i>Geological Survey and Mines Department</i> [1973], <i>Hotin and Ouedraogo</i> [1976], <i>Kazmin</i> [1972], <i>Marzoli et al.</i> [2000], <i>Michel</i> [1973], <i>Nehlig et al.</i> [2006], <i>Németh et al.</i> [2003], <i>Persits et al.</i> [2002], <i>Saadi</i> [1982], <i>Schlüter</i> [2005], and <i>Wright</i> [1965] <i>ESRI</i> [2008]	Shapefile	7.5
Cape Verde	<i>Economic and Social Research Institute (ESRI)</i> [2008]		Shapefile	10 ^a
Madagascar	<i>Besairie</i> [1964]	<i>Besairie</i> [1964]	Mapinfo	1
Mozambique	<i>Pinna et al.</i> [1987]	<i>Pinna et al.</i> [1987]	Mapinfo	1
Namibia	<i>Geological Survey of Namibia</i> [1980]	<i>Geological Survey of Namibia</i> [1980]	Shapefile	1
Somalia	<i>Africover</i> [2002]	<i>Africover</i> [2002] and <i>Nyagah</i> [1995]	Vector PDF	1
South Africa	<i>Keyser</i> [1998]	<i>Keyser</i> [1998]	Shapefile	1
Spain (Canaries)	<i>Instituto Geológico y Minero de Espana</i> [1994]	<i>Instituto Geológico y Minero de Espana</i> [1994]	Shapefile	1
Antarctica	Scientific Committee on Antarctic Research (Antarctic digital database, 2012) and <i>Tingey</i> [1991a]	<i>Antarctica</i> Scientific Committee on Antarctic Research (Antarctic digital database, 2012) and <i>Tingey</i> [1991a, 1991b]	Shapefile	10
Andaman Islands Arab Peninsula	<i>Steinshouer et al.</i> [1999] <i>Pollastro et al.</i> [1997a]	<i>Asia</i> <i>Allen et al.</i> [2008] <i>Abed and Amireh</i> [1999], <i>Al-Hafidh and Qasim</i> [1992], <i>Alsharhan and Nairn</i> [1994, 1995, 1997], <i>Alsharhan et al.</i> [1993], <i>As-Saruri et al.</i> [2010], <i>Hegner and Pallister</i> [1989], <i>Jassim and Goff</i> [2006], <i>Lustrino and Sharkov</i> [2006], <i>McClure</i> [1978], <i>Moshirif</i> [1984], <i>Natural Resources Project</i> [1990], <i>Pollastro et al.</i> [1997a], <i>Qari</i> [1989], <i>Sharief</i> [1982], and <i>Soliman and Elfetouh</i> [1970] <i>Alam et al.</i> [2003] and <i>Persits et al.</i> [2001] <i>CCOP</i> [2004] and <i>Fromaget and Saurin</i> [1952]	Shapefile Shapefile	5 4.5
Bangladesh Cambodia	<i>Persits et al.</i> [2001] <i>Coordinating Committee for Geoscience Programmes in East and Southeast Asia (CCOP)</i> [2004]		Shapefile Shapefile	1 2
China China Nei Mongol	<i>China Geological Survey</i> [2001] <i>Ministry of Geology and Mineral Resources of the People's Republic of China</i> [1991]	<i>China Geological Survey</i> [2001] <i>Ministry of Geology and Mineral Resources of the People's Republic of China</i> [1991]	Shapefile Shapefile	2.5 1.5

Table A1. (continued)

Country (Region)	Map Sources	Lithology Sources	Original Format	Scale (1:X mil)
China Xinjiang	Bureau of Geology and Mineral Resources of Xinjiang [1992]	Bureau of Geology and Mineral Resources of Xinjiang [1992]	Shapefile	1.5
Himalaya	Geological Survey of India [2005]	Ahmad and Bhat [1987], Ahmad and Tarney [1991], Ahmad and Bhat [2006], Bhat et al. [1994a], Balasubrahmanyam [2006], Bhat et al. [1994b], Bhat et al. [1994b], Dunlap and Wyszczanski [2002], Fuchs [1975], Ganesan et al. [1982], Geological Survey of India [1979, 2005], Hambrey et al. [1981], Joshi et al. [1990], Kumar and Singh [1983], Macfarlane et al. [1999], Radhakrishna et al. [1984], Sabir Khan [1994], Singh and Jain [2003], Sinha [1989], Sinha and Nautiyal [1981], Sinha et al. [1999], Srimal [2005], M. J. Streule et al. [2009], and Valdiya and Bhatia [1980]	Paper map	1
India	Dasgupta and Chakravorty [1998]	Bhatia and Bhatia [1973], Bhushan [1998], Bhushan et al. [1991], Central Ground Water Board [2009], Clark [2005], Dasgupta and Chakravorty [1998], Deb et al. [1978], Deb et al. [2002], Dunlap and Wyszczanski [2002], Fuchs and Sinha [1978], Geological Survey of India [2009], Ghose et al. [2010], Krishnan [1968], Kumar et al. [1984], Kumar and Pankaj [2009], Kumar [1985], Lakhera et al. [1980], Mazumdar and Bhattacharya [2004], Mishra [2009], Mukherjee et al. [1992], Myrow et al. [2006], Nag et al. [1999], Nayak et al. [2009], Pande and Kumar [1965], Patwardhan [2010], Rana et al. [2005], Sharma and Bhole [2005], Sharma [2007], Sharma and Thomas [2005], Shekhawat and Prabhulingaiah [2010], Singh [2010], Singh [2011], M. J. Streule et al. [2009], Sundaram et al. [2001], Sur et al. [2006], Taylor and Mitchell [2000], Tewari and Seckbach [2011], Thakur [1998], Tobgay et al. [2009], Valdiya and Jhingran [1973], Valdiya [1995], and Wadia [1975]	Paper map	2
Iran	Pollastro et al. [1997b]	Alavi [1994], Alsharhan and Nairn [1997], Boccaletti et al. [1976], Farhoudi and Karig [1977], Fürsich et al. [2009], Mehrabi et al. [1999], Moores and Fairbridge [1997], Pollastro et al. [1997b], Seyed-Emami [2003], and Wendt et al. [2005] Geological Survey of Japan [2003] and Takai et al. [1963]	Shapefile	2.5
Japan	Geological Survey of Japan [2003]		Shapefile	1



Table A1. (continued)

Country (Region)	Map Sources	Lithology Sources	Original Format	Scale (1:X mil)
Korea	CCOP [2004]	CCOP [2004], Egawa and Lee [2009], Kim and Lee [1996], Lee and Chough [2006], Moores and Fairbridge [1997], and Sim and Lee [2006]	Shapefile	2
Laos	CCOP [2004]	CCOP [2004] and Fromaget and Saurin [1952]	Shapefile	2
Malaysia	CCOP [2004]	Balaguru and Nichols [2004], CCOP [2004], Cocks et al. [2005], Heng [1992], Metcalfe [1990], Wang and Sugiyama [2002], and Williams et al. [1988]	Shapefile	2
Maldives	GADM database of global administrative areas (version 1.0, 2009)	Moores and Fairbridge [1997]	Shapefile	1 ^a
Mid East (used for Pakistan and Afghanistan)	Haghipour and Saidi [2010]	Doeblich and Wahl [2006], Haghipour and Saidi [2010], and Khan et al. [1964]	Shapefile	5
Mongolia	Steinshouer et al. [1999]	Bayasgalan et al. [2007], Dashzeveg et al. [1995], Hanzl and Krejci [2005], Jahn et al. [2000], Mineral Resources Authority of Mongolia [1998], Steinshouer et al. [1999], and Traynor and Sladen [1995]	Shapefile	5
Myanmar	CCOP [2004]	CCOP [2004], Earth Sciences Research Division [1977], Helmcke [1985], Khin and Myitta [1999], Latt et al. [2008], Loveman [1919], Mitchell [1992], and Myanmar Ministry of Science and Technology (Sample questions and worked out examples for Geol-03042 Engineering Geology for Mining Engineering, 2010)	Shapefile	2
Sri Lanka	Economic and Social Commission for Asia and the Pacific [1989]	Economic and Social Commission for Asia and the Pacific [1989]	Shapefile	1
South Asia (used for parts of Nepal and Bhutan)	Wandrey and Law [1998]	Castelli and Lombardo [1988], Corrie and Kohn [2011], Corvinus and Rimal [2001], Dietrich and Gansser [1981], Long et al. [2011], Rai et al. [2007], and Wandrey and Law [1998]	Shapefile	10
Soviet Union (former)	Karpinsky [1983]	Dolginow et al. [1994], Karpinsky [1983], and Zhamojda [1968]	Paper map	2.5
Thailand	CCOP [2004]	CCOP [2004] and Dheeradilok et al. [1992]	Shapefile	2
Turkey	Institute of Mineral Research and Exploration [1961]	Institute of Mineral Research and Exploration [1961], Okay et al. [2002, 2008], and Yilmaz [1993]	Mapinfo	0.5
Vietnam	CCOP [2004]	CCOP [2004], Fromaget and Saurin [1952], and Lepvrier et al. [2011]	Shapefile	2
Yemen	Natural Resources Project [1990]	Alsharhan and Naim [1997], Natural Resources Project [1990], Purser [1998], and Yemen Geological Survey and Mineral Resources Board [2009]	Shapefile	1



Table A1. (continued)

Country (Region)	Map Sources	Lithology Sources	Original Format	Scale (1:X mil)
Queensland	<i>Whitaker et al.</i> [2007]	<i>Australasia</i>	Shapefile	1
New South Wales	<i>Raymond et al.</i> [2007c]	<i>Whitaker et al.</i> [2007]	Shapefile	1
Tasmania	<i>Raymond et al.</i> [2007b]	<i>Raymond et al.</i> [2007b]	Shapefile	1
South Australia	<i>Raymond et al.</i> [2007a]	<i>Raymond et al.</i> [2007a]	Shapefile	1
Western Australia	<i>Whitaker et al.</i> [2008]	<i>Whitaker et al.</i> [2008]	Shapefile	1
Northern Territory	<i>Stewart et al.</i> [2008]	<i>Stewart et al.</i> [2008]	Shapefile	1
Brunei	<i>Liu et al.</i> [2006]	<i>Liu et al.</i> [2006]	Shapefile	1
Fiji	<i>CCOP</i> [2004]	<i>CCOP</i> [2004] and <i>Heng</i> [1992]	Shapefile	2
	<i>Colley</i> [1976]	<i>Bradshaw</i> [1992], <i>Colley</i> [1976, 2009], and <i>Rodda</i> [1967]	Mapinfo	0.25
Indonesia	<i>Geological Survey Institute of Indonesia</i> [1993]	<i>Geological Survey Institute of Indonesia</i> [1993]	Shapefile	1
Malaysia Sabah	<i>Yin</i> [1985]	<i>Yin</i> [1985]	Mapinfo	0.5
Malaysia Sarawak	<i>Heng</i> [1992]	<i>Heng</i> [1992]	Mapinfo	0.5
New Caledonia	<i>Direction de l'Industrie, des Mines et de l'Energie (DIMENC)</i> [1981]	<i>DIMENC</i> [1981]	Shapefile	0.2
New Zealand	<i>New Zealand Geological Survey</i> [1972]	<i>New Zealand Geological Survey</i> [1972]	Shapefile	1
Papua New Guinea	<i>CCOP</i> [2004]	<i>CCOP</i> [2004] and <i>Stead</i> [1990]	Shapefile	2
Philippines	<i>CCOP</i> [2004]	<i>CCOP</i> [2004], <i>Mitchell et al.</i> [1986], and <i>Moore and Fairbridge</i> [1997]	Shapefile	2
Solomon Islands	<i>Steinshouer et al.</i> [1999] and <i>Turner</i> [1978]	<i>Peterson et al.</i> [1999] and <i>Turner</i> [1978]	Mapinfo	5 (0.1)
Vanuatu	<i>Mollock</i> [1974]	<i>Mollock</i> [1974]	Mapinfo	1
Austria	<i>Egger et al.</i> [1999]	<i>Europe</i>	Shapefile	1.5
Balkan	<i>Pawlewicz et al.</i> [1997]	<i>Dinter and Royden</i> [1993], <i>Grubić</i> [1980], <i>Iancu et al.</i> [2005], <i>Pawlewicz et al.</i> [1997], <i>Robertson and Shallo</i> [2000], <i>Schefer et al.</i> [2011], <i>Sotiropoulos et al.</i> [2008], and <i>Szabó et al.</i> [1992]	Shapefile	5
Belgium	One Geology Europe Consortium (Surface geological maps of Europe, 2010, available at http://www.onegeology-europe.org/ , accessed 17 January 2011) (hereinafter referred to as One Geology Europe Consortium 2010)	One Geology Europe Consortium 2010	Shapefile	1
Britain	<i>British Geological Survey</i> [2007]	<i>Archer</i> [1977], <i>Brenchley and Rawson</i> [2007], <i>British Geological Survey</i> [2007], <i>Cook</i> [1995], <i>Cope et al.</i> [1992], <i>Cox et al.</i> [1992], <i>Crimes et al.</i> [1992], <i>Davies</i> [1967], <i>Douglas and Arkell</i> [1928], <i>Gallois and Etches</i> [2001], <i>Gallois</i> [1976], <i>Greensmith</i> [1957], <i>Hallam and Selwood</i> [1976], <i>Tunbridge</i> [1981],	Shapefile	0.65



Table A1. (continued)

Country (Region)	Map Sources	Lithology Sources	Original Format	Scale (1:X mil)
Czech Republic	One Geology Europe Consortium 2010	<i>John and Fisher</i> [1984], <i>Macdougall and Prentice</i> [1964], <i>Milodowski and Zalasiewicz</i> [1991], <i>Peter</i> [1986], <i>Pharaoh and Carney</i> [2000], <i>Edwards</i> [1976], <i>Rushon</i> [1979], <i>Smith and Edwards</i> [1991], <i>Watts</i> [1962], <i>Williams and Floyd</i> [2000], <i>Wright and Knight</i> [1995], and <i>Wright</i> [1856]	Shapefile	1
Denmark	One Geology Europe Consortium 2010	One Geology Europe Consortium 2010	Shapefile	1
Estonia	One Geology Europe Consortium 2010	One Geology Europe Consortium 2010	Shapefile	1
France	<i>Bureau de Recherches Géologiques et Minières (BRGM)</i> [2003]	<i>BRGM</i> [2003]	Shapefile	1
Germany	<i>Trumit et al.</i> [2003]	<i>Trumit et al.</i> [2003]	Shapefile	1
Hungary	One Geology Europe Consortium 2010	One Geology Europe Consortium 2010	Shapefile	1
Ireland	<i>Geological Survey of Ireland</i> [2006]	<i>Geological Survey of Ireland</i> [2006]	Shapefile	0.5
Italy	One Geology Europe Consortium 2010	One Geology Europe Consortium 2010	Shapefile	1
Luxemburg	One Geology Europe Consortium 2010	One Geology Europe Consortium 2010	Shapefile	1
Netherlands	One Geology Europe Consortium 2010	One Geology Europe Consortium 2010	Shapefile	1
Northern Europe	<i>Sigmond</i> [2002]	<i>Sigmond</i> [2002]	Shapefile	4
Poland	One Geology Europe Consortium 2010	One Geology Europe Consortium 2010	Shapefile	1
Portugal	One Geology Europe Consortium 2010	One Geology Europe Consortium 2010	Shapefile	1
Portugal Islands	<i>Billett and Cowden</i> [1980] and GADM database of global administrative areas (version 1.0, 2009)	<i>Billett and Cowden</i> [1980] and <i>Diirr et al.</i> [2005]	Paper map/shapefile	1
Slovak Republic	One Geology Europe Consortium 2010	One Geology Europe Consortium 2010	Shapefile	1
Slovenia	One Geology Europe Consortium 2010	One Geology Europe Consortium 2010	Shapefile	1
(former) Soviet Union	<i>Karpinsky</i> [1983]	<i>Dolginow et al.</i> [1994], <i>Karpinsky</i> [1983], and <i>Zhamojda</i> [1968]	Paper map	2.5
Spain	<i>Instituto Geológico y Minero de Espana</i> [1994]	<i>Instituto Geológico y Minero de Espana</i> [1994]	Shapefile	1
Sweden	<i>Pawlewicz et al.</i> [1997]	<i>Manten</i> [1971]	Shapefile	5
Switzerland	<i>Bundesamt für Landestopografie</i> [2005]	<i>Bundesamt für Landestopografie</i> [2005], <i>Hsü and Briegel</i> [1991], <i>Walter</i> [1995], and <i>Weissert and Stössel</i> [2010]	Shapefile	0.5
Alaska	<i>Moll et al.</i> [1997]	North America	Shapefile	2.5
Canada	<i>Wheeler et al.</i> [1997]	<i>Beikman</i> [1980]	Shapefile	5
		<i>Fyffe and Richard</i> [2007], <i>Ludington et al.</i> [2005], <i>Ministère des Ressources Naturelles</i> [2002], <i>Nicholson et al.</i> [2006], and <i>Wheeler et al.</i> [1997]		
		<i>Buschkuhle</i> [2003], <i>Douglas et al.</i> [1974, 1970], <i>Frazier and Schwimmer</i> [1987], <i>Hamblin</i> [1997], <i>Hamilton et al.</i> [2004], <i>Hayes et al.</i> [1994],		
Canada - Alberta	<i>Hamilton et al.</i> [2004]		Shapefile	1



Table A1. (continued)

Country (Region)	Map Sources	Lithology Sources	Original Format	Scale (1:X mil)
Canada – Baffin Island	Scott and de Kemp [1998]	Irish [1971], Macdonald and Slimmon [1999], Massey et al. [2005], McMechan and Dawson [1995], Okulich [2006], Ollerenshaw [1970], Pollock et al. [2000], Powers [1931], Pruett and Murray [1991], Richardson et al. [1990], and Yeo et al. [2002]	Shapefile	1
Canada – British Columbia	Massey et al. [2005]	Blackadar et al. [1968], Johns and Young [2006], and Scott and de Kemp [1998]	Shapefile	0.25
Canada - Manitoba	Schledewitz and Lindal [2005]	Massey et al. [2005]	Shapefile	0.25
Canada - Newfoundland	Colman-Sadd and Crisby-Whittle [2005]	Schledewitz and Lindal [2005]	Shapefile	0.1
Canada – North West Territories	Irwin [2005]	Colman-Sadd and Crisby-Whittle [2005]	Shapefile	0.5
Canada Nova Scotia	Fisher and Poole [2006]	Irwin [2005]	Shapefile	0.5
Canada - Ontario	Ontario Geological Survey [1993]	Fisher and Poole [2006]	Shapefile	1
Canada - Saskatchewan	Macdonald and Slimmon [1999]	Ontario Geological Survey [1993]	Shapefile	1
		Douglas et al. [1970], Fuzesy [1979], Gendzwill and Meieshin [1996], Harlaub et al. [2004], Macdonald and Slimmon [1999], Marsh and Heinemann [2006], and Millard [1996]		
Canada – Slave Province	Stubley [2005]	Stubley [2005]	Shapefile	0.1
Canada – Western Churchill	Paul et al. [2002]	Aspler and Chiarenzelli [1997], Ernst and Buchan [2004], Hadlari and Rainbird [2000], Hadlari et al. [2006], Paul et al. [2002], Schau [1993], and Tella et al. [2007]	Shapefile	1
Canada Yukon	Gordey and Makepeace [2000]	Gordey and Makepeace [2000]	Shapefile	0.25
Conterminous USA	Dicken et al. [2005, 2007], Ludington et al. [2005], Nicholson et al. [2005, 2006, 2007], and Stoeser et al. [2005]	Dicken et al. [2005, 2007], Ludington et al. [2005], Nicholson et al. [2005, 2006, 2007], and Stoeser et al. [2005]	Shapefile	0.5
Costa Rica	Schruben [1996]	Schruben [1996]	Shapefile	0.5
Greenland	Escher and Pulvertaft [1995]	Escher and Pulvertaft [1995] and Henriksen et al. [2009]	Shapefile	2.5
Honduras	Wieczorek et al. [1998]	Wieczorek et al. [1998]	Shapefile	0.5
Mexico	Servicio Geologico Mexicano [2007]	Servicio Geologico Mexicano [2007]	Shapefile	0.5
Middle America General	Garrity and Solter [2009]	Alminas et al. [1994], Arengi and Hodgson [2000], Brown [1913], Brunt et al. [1973], Bureau des Mines et de l'Energie [1988], Candanedo et al. [1998], Christman [1953], Draper et al. [1994, 1996], Frank et al. [1998], Gallienne [1975], Garrity and Solter [2009], Geological Survey Department Jamaica [1958], Gunawan et al. [2008], James and Ginsburg [1979], Kesler et al. [1977], Linas [2005], Mesolella	Shapefile	5



Table A1. (continued)

Country (Region)	Map Sources	Lithology Sources	Original Format	Scale (1:X mil)
Puerto Rico		<i>et al.</i> [1969], <i>Ministerio de Medio Ambiente y Recursos Naturales</i> [2002], <i>Myloie and Carew</i> [1995], <i>Palmer</i> [1945], <i>Schruben</i> [1996], <i>Servicio Geológico Mexicano</i> [2007], and <i>Wieczorek et al.</i> [1998]		
USA Hawaii	<i>Bawiec</i> [1999] <i>Sherrod et al.</i> [2007]	<i>Bawiec</i> [1999] <i>Sherrod et al.</i> [2007]	Shapefile Shapefile	0.1 0.1
Argentina		<i>South America</i>		
Brazil	<i>Servicio Geológico Minero Argentino</i> [1997] <i>Serviço Geológico do Brasil</i> [2004]	<i>Servicio Geológico Minero Argentino</i> [1997] <i>Serviço Geológico do Brasil</i> [2004]	Map Info Shapefile	2.5 1
Bolivia	<i>Paraeija and Ballon</i> [1978]	<i>Paraeija and Ballon</i> [1978]	MapInfo file	1
Chile	<i>Servicio Nacional de Geología y Minería</i> [2004]	<i>Servicio Nacional de Geología y Minería</i> [2002, 2004]	Shapefile	1
Colombia	<i>Gómez et al.</i> [2007]	<i>Gómez et al.</i> [2007]	Paper map	2.8
Ecuador	<i>Ortega et al.</i> [1982]	<i>Ortega et al.</i> [1982]	Paper map	1
French Guyana	<i>BRGM</i> [2001]	<i>BRGM</i> [2001] and <i>Ledru et al.</i> [1994]	Shapefile	0.5
Guyana & Suriname	<i>Schobbenhaus and Bellizia</i> [2001]	<i>Prasad</i> [1983] and <i>Schobbenhaus and Bellizia</i> [2001]	Shapefile	5
Paraguay	<i>González</i> [2000]	<i>Berrocá and Fernandes</i> [1996], <i>Comin-Chiaromonti et al.</i> [1999], <i>Gomes et al.</i> [1996], and <i>González</i> [2000]	Pixel image	2.5
Peru	<i>Instituto de Geología y Minería</i> [1975]	<i>Anderson</i> [1944], <i>Bahlburg et al.</i> [2006], <i>Bendezú and Fontboté</i> [2002], <i>Boucot et al.</i> [1980], <i>Bush et al.</i> [1994], <i>Callot et al.</i> [2008], <i>Campbell et al.</i> [2006], <i>Carlotto et al.</i> [2009], <i>Carrascal-Miranda and Suarez-Ruiz</i> [2004], <i>Chen et al.</i> [2010], <i>Clark et al.</i> [1990], <i>DeVries</i> [1988], <i>Fildani et al.</i> [2005], <i>Frizzell</i> [1943], <i>Haeblerlin et al.</i> [2004], <i>Hillebrandt</i> [1970], <i>Instituto de Geología y Minería</i> [1975], <i>Jacay and Sempere</i> [2005], <i>Jacay et al.</i> [2008], <i>Jaillard et al.</i> [2005], <i>Ligarda et al.</i> [1993], <i>Lindell et al.</i> [2010], <i>Mathalone and Montoya</i> [1995], <i>Mégard and Philip</i> [1976], <i>Mégard et al.</i> [1984], <i>Noble et al.</i> [1974, 1979, 1990], <i>Petersen et al.</i> [1977], <i>Portugal</i> [1974], <i>Ramirez and Cisneros</i> [2007], <i>Riccardi et al.</i> [1992], <i>Rod and Maync</i> [1954], <i>Sandeman et al.</i> [1996], <i>Szekely</i> [1966], <i>Taylor et al.</i> [1998],	Shapefiles	1

Table A1. (continued)

Country (Region)	Map Sources	Lithology Sources	Original Format	Scale (1:X mil)
Trinidad and Tobago Uruguay	Schobbenhaus and Bellizia [2001] Dirección Nacional de Minería y Geología [1985]	Tosdal <i>et al.</i> [1981], Tsuchi [1990], Wilson [1963], Winkler <i>et al.</i> [2005], and Zapata <i>et al.</i> [2005] Snoke [2001] Anton [1993], Collazo Carabello [2006], Dirección Nacional de Minería y Geología [1985], Lustrino <i>et al.</i> [2005], Manganeli <i>et al.</i> [2007], Martínez and Rojas [2004], Masquelin <i>et al.</i> [2009], Muzio <i>et al.</i> [2009], Preciozzi Porta [1993], Richards [1974], Tófolo and Morrás [2009], Tófolo and Pazos [2010], Ubilla <i>et al.</i> [2004], Verde and Martínez [2004], and Veroslavsky and Ubilla [2007] Garrity <i>et al.</i> [2006]	Shapefile PDF File	5 0.5
Venezuela	Garrity <i>et al.</i> [2006]		Shapefile	0.75

^aNo geological map was available for these islands, thus other available spatial representations were used and given a homogenous lithology.




Figure A1. Simplified workflow of the GLiM development.

The geological map of Antarctica [Tingey, 1991a] did not include ice areas. The Antarctic digital database (Scientific Committee on Antarctic Research, Antarctic digital database, 2012) was used to represent the ice extent around the geological units.

[29] A few island areas had no geological maps even after intensive research. However, these were all volcanic or carbonate islands. Thus, their shape was taken from administrative data sets, and a uniform lithology was assumed, based on geological literature. Affected areas are: Azores, Cape Verde Islands, Maldives.

[30] In order to build a topologically sound data set, the regional national geological maps were not simply all combined into one map, but their shapes adapted to perfectly match on a per-continent basis.

[31] First, the parts of geological maps which were overlapping with their neighbors were erased (GIS tool), favoring 1) maps of higher resolution and 2) more trusted maps (e.g., available in better format or with more concise description). The erase order per continent is given below. Maps named first were given preference to those named afterwards. If the erased map had greater terrestrial coverage than the favored map, the resulting fragments were deleted manually.

[32] *Africa*. South Africa, Namibia, Mozambique, USGS Africa.

[33] *Asia*. China Neii Mongol, China Xinjiang, Japan, China, Korea, Vietnam, Laos, Cambodia, Thailand, Malaysia, Himalaya, Sri Lanka, India, Myanmar, Bangladesh, South Asia, Yemen, Turkey, former Soviet Union, Iran, Arab Peninsula, Mid East, Mongolia, Maldives, Andaman Islands.

[34] *Antarctica*. No classical border conflicts were encountered in Antarctica. However, the geological map, which did not represent ice areas, was erased from a map representing those areas.

[35] *Australia and Oceania*. Australia-NSW, Australia-NT, Australia-QLD, Australia-SA, Australia-TAS, Australia-VIC, Australia-WA, Malaysia Sabah, Malaysia Sarawak, Brunei, Papua New Guinea, Indonesia, Solomon Islands Shortland, Solomon Islands general.

[36] *Europe*. Ireland, Britain, Spain, Portugal, Portugal Islands, Luxemburg, Belgium, Italy, Netherlands, France, Denmark, Poland, Switzerland, Slovenia, Slovakia, Czech Republic, Hungary, Austria, Germany, Estonia, Sweden Gotland, Balkan USGS, North Europe, European part of the GLiM Asia (which is then eliminated there).

[37] *North America*. Conterminous U.S., North Part, South Part.

[38] 1. Conterminous U.S.: All maps of the conterminous U.S. were trusted on the same level, and overlaps were so small that they were not erased of each other, but converted into blank features, which then were automatically merged (ArcGIS tool “Eliminate”) to the neighboring feature with the longest common border.

[39] 2. North Part: Yukon, Alaska, British Columbia, Alberta, Saskatchewan, Ontario, Manitoba, Nova Scotia, Newfoundland, Slave Province, Western Churchill, Baffin Island, Northwest Territories, Canada.

[40] 3. South Part: The south part of the North American map does not contain border conflicts between regional maps, but all parts of the geological map of North America were deleted by hand which overlapped regional maps in this region.

[41] *South America*. Brazil, French Guyana, Venezuela, Guyana, Suriname, Trinidad and Tobago, Ecuador, Chile, Peru, Colombia, Bolivia, Uruguay, Argentina, Paraguay.

[42] Following the erase procedure deleting overlapping polygons, the maps were joined with the translation tables containing the regional lithology translations (see section A3) and merged to a continental map. However, this continental map still contained numerous topological errors, mainly overlaps within individual regional maps or gaps at map borders. These were now solved by hand using the topology tools of ArcGIS. If during topology works polygons were generated whose size fell below the cluster limit of the geodatabase (of 0.1 m) and thus became zero, these polygons were deleted.

[43] The topological errors were distinguished into minor and major errors. Minor errors were small (usually smaller than 10 km², on average they were smaller than 1 km²) and induced no bias. However, if a smaller unit was completely covered by a larger unit, this was classified a major error, as everything else that was not a minor error. Minor topological errors were treated automatically, as there were more than 120,000 of them. Thus manual solving was not reasonable. Each minor gap and each minor overlap was converted into a blank feature, which then was automatically merged (using the ArcGIS tool “Eliminate”) to the neighboring feature with the longest border. The effects of the automatic procedure were visually checked against the original data at some locations, where it induced no

major inconsistencies (as expected for these very small features).

[44] The major errors were treated manually according to their cause. Most of these treatments are listed below:

[45] *Africa*. In Mozambique, two smaller units were overlain by a larger polygon in the original data. This was resolved by erasing their shapes from the overlapping polygon.

[46] *Antarctica*. No major topological problems were encountered in Antarctica.

[47] *Asia*. (1) Four major overlaps in China Neij Mongol, were solved favoring the smaller polygon, which would otherwise have been erased. (2) Five small islands in the Pacific were deleted that had no defined lithology and did not border to any known lithology. (3) The geological map of the former Soviet Union had been digitized by hand by student assistants. Many small topology errors were introduced by this procedure, which were treated separately before the Soviet Union map was included in the Asian map. Some major overlaps also occurred, which were solved favoring the smaller units to avoid their deletion. The reference scale of the map was reduced from 1:2.5 million originally to 1:3 million because of possible errors in digitalization.

[48] *Australia and Oceania*. (1) Several map-internal overlaps, particularly in Indonesia, between nearly identical rocks were solved manually. (2) In Indonesia, some gaps were relabeled as water bodies after visual inspection of satellite imagery.

[49] *Europe*. Particularly in Belgium, many topographical errors existed. Major errors were treated by hand. However, about 90,000 small overlaps and gaps were treated automatically. This may have caused some flaws in the representation of the Belgian geology.

[50] *North America*. (1) In the geological map of Canada, some duplicate polygons (i.e., a situation where two exactly the same polygons overlap) were resolved by deleting one of them. (2) In Hawaii, Mexico, Ontario and Saskatchewan, some smaller units were covered with larger ones. Their shapes were cut from the larger to resolve the overlap. (3) In Mexico, two major areas overlapped. One was a duplicate polygon with exactly the same attributes; the other overlap was by two polygons sharing the same geometry but classified as either basalt or alluvial. The alluvial was preferred, because of the polygon shape. (4) In Puerto Rico, a swamp classed polygon overlaps a number of other

smaller units, which were cut from the larger polygon to retain them. Also one water polygon overlapped another polygon, which was favored because satellite imagery did not show water at that position.

[51] *South America*. (1) The 100 largest of the automatically filled gaps were manually checked if they were water bodies, using satellite imagery (MS Bing), and labeled accordingly. Approximately half of the polygons were water bodies. (2) 29 small holes (originally treated as minor errors) had to be merged with a neighboring polygon by hand because they caused errors when merged to their nearest neighbor automatically (actually the automatic procedure erroneously erased the nearest neighbor and themselves in the process likely due to an internal error in ArcGIS). In these cases, a new polygon was created (using the edit tool "Auto Complete Polygon") and merged to the subjectively best suited neighbor.

[52] Finally, border conflicts between the continental maps were solved based on the time of finalization. Continental maps finalized earlier were favored over those finished later. The order was: South America, North America, Africa, Asia, and Europe. Australia and Oceania had no conflicts which needed to be resolved.

A3. Lithological Translation Methods

[53] The translation from stratigraphic units provided by most of the geological maps to lithological units required two steps: First, the rock types associated with the stratigraphic unit were identified; second, this detailed information was translated into the general classes applied here.

[54] Ideally, the rock type information was already provided by the geological map data sets as an additional attribute. In this case that information was used and translated. The depth of the detail of rock description varies largely between maps, however. For example, the U.S. State maps provide major and minor rock types (maximum two rock types and standardized names), while the Australian regional maps provide a detailed description of all rocks occurring within the respective unit, often more than 250 characters long. The geological map of China and the former Soviet Union provided rock type information only in their native languages. These were translated to English by graduate assistants fluent in these languages.

[55] If the geological maps did not provide any information on rock types, descriptions of the

Table A2. List of the Lithological Classes and Subclasses in the Three Levels

Code	Description
<i>Level 1: xx</i>	
su	Unconsolidated sediments
ss	Siliciclastic sedimentary rocks
py	Pyroclastics
sm	Mixed sedimentary rocks
sc	Carbonate sedimentary rocks
ev	Evaporites
va	Acid volcanic rocks
vi	Intermediate volcanic rocks
vb	Basic volcanic rocks
pa	Acid plutonic rocks
pi	Intermediate plutonic rocks
pb	Basic plutonic rocks
mt	Metamorphics
wb	Water Bodies
ig	Ice and Glaciers
nd	No Data
<i>Second Level: yy</i>	
ad	Alluvial deposits
ds	Dune sands
lo	Loess
la	Laterites
or	Organic sediment
mx	Mixed grain size
sh	Fine grained
ss	Coarse grained
am	Mafic metamorphics mentioned
gr	Greenstone mentioned
pu	(Pure) carbonate
py	Pyroclastics mentioned
<i>Third Level: zz</i>	
bs	Black shale mentioned
cl	Fossil plant organic material mentioned
ch	Chert mentioned
fe	Iron minerals mentioned
ph	Phosphorous-rich minerals mentioned
pt	Pyrite mentioned
gl	Glacial influence mentioned
mt	Metamorphic influence mentioned
ev	Subordinate evaporites mentioned
vr	Subordinate volcanics mentioned
pr	Subordinate plutonics mentioned
sr	Subordinate sedimentary rocks mentioned
su	Subordinate unconsolidated sediments mentioned
we	Intensive weathering mentioned

named stratigraphic units were sought in regional literature. The quality of the literature was variable and may have introduced some uncertainty. In some rare locations, the rock type information of digital geological map vector data sets was derived from paper maps, which were georeferenced and visually assigned to the units of the digital maps.

[56] The different rock type information was translated into the lithological classes listed below, usually emphasizing more reactive rock types.

However, determining the dominant rock types within a unit was not always straightforward. In accordance with abundant mapping guidelines it was assumed that rocks mentioned foremost are more abundant than those mentioned later in rock unit descriptions. In some cases combinations of certain rock types led to a specific classification. Technically, all translations were done in a join-table, which combined the attribute of the geological map that the join was based upon (usually defining the stratigraphic unit), with the rock description, lithology class, age, source and quality information as well as additional comments. In very few cases, two stratigraphic units were given different (but similar) rock compositions in the source map. In this case, only one of the different compositions was used for translation. Affected Regions where this occurred include British Columbia, Manitoba, Nova Scotia and Yukon in Canada. The same process was applied where the age attribute differed, where e.g., three units in British Columbia had two different ages assigned, of which only the first was used.

A4. The Lithological Classification

[57] The lithological classification is an extension of the classification system by Dürr *et al.* [2005], with additional classes and two new levels of information [cf. Jansen *et al.*, 2010; Moosdorf *et al.*, 2010]. The lithological classification is now represented by a six-symbol code: “xxyyzz,” where “xx” represents the code of the first level information, comparable to the existing lithological nomenclature. In addition, “yy” gives more detailed information on rock types, while “zz” provides special information about the mapped unit. The second and third level information is optional, and not all units provide additional information (represented by the code “_”). This structure reduces information to few general classes but has additional detail information when needed and available. Table A2 lists all lithological classes and subclasses.

A4.1. First Level: xx

[58] The first level lithological class was mandatory for all units. It describes the general rock types in the mapped unit.

A4.1.1. Siliciclastic Sediments (su, ss, py)

[59] *Unconsolidated sediment (su)* represents young, not yet consolidated sediments, usually of Cenozoic age. It comprises all grain sizes, which are indicated by the second level information.

Additionally, more specific unconsolidated sediments can be indicated there, i.e., dune sands, alluvial deposits, loess and swamps. Examples of unconsolidated sediments are sands, mud, swamp deposits, dunes, and beach sands. Where carbonate is reported in the unconsolidated sediment, the lithological class sm (see below for description) is usually selected, thus losing the information on the consolidation, but maintaining the information on the carbonate content.

[60] *Siliciclastic sedimentary rocks (ss)* represent e.g., sandstone, mudstone and greywacke, but may also show a small degree of metamorphic alteration (e.g., shale; this would be indicated by the third level information mt). Siliciclastic sediment is usually accompanied by the second level information of grain size. Where carbonate was named in the rock description of the mapped unit, the lithological unit sm was used, so siliciclastic sedimentary rocks are without mapped carbonate influence. Note that in some cases the carbonate presence (e.g., as matrix) may not be named in the rock description, and siliciclastic sediments may still contain carbonate in reality.

[61] *Pyroclastics (py)* are sediments of volcanic origin. Typical pyroclastics are tuff, volcanic breccias, or ash. If a unit appears to be dominated by pyroclastics, its first level information was set to py. However, if only smaller amounts of pyroclastics are described in the rock description, instead the second level information can be set to py.

A4.1.2. Carbonate-Rich Sedimentary Rocks and Evaporites (sc, sm, ev)

[62] *Mixed sedimentary rocks (sm)* represent all sediments where carbonate is mentioned but not dominant, plus some units that were identified as sediments, but no information on the type of sediment was available. Mixed sedimentary rocks are usually a combination of different rock types (e.g., interlayered sandstone and limestone). Another classical representative rock of that class would be e.g., shaley marl. The class sm is usually accompanied by the second level information of the grain size of its siliciclastic fraction. Sediments classed as sm may in some cases also be unconsolidated, in which case the information on the carbonate content was retained rather than the information on the consolidation state.

[63] *Carbonate sedimentary rocks (sc)* are dominated by carbonate rocks. Examples of sc units are limestone, dolomite and marl. As usually the rock

descriptions of the mapped units do not give relative abundances of the rock types which they encompass, units were classed as sc if the first named rock type was a carbonate rock, if the majority of rock types were carbonates or if the named order otherwise led to the impression of a domination by carbonates. If siliciclastic sediments were mentioned in a carbonate-dominated unit, the grain size of the siliciclastic fraction is usually given in the second information level. If that was not the case, the second level code “pu” indicates the unit as a pure carbonate.

[64] *Evaporite (ev)* units contain substantial amounts of evaporitic rocks. The typical encountered evaporite rock was gypsum, but also anhydrite, halite or nomenclatures as “salt pan.” If a map unit was interpreted as dominated by evaporites, it was classed as ev, regardless of other mentioned rocks. This implies, that ev units may additionally contain, e.g., carbonates. This would usually be indicated by the available subclasses.

A4.1.3. Volcanics (va, vi, vb)

[65] Volcanic rocks are divided in three main lithological classes, based on their composition.

[66] *Acid volcanics (va)* are typically rhyolites, trachytes or dacites and classed after *Le Maitre* [2004].

[67] *Intermediate volcanics (vi)* are classically andesites. However, if basic or acid volcanics are mentioned in addition to intermediate volcanics, then the former classes are selected. Units featuring only acid and basic volcanics or, rarely, volcanics of unknown type are also classed as intermediate volcanics.

[68] *Basic volcanics (vb)* are usually basalt-type rocks. Rock types classified as basic volcanics (following *Le Maitre* [2004]) apart from basalts are e.g., tephrites, tholeites, and lamprophyres.

A4.1.4. Plutonics (pa, pi, pb)

[69] *Acid plutonics (pa)* represent plutonic rocks containing quartz. Granites and their relatives are grouped in this class in particular, but also quartz-diorites and quartz-monzonites. In addition, there is an overlap with the metamorphic rocks, as some migmatites may also be referred to as granite in the geological maps, so would be classed as pa, as would granitic gneiss, or slightly folded granite. Although these are strictly metamorphic rocks in that they experienced metamorphosis, but their

attributes are likely to be more related to the acid plutonics than to some of the members of the very heterogeneous group of metamorphic rocks.

[70] *Intermediate plutonics (pi)* encompass a group of non-mafic plutonic rock types mainly defined by their ‘relative’ absence of quartz. This class is dominated by diorite, monzonite, syenite and their subtypes. In addition, this class was used in the few cases in which no clear character of a mapped plutonic rock could be identified.

[71] *Basic plutonics (pb)* include plutonic rocks rich in mafic minerals, like gabbro and peridotite, as well as ultrabasic species like norite. Ophiolite structures would be classified as basic plutonic. Basic plutonic rocks can, like their acid and intermediate counterparts, show a certain degree of metamorphism but still be classified as basic plutonic.

A4.1.5. Metamorphics (mt)

[72] *Metamorphics (mt)* is the ‘broadest’ lithological class. It encompasses a wide variety of rocks from shales to gneiss, from amphibolite to quartzite. If the metamorphics contain marble, they were assigned the yy attribute ‘pu’, indicating the presence of carbonate outside sc or sm units. If they contained mafic metamorphic rocks (e.g., amphibolite), the yy-attribute was set to ‘am’. If the metamorphism seemed only weak, the original rock was used to define the lithological class. For example the rock description “foliated, slightly folded granite” triggered classification as acid plutonic rock (xx = pa), accompanied by a zz-value of ‘mt’, indicating a metamorphosis of the rock.

A4.1.6. Other Units (wb, ig, nd)

[73] *Water bodies (wb)* were included although not a rock type. Some geological maps either include water bodies as a geological unit or leave the water covered areas blank, leading to gaps in the digital data set. Such areas are classed as water bodies here. They encompass lakes, rivers, but also parts of some coastal oceans. The water body areas are not meant to be used as indication of water areas in general, as they are not a priority of this data set. Here, lithology is prioritized, but if the geological map had water as ‘rock type’, ‘water bodies’ was assigned.

[74] *Ice and glaciers (ig)* are mainly situated in Antarctica, Greenland and on some mountains. However, the coverage is not representative for an ice extent, as the priority of this map is on lithology. If the source geological maps identified ice, then class ig was used. The only exception is Antarctica, where the

geological map did not represent ice areas and an independent data set was used for their representation.

[75] *Non-defined (nd)* units are blank areas of the map. Although much effort was put into representing all areas of the globe, the rock types of some units could not be identified and are classed as non-defined. Most are classed as undefined in the source geological maps. A significant proportion of these units seems to be situated under glaciers.

A4.2. Second Level: yy

[76] The second level lithological information was introduced to further describe the mapped rocks where possible and appropriate. Information of this level can be combined with the first level to adapt the classification to answer specific questions. However, contrary to the first level information, the second and third level information is not mandatory and has not been assigned in all cases.

A4.2.1. Specific Unconsolidated Sediments (ad, ds, la, lo, or)

[77] Some specific unconsolidated sedimentary units were dealt with separately because of their specific attributes, composition or genesis.

[78] *Alluvial deposits (ad)* are very young sediments associated with fluvial systems. They are mapped if the original map referred to alluvial units, e.g., alluvial fans or terraces. However, their representation in the different maps is very variable, because many geological maps do not differentiate unconsolidated units.

[79] *Dune sands (ds)* are aeolian sediments, occurring mostly at beaches or in deserts. Again, their representation is incomplete, as many geological maps did not contain this information.

[80] *Laterites (la)* were included if deeply weathered soils are described in the source literature. As the reference depth for the GLiM cannot be consistent, the subclass la indicates a thick weathered soil horizon above the rocks defining the lithological class.

[81] *Loess (lo)* is an aeolian sediment deposited during glacial times. However, the representation of loess is complicated, as it is often only a thin cover, which is ignored by the geological maps. No additional sources were used for this attribute, thus its coverage, compared to specialized maps focusing loess cover [e.g., Bettis *et al.*, 2003; Haase *et al.*, 2007; Pécsi, 1990; Zárate, 2003], is very restricted.

[82] *Organic sediment (or)* is mapped if the geological map mentions swamp deposits. Organic sediment has a very specific environment, which can be important for numerous applications. However, again, not all swamps were identified in the map, as no additional sources apart from the geological maps were used.

A4.2.2. Sediment Grain Sizes (mx, sh, ss)

[83] If it was possible to define a dominating grain size in siliciclastic sediments, this was indicated by a second level attribute. These attributes were added to consolidated and unconsolidated sedimentary lithology classes.

[84] *Mixed grain sizes (mx)* indicate the lack of a dominating grain size, or the absence of grain size information.

[85] *Fine grains (sh)* indicate a dominance of grains finer than sand.

[86] *Coarse grains (ss)* indicate a dominating grain size of at least sand diameter.

A4.2.3. Metamorphic Rocks (am, gr)

[87] The lithological class of metamorphic rocks encompasses a multitude of different rock types. The reference to some specific metamorphic rocks resulted in the assignation of a second level attribute, even in cases of non-metamorphic lithological classes (e.g., plutonic dominated units).

[88] *Mafic metamorphic rocks (am)* are classed if rocks like amphibolite or serpentinite are indicated in the geological units. Amphibolites, for instance, have weathering rates that are about four times higher than gneiss [Meybeck, 1987] and are thus highlighted by this attribute.

[89] *Greenstones (gr)* are mapped to highlight their presence. The greenstone belts are old and complex structures and may thus be of interest for a number of applications. However, not all greenstone belts are represented in the lithological map, only those indicated by their source maps. Thus gr should not be used as a comprehensive reference of the occurrence of global greenstone.

A4.2.4. Carbonate and Pyroclastics Occurrence (pu, py)

[90] *Pure carbonate rocks (pu)* was intended to indicate sc, pure carbonate units without siliciclastic contents. However, the pu-attribute is also used in non-sedimentary units to indicate minor

carbonate occurrences that are not sufficient enough to justify setting the first level to sc.

[91] *Pyroclastic sediments (py)* can also be attributed in the second level. In contrast to the first level py, the second level attribute indicates some occurrences of pyroclastics rather than the dominant rock type of the mapped unit. This subclass is a recognition of the potential importance of the presence of pyroclastics, which was reported to exhibit very high weathering rates when fresh [Dahlgren et al., 1999; Hartmann et al., 2010].

A4.3. Third Level: zz

[92] The third level adds information on individual aspects of the rocks and other mapped rock types, if available. Again, the third level information is neither mandatory nor exhaustive. The absence of third level information does not necessarily indicate the absence of the associated rock attribute. However, the presence of a third level zz-code clearly indicates the presence of the respective rock attribute.

A4.3.1. Fossil Carbon Rich Sediments (bs, cl)

[93] The occurrence of fossil carbon rich sediments can be relevant for some scientific questions, like the quantification of processes in the carbon cycle [Copard et al., 2007].

[94] *Black shale (bs)* is mapped where, e.g., black shale, oil shale, organic-rich shale, or similar rock units are named in the rock description.

[95] *Coal (cl)* is mapped if significant occurrences of any kinds of lignite or coal, up to anthracite, are noted in the rock description. It is often associated with siliciclastic or mixed sediment.

A4.3.2. Special Occurrences (ch, fe, ph, pt)

[96] Occurrences of certain materials in rock units, which may be interesting for individual studies are indicated by this third level attribute group. However, the latter three units are very rare and do not represent all occurrences of the materials.

[97] *Chert (ch)* occurrences are mapped individually. Example uses for chert include estimating silica exports from carbonate rock units as reported by Jansen et al. [2010].

[98] *Iron (fe)* is mentioned if the geological maps noted iron deposits, e.g., banded iron formations or magnetite.

[99] *Phosphorous (ph)* was noted if, e.g., phosphorite or apatite were indicated in the geological

maps or the description of the rocks. This information was added, because phosphorous is an important nutrient and thus the supply of that element to the biosphere via chemical rock weathering may be essential in many areas.

[100] *Pyrite (pt)* is occasionally noted in the geological maps or the literature describing the rocks. Pyrite weathering can be important for matter flux mass balances and impacts the pH of adjacent waters bodies.

A4.3.3. Genesis Aspects (gl, mt, we)

[101] Rock properties are of course strongly affected by their genesis which is only seldom represented in the geological maps. However, three attributes containing particular information about the rock genesis are indicated at the zz level.

[102] *Glacially overprinted units (gl)* indicate references to previous glacial activity. This encompasses directly glacial sediment, e.g., glacial till, as well as glacial overprinting in the unit description of other rock units. Again, this is in no way a complete description of all glacially affected units globally.

[103] *Metamorphic rocks (mt)* are highlighted in the third level in addition to the first level as well as the second level focusing, which focused on two certain rock types solely (yy = 'am' or 'gr'). Third level 'mt' occurs in two possible settings. If rocks are slightly metamorphosed, but their source rock was clearly named in the geologic description, e.g., foliated granite, the original rock type defined the lithological class (in this case xx = pa), and the third level information mt was added. If the properties of the slightly metamorphosed rocks are still similar to their source rocks, this procedure defines these rocks better than setting xx = mt, which would add them to the very broad group of metamorphics. For example, in order to emphasize the carbonate content, a marble would be classed scpmt (a metamorphosed pure carbonate rock). The second meaning of the third level information mt is that within the mapped units, minor occurrences of metamorphic rocks are named in a unit dominated by a rock type leading to a different first level attribute than mt.

[104] *Intensive weathering (we)* is not directly a genetic attribute of the original rock, but has altered the attributes of the rock which is now encountered in the unit where it is mapped. The subclass "we" was attributed where the source geological map indicates rocks as weathered.

A4.3.4. Minor Occurring Rock Types (ev, vr, pr, sr, su)

[105] If rock types other than the dominating rock that defines the first level lithological class are noted within a map unit, they can be added to the third level aggregated in broad rock groups.

[106] *Evaporites (ev)* define only small occurrences of evaporitic sediments. Evaporites weather very fast, and thus have a big impact on the chemistry of natural waters and are always mentioned in the lithological classification if they are mentioned in the geological maps (either as first or third level lithological information).

[107] *Volcanic rocks (vr)* highlight the mapped occurrence of minor acid, intermediate or basic volcanic rocks in a map unit of a different lithological class.

[108] *Plutonic rocks (pr)* are classed at the third level if minor occurrences of acid, intermediate or basic plutonics are noted in the geological information.

[109] *Sedimentary rocks (sr)* highlight minor sedimentary strata in the described unit. If these minor units contain mapped carbonates, the third level attribute sr is accompanied by the second level attribute pu. For example, a unit of basalt lava flows interbedded with minor limestone would be classed as vbpusr.

[110] *Unconsolidated sediments (su)* highlights a dominating unit covered with unconsolidated sediments. Unfortunately, geological maps do not follow a common rule concerning the thickness of cover necessary to map unconsolidated sediments as individual units in their own right or to note them at all. Thus, the absence of the attribute does not indicate that the mapped bedrock unit is outcropping right below the topsoil horizon, but its presence indicates that certainly a significant cover of unconsolidated sediment is situated on top of the mapped lithological class. This could be important e.g., for hydrogeological applications.

A4.4. Conflict Management

[111] The applied lithological classification system allows only one single description per level of information. This of course causes conflicts if the rock description of any mapped unit allows attribution of multiple classes. The conflicts were solved in general by selecting the class with the greatest impact on matter fluxes by chemical weathering (which was the original purpose of the map). This




Figure A2. Spatial map quality (map resolution) and point lithology information for comparison.

leads to a general order of classification as follows (first named classes are favored above those following later): First level: ev, py, sc, vb, sm, pb, vi, va, pi, pa, su, ss, mt, wb, ig, nd; Second level: py, pu, la, am, gr, or, lo, ad, ds, sh, ss, mx; Third level: ev, ph, bs, cl, gl, fe, pt, vr, we, mt, ch, pr, su, sr.

[112] In specific cases the previously named order was changed if the rock description indicated a dominance of a minor attribute. For instance a unit mapped as swamp with very small occurrences of pyroclastics would be classed as “suor_,” although the general pattern would demand to name it “supy_.”

A5. Quality Classification System and Evaluation

[113] The presented global lithological map is a complex construct based on a large number of sources. To give some estimate of the reliability of the contained information at each position, a quality classification system was introduced.

[114] The spatial quality attribute ‘sqa’ is based on the scale of the source geological map as a metric attribute. To simplify the reading, it is defined as ‘sqa’ = ‘source scale/10⁶’ (Figure A2, source scales are provided in Table A1).

[115] The applied ‘thematic quality attribute’ is more difficult to interpret. It contains a rating of A, B, or C to provide a broad sense of the reliability of the lithological classification. While in theory, every lithological class should correctly describe the real rock types at its location, this is more likely where the geological map directly provided the information for the lithological classification. These areas get a thematic quality attribute of “A,” representing the highest reliability. If additional regional literature had to be consulted to identify the rock types associated with the stratigraphic units named in the geological source maps, the attribute “B” is given. This also applies to areas where the classification based on the geological maps was uncertain. The weakest quality, indicated by the “C” value of the attribute, indicates that only very general information about the rock types of this unit was available.

[116] However, the quality attributes were not always individually assigned to each polygon. In particular the thematic quality is only a guide to the general reliability of the information of the areas, and should be treated with care.

[117] To evaluate the accuracy of the GLiM at the point-scale, a set of 290 000 data points with rock type information was extracted from three large

databases (PetDB [Lehnert *et al.*, 2000]: www.petdb.org, SedDB: www.seddb.org, and VentDB: www.ventdb.org via the EarthChemPortal: www.earthchem.org; USGS National Geochemical Database NGDB [U.S. Geological Survey, 2008] and OZCHEM [Champion *et al.*, 2008]). The reported rock names were translated into lithological classes where possible and where the rock did not indicate a very local phenomena and exotic rock types (e.g., ore bodies). However, only the USGS NGDB data are outcrop data with certainty, while the others also include drill-core interpretations or surface samples. To exclude as many of these cases as possible, unconsolidated sediments were omitted from the evaluation, as well as points where the lithological map reports water, ice/glaciers or an unknown rock type. The EarthChem portal included spatial accuracy information in their data, which was restricted to data sets with an accuracy $<0.01^\circ$. After the deselection processes, 164,953 points remained, of which in 40% of the cases, the lithology of the GLiM and the lithology classification of the independent sample match. This is a reasonable result when inaccuracies in the sample spatial positions are taken into account (e.g., inaccuracies of up to 10 km may occur in the OZCHEM database). If similar lithological classes are also taken into account (e.g., for mapped mixed sediments in the GLiM, the siliciclastic and carbonate lithologies of points would be accepted as “similar,” or for basic or acid igneous classes also their intermediate neighbors), the match increases to 64% of the points with similar lithologies to those mapped in the GLiM (Figure A2).

A6. Contributors to GLiM

[118] Thank you to all contributors; and if anyone was forgotten in the multitude of people and organizations contributing to the map – please excuse us.

[119] The map needed extensive manual digitalization, which was done by Birte Meier (Argentina, Soviet Union, and early version Africa), Svetlana Didorenko (former USSR), Andre Paul (Austria, Colombia, Ecuador, Paraguay, Soviet Union, and Uruguay), Mercedes Pordzik (Himalaya). The rock descriptions of the geological maps of the former USSR and China had to be translated into English, which was done by Svetlana Didorenko and Yixi Gu. Some regions were translated into lithological classes as part of graduate theses by Nadja Hoppe (Eastern Australia: Queensland, New South Wales, Victoria, Tasmania), Birte Meier (early versions of

Africa, South Africa), Svetlana Didorenko (Siberia) Henning Kedenburg (India) and Mercedes Pordzik (Himalaya).

[120] Several people from all over the world helped with map acquisition, which were Thomas Schlüter, Elisabeth Sillmann (Africa); James Passmore and Oliver Raymond (Antarctica); Loretta Crisby-Whittle (Newfoundland, Canada); Andrea Salas, Andreas Weiss, Caroline Raulf (Chile); Martin Schreck (Costa Rica); Jean- Paul Cadet, Philippe Rossi, CGMW (diverse maps); Rainer Lezius (India); Ana Patricia Mora Aguilar, Carlos Alcocer Valdés, Margarita Aldana (Mexico); A. Saidi (Middle East); Pavel Hanzl (Mongolia); Mark Rattenbury (New Zealand); Heiko Dirks (Arab Peninsula); Carlos Schobenhau (South America); Arthur W. Snoke (Tobago), and Paul Hackley (Venezuela).

[121] The project ran on very limited funding and relied on mostly free access to source data. Many agencies and companies provided maps for free: Alberta Geological Survey (Alberta, Canada); BRGM (Nouvelle Calédonie, Guyane); British Columbia Ministry of Energy and Mines (British Columbia, Canada); Geological Survey of Britain (United Kingdom); CGMW (Middle East, South America, World); Serviço Geológico do Brasil (Brazil); China Geological Survey (China); FAO (Africa); FAO Africover Project (Somalia); Geological Survey Institute of Indonesia (Indonesia); Geological Survey New Zealand (New Zealand); Geological Survey of Canada (Canada and Western Churchill, Canada); Geoscience Australia (Australia state maps); Geologische Bundesanstalt (Austria); INGEOMINAS, Colombia (Colombia); Instituto Geológico y Minero de España (Spain); Manitoba Geological Survey (Manitoba, Canada); Natural Resources Canada (northern Baffin Island, Canada); Newfoundland and Labrador Department of Natural Resources (Newfoundland, Canada); Northwest Territories Geoscience Office (Northwest Territories, Slave Craton, Canada); Nova Scotia Department of Natural Resources, Mineral Resources Branch (Nova Scotia, Canada); One Geology (providing links to data sources); Data from OneGeology-Europe were used and/or reproduced with the permission of the geological survey organization right holders who comprised the OneGeology-Europe project (Geological Survey of Belgium, Česká Geologická Služba, De Nationale Geologische Onderzoeksinstituten voor Nederland en België, Eesti Geoloogiakeskus OÜ, Istituto Superiore per la Protezione e la Ricerca Ambientale, Geological Survey of the Netherlands, Service géologique du Luxembourg, Laboratório Nacional de Energia e

Geologia, Štátny geologický ústav Dionýza Štúra, Geološki zavod Slovenije, Państwowy Instytut Geologiczny, Magyar Állami Földtani Intézet;). All Rights Reserved); Ontario Geological Survey (Ontario, Canada); Orr and associates (www.orr-bodies.com; China Nei Mongol, China Xinjiang, Madagascar, Malaysia Sabah, Malaysia Sarawak, Mozambique, Solomon Islands, Turkey, Vanuatu); Raafat S. Wassef (Egypt); Saskatchewan Industry and Resources (Saskatchewan, Canada); Serviço Geológico do Brasil (Brazil); Servicio Geológico Mexicano (Mexico); U.S. Geological Survey (Puerto Rico, Afghanistan, Venezuela, Arabian Peninsula, Iran, Virgin Islands, Honduras, Africa); Viceministerio de Minas y Energía, Paraguay (Paraguay); Yukon Region, Indian and Northern Affairs Canada (Yukon, Canada). In addition, the Council for Geoscience South Africa (South Africa) and the Geological Survey of Namibia (Namibia) provided their respective maps for a reduced price. The providers of the EarthChem Portal (www.earthchem.org), the National Geochemical Rock Database (USGS), and OZCHEM (Geoscience Australia),

[122] Hans Dürr, Ronny Lauerwald, Tom Gleeson, and Clement Bataille improved the map with general consulting. Mark Mulligan, Stephen Pordor, and Clement Bataille helped to improve the readability of the manuscript.

Acknowledgments

[123] The GLiM is a product of a series of projects, partly funded by the German Science foundation (grants HA4472/6-1 and Cluster of Excellence CLISAP, EXC107) and supported by the TU Darmstadt via the working group of Stephan Kempe. A large number of people and institutions contributed to the map, they are acknowledged here and are named in Appendix A. In particular, Hans Dürr supported the project with his knowledge about lithological classification. Two anonymous reviewers are acknowledged for their constructive criticism of the manuscript.

References

- Abed, A. M., and B. S. Amireh (1999), Sedimentology, geochemistry, economic potential and palaeogeography of an Upper Cretaceous phosphorite belt in the southeastern desert of Jordan, *Cretaceous Res.*, **20**(2), 119–133, doi:10.1006/cres.1999.0147.
- Africover (2002), Geology of Somalia at scale of 1:1,000,000, Land and Water Dev. Div., Food and Agric. Organ., Rome.
- Ahmad, T., and M. I. Bhat (1987), Geochemistry and petrogenesis of the mandidarla volcanics, northwestern Himalayas, *Precambrian Res.*, **37**(3), 231–256, doi:10.1016/0301-9268(87)90069-6.
- Ahmad, T., and J. Tarney (1991), Geochemistry and petrogenesis of Garhwal volcanics: Implications for evolution of the north Indian lithosphere, *Precambrian Res.*, **50**(1–2), 69–88, doi:10.1016/0301-9268(91)90048-F.
- Alam, M., M. M. Alam, J. R. Curray, A. L. R. Chowdhury, and M. R. Gani (2003), An overview of the sedimentary geology of the Bengal Basin in relation to the regional tectonic framework and basin-fill history, *Sediment. Geol.*, **155**(3–4), 179–208, doi:10.1016/S0037-0738(02)00180-X.
- Alavi, M. (1994), Tectonics of the Zagros orogenic belt of Iran: New data and interpretations, *Tectonophysics*, **229**(3–4), 211–238, doi:10.1016/0040-1951(94)90030-2.
- Alberti, A., F. Castorina, P. Censi, P. Comin-Chiaramonti, and C. B. Gomes (1999), Geochemical characteristics of Cretaceous carbonatites from Angola, *J. Afr. Earth Sci.*, **29**(4), 735–759, doi:10.1016/S0899-5362(99)00127-X.
- Al-Hafidh, N. M., and S. A. Qasim (1992), Petrochemistry and geotectonic setting of the Shalair granite, NE Iraq, *J. Afr. Earth Sci.*, **14**(3), 429–441, doi:10.1016/0899-5362(92)90046-F.
- Allen, R., et al. (2008), New constraints on the sedimentation and uplift history of the Andaman-Nicobar accretionary prism, South Andaman Island, *Spec. Pap. Geol. Soc. Am.*, **436**, 223–255.
- Alminas, H. V., E. E. Foord, and R. E. Tucker (1994), Geochemistry, mineralogy, and geochronology of the U.S. Virgin Islands, *U.S. Geol. Surv. Bull.*, **2057**, 36 pp.
- Alsharhan, A. S., and A. E. M. Nairn (1994), The Late Permian carbonates (Khuff formation) in the western Arabian Gulf: Its hydrocarbon parameters and paleogeographic aspects, *Carbonate Evaporite*, **9**(2), 132–142, doi:10.1007/BF03175226.
- Alsharhan, A. S., and A. E. M. Nairn (1995), Tertiary of the Arabian Gulf: Sedimentology and hydrocarbon potential, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, **114**(2–4), 369–384, doi:10.1016/0031-0182(94)00089-Q.
- Alsharhan, A. S., and A. E. M. Nairn (1997), *Sedimentary Basins and Petroleum Geology of the Middle East*, Elsevier, Amsterdam.
- Alsharhan, A. S., A. E. M. Nairn, and A. A. Mohammed (1993), Late Palaeozoic glacial sediments of the southern Arabian Peninsula: Their lithofacies and hydrocarbon potential, *Mar. Pet. Geol.*, **10**(1), 71–78, doi:10.1016/0264-8172(93)90101-W.
- Amiotte Suchet, P., J. L. Probst, and W. Ludwig (2003), Worldwide distribution of continental rock lithology: Implications for the atmospheric/soil CO₂ uptake by continental weathering and alkalinity river transport to the oceans, *Global Biogeochem. Cycles*, **17**(2), 1038, doi:10.1029/2002GB001891.
- Anderson, J. L. (1944), Clastic dikes of the Chira and Verdun formations northwestern Peru, *J. Geol.*, **52**(4), 250–263, doi:10.1086/625215.
- Anton, D. J. (1993), *Thirsty Cities: Urban Environments and Water Supply in Latin America*, 197 pp., Int. Dev. Res. Cent., Ottawa, Ont., Canada.
- Archer, J. B. (1977), Llanvirn stratigraphy of the Galway-Mayo border area, western Ireland, *Geol. J.*, **12**(1), 77–98, doi:10.1002/gj.3350120106.
- Arengi, J. T., and G. V. Hodgson (2000), Overview of the geology and mineral industry of Nicaragua, *Int. Geol. Rev.*, **42**(1), 45–63, doi:10.1080/00206810009465069.
- Aspler, L. B., and J. R. Chiarenzelli (1997), Initiation of similar to 2.45–2.1 Ga intracratonic basin sedimentation of the Hurwitz Group, Keewatin Hinterland, Northwest Territories,

- Canada, *Precambrian Res.*, 81(3–4), 265–297, doi:10.1016/S0301-9268(96)00038-1.
- As-Saruri, M. A., R. Sorkhabi, and R. Baraba (2010), Sedimentary basins of Yemen: Their tectonic development and lithostratigraphic cover, *Arabian J. Geosci.*, 3(4), 515–527, doi:10.1007/s12517-010-0189-z.
- Auchapt, A., C. Dupuy, J. Dostal, and M. Kanika (1987), Geochemistry and petrogenesis of rift-related volcanic rocks from South Kivu (Zaire), *J. Volcanol. Geotherm. Res.*, 31(1–2), 33–46, doi:10.1016/0377-0273(87)90004-7.
- Bahlburg, H., V. Carlotto, and J. Cardenas (2006), Evidence of Early to Middle Ordovician arc volcanism in the Cordillera Oriental and Altiplano of southern Peru, Ollantaytambo Formation and Umachiri beds, *J. South Am. Earth Sci.*, 22(1–2), 52–65, doi:10.1016/j.jsames.2006.09.001.
- Balaguru, A., and G. Nichols (2004), Tertiary stratigraphy and basin evolution, southern Sabah (Malaysian Borneo), *J. Asian Earth Sci.*, 23(4), 537–554, doi:10.1016/j.jseas.2003.08.001.
- Balasubrahmanyam, M. N. (2006), *Geology and Tectonics of India: An Overview*, 204 pp., Int. Assoc. for Gondwana Res., Kochi, Japan.
- Bataille, C. P., and G. J. Bowen (2012), Mapping ⁸⁷Sr/⁸⁶Sr variations in bedrock and water for large scale provenance studies, *Chem. Geol.*, 304–305(0), 39–52, doi:10.1016/j.chemgeo.2012.01.028.
- Bawiec, W. J. (Ed.) (1999), *Geology, geochemistry, geophysics, mineral occurrences and mineral resource assessment for the Commonwealth of Puerto Rico* [online], *U.S. Geol. Surv. Open File Rep.*, 98-038.
- Bayasgalan, A., R. Carson, B. Jordon, and K. Wegmann (2007), *Geology of the Hangay Nuruu, Central Mongolia*, paper presented at Twentieth Annual Keck Research Symposium in Geology, Keck Geology Consortium, Wooster, OH.
- Beikman, H. M. (1980), *Geologic map of Alaska*, U.S. Geol. Surv., Reston, Va.
- Bendezú, R., and L. Fontboté (2002), Late timing for high sulfidation cordilleran base metal lode and replacement deposits in porphyry-related districts: The case of Colquijirca, central Peru, *SGA News*, 13(1), 1–9.
- Berner, R. A. (1994), Geocarb-II: A revised model of atmospheric CO₂ over Phanerozoic time, *Am. J. Sci.*, 294(1), 56–91, doi:10.2475/ajs.294.1.56.
- Berrocal, J., and C. Fernandes (1996), Seismicity in Paraguay and neighbouring regions, in *Alkaline Magmatism in Central-Eastern Paraguay: Relationships With Coeval Magmatism in Brazil*, edited by P. Comin-Chiaromonti and C. B. Gomes, pp. 57–66, State of São Paulo Res. Found., Sao Paulo, Brazil.
- Besairie, H. (1964), *Carte Géologique de Madagascar*, Serv. Géol. de Madagascar, Antananarivo.
- Bettis, E. A., D. R. Muhs, H. M. Roberts, and A. G. Wintle (2003), Last glacial loess in the conterminous USA, *Quat. Sci. Rev.*, 22(18–19), 1907–1946, doi:10.1016/S0277-3791(03)00169-0.
- Bhat, M. I., P. Lefort, and T. Ahmad (1994a), Bafliaz volcanics, NW Himalaya: Origin of a bimodal, tholeiite and alkali basalt suite, *Chem. Geol.*, 114(3–4), 217–234, doi:10.1016/0009-2541(94)90054-X.
- Bhat, M. I., P. Le Fort, and T. Ahmad (1994b), Bafliaz volcanics, NW Himalaya: Origin of a bimodal, tholeiite and alkali basalt suite, *Chem. Geol.*, 114(3–4), 217–234, doi:10.1016/0009-2541(94)90054-X.
- Bhatia, T. R., and S. K. Bhatia (1973), Sedimentology of the slate belt of Ramban-Banihal area, Kashmir Himalaya, *Himalayan Geol.*, 3, 116–134.
- Bhushan, S. K. (1998), Geochemistry of the Archaean and Proterozoic clastic and carbonate rocks of Rajasthan, India, *Gondwana Res.*, 1(2), 257–265, doi:10.1016/S1342-937X(05)70836-4.
- Bhushan, S. K., C. M. Bindal, and R. K. Aggarwal (1991), Geology of Bomdila Group in Arunachal Pradesh, *J. Himalayan Geol.*, 2(2), 207–214.
- Billet, M. F., and A. Cowden (1980), A geological visit to Madeira, *Edinburgh Geol.*, 8, 1–7.
- Blackadar, R. G., W. L. Davison, and H. P. Trettin (1968), *Geology Moffet Inlet-Fitzgerald Bay, District of Franklin*, Geol. Surv. of Can., Ottawa, Ont., doi:10.4095/107497.
- Bluth, G. J. S., and L. R. Kump (1991), Phanerozoic paleogeology, *Am. J. Sci.*, 291(3), 284–308, doi:10.2475/ajs.291.3.284.
- Boccaletti, M., F. Innocenti, P. Manetti, R. Mazzuoli, A. Motamed, G. Pasquare, F. di Brozolo, and E. Sobhani (1976), Neogene and quaternary volcanism of the Bijar Area (western Iran), *Bull. Volcanol.*, 40(2), 121–132, doi:10.1007/BF02599857.
- Boucot, A. J., P. E. Isaacson, and G. Laubacher (1980), Early Devonian, eastern Americas realm faunule from the coast of southern Peru, *J. Paleontol.*, 54(2), 359–365.
- Bow, J., M. Brown, and R. Sayre (2009), *Surficial lithology: Africa Terrestrial Ecological Footprint Mapping Project*, Nat. Conserv., Arlington, Va.
- Bradshaw, M. (1992), BMR-APIRA Phanerozoic history of Australia project, South West Pacific data package, 81 pp., Aust. Geol. Surv. Organ., Canberra.
- Brenchley, P. J., and P. F. Rawson (2007), *The Geology of England and Wales*, 559 pp., Geol. Soc., London.
- British Geological Survey (2007), *Digital Geological Map of Great Britain (DiGMapGB)*, Br. Geol. Surv., Nottingham, U. K.
- Brown, A. P. (1913), Notes on the geology of the island of Antigua, *Proc. Acad. Natl. Sci. Philadelphia*, 65(3), 584–616.
- Brunt, M. A., M. E. Giglioli, J. D. Mather, D. J. W. Piper, and H. G. Richards (1973), Pleistocene rocks of Cayman Islands, *Geol. Mag.*, 110(3), 209–221, doi:10.1017/S0016756800036037.
- Bundesamt für Landestopografie (2005), *Geologische Karte der Schweiz*, 1:500,000, Wabern, Switzerland.
- Bureau de Recherches Géologiques et Minières (BRGM) (2001), *Carte géologique de la Guyane*, Orléans, France.
- Bureau de Recherches Géologiques et Minières (BRGM) (2003), *Carte Géologique de la France à 1/1 000 000 - 6ème édition révisée*, BRGM, Paris.
- Bureau des Mines et de l’Energie (1988), *Carte géologique de la République d’Haïti*, scale 1:250,000, Port-au-Prince.
- Bureau of Geology and Mineral Resources of Xinjiang (1992), *Geological map of Xinjiang Uygur, Autonomous Region, China*, version 2, scale 1:1,500,000, Geol. Publ. House, Beijing.
- Buschkuhle, M. (2003), Sedimentology and stratigraphy of middle and upper devonian carbonates in northern Alberta: A contribution to the carbonate-hosted Pb-Zn (MVT) targeted geoscience initiative, *Rep. EUB/AGS Geo-Note 2002-14*, Alberta Geol. Surv., Edmonton, Alberta, Canada.
- Bush, V. A., L. D. Vinogradov, and A. I. Titov (1994), Tectonic breccia and thrust tectonics of the tertiary deposits in northwestern Peru, *Geotectonics. Engl. Transl.*, 28(2), 159–169.
- Callot, P., T. Sempere, F. Odonne, and E. Robert (2008), Giant submarine collapse of a carbonate platform at the Turonian-Coniacian transition: The Ayabacas Formation, southern

- Peru, *Basin Res.*, 20(3), 333–357, doi:10.1111/j.1365-2117.2008.00358.x.
- Campbell, K. E., C. D. Frailey, and L. Romero-Pittman (2006), The Pan-Amazonian Ucayali Peneplain, late Neogene sedimentation in Amazonia, and the birth of the modern Amazon River system, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 239(1–2), 166–219, doi:10.1016/j.palaeo.2006.01.020.
- Candanedo, C., O. Fabrega, V. de Arjona, J. D. Villa, S. Sanchez, R. Arjona, R. Bolanos, and D. Laguna (1998), *Mapa Hidrogeológico de Panama*, Empresa de Transmission Electr., Gerencia de Hidrometeorol., Panama City.
- Carlotto, V., J. Cardenas, V. Cano, and R. Rodriguez (2009), Paleogeographic boundary in the evolution of the Pucara Basin (Triassic-Liassic) and the Arequipa basin (Lias-Dogger): An inheritance of the block accreted during the Mesoproterozoic, in *International Lateinamerika-Kolloquium*, edited by G. Wörner and S. Möller-McNett, pp. 50–52, Universitätsverlag Göttingen, Göttingen, Germany.
- Carrascal-Miranda, E. R., and I. Suarez-Ruiz (2004), Short description of the Peruvian coal basins, *Int. J. Coal Geol.*, 58(1–2), 107–117, doi:10.1016/j.coal.2003.05.004.
- Castelli, D., and B. Lombardo (1988), The Gophu La and western Lunana granites: Miocene muscovite leucogranites of the Bhutan Himalaya, *Lithos*, 21(3), 211–225, doi:10.1016/0024-4937(88)90010-2.
- Central Ground Water Board (2009), Groundwater brochure of Rudraprayag District, Uttarakhand, Minist. of Water Resour., New Delhi.
- Champion, D. C., A. R. Budd, and L. A. I. Wyborn (2008), *OZCHEM National Whole Rock Geochemistry Database*, Geosci. Aust., Canberra, A. C. T.
- Chen, H. Y., A. H. Clark, T. K. Kyser, T. D. Ullrich, R. Baxter, Y. M. Chen, and T. C. Moody (2010), Evolution of the giant Marcona-Mina Justa iron oxide-copper-gold district, south-central Peru, *Econ. Geol.*, 105(1), 155–185, doi:10.2113/gsecongeo.105.1.155.
- China Geological Survey (2001), 1:2,500,000-scale digital geological map database of China, Beijing.
- Christman, R. A. (1953), Geology of St. Bartholomew, St. Martin, and Anguilla, Lesser Antilles, *Geol. Soc. Am. Bull.*, 64(1), 65–96, doi:10.1130/0016-7606(1953)64[85:GOSBSM]2.0.CO;2.
- Clark, A. H., R. M. Tosdal, E. Farrar, and A. Plazolles (1990), Geomorphologic environment and age of supergene enrichment of the Cuajone, Quellaveco, and Toquepala porphyry copper deposits, southeastern Peru, *Econ. Geol.*, 85(7), 1604–1628, doi:10.2113/gsecongeo.85.7.1604.
- Clark, R. J. (2005), Structural constraints on the exhumation of the Tso Moriri Dome, NW Himalaya, Master's thesis, 52 pp., Mass. Inst. of Technol., Cambridge.
- Cocks, L. R. M., R. A. Fortey, and C. P. Lee (2005), A review of Lower and Middle Palaeozoic biostratigraphy in west peninsular Malaysia and southern Thailand in its context within the Sibumasu Terrane, *J. Asian Earth Sci.*, 24(6), 703–717, doi:10.1016/j.jseas.2004.05.001.
- Collazo Carabello, M. P. (2006), *Investigación hidrogeológica del acuífero guaraní en el área aflorante de los departamentos Rivera y Tacuarembó*, Uruguay, Univ. de Buenos Aires, Buenos Aires.
- Colley, H. (1976), Mineral deposits of Fiji (metallic deposits), accompanied by 2 maps: Metallogenic map of Vanua Levu and metallogenic map of Vitu Levu, scale 1:250,000, 123 pp., Miner. Resour. Div., Suva, Fiji.
- Colley, H. (2009), Fiji, geology, in *Encyclopedia of Islands*, edited by R. G. Gillespie and D. A. Clague, pp. 305–309, Univ. of Calif. Press, Berkeley.
- Colman-Sadd, S. P., and L. V. J. Crisby-Whittle (2005), Partial bedrock geology dataset for the Island of Newfoundland (NTS 02E, 02F, 02L, 02M, 11O, 11P, 12A, 12B, 12G, 12H, 12I, 12P and parts of 01M, 02D), version 6.1, *Open File, NFLD/2616*, Newfoundland and Labrador Dep. of Nat. Resour., Geol. Surv., St. John's, Newfoundland, Canada.
- Comin-Chiaramonti, P., A. Cundari, J. M. DeGraff, C. B. Gomes, and E. M. Piccirillo (1999), Early Cretaceous-Tertiary magmatism in eastern Paraguay (western Parana basin): Geological, geophysical and geochemical relationships, *J. Geodyn.*, 28(4–5), 375–391, doi:10.1016/S0264-3707(99)00016-2.
- Cook, E. (1995), Taphonomy of two non-marine Lower Cretaceous bone accumulations from southeastern England, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 116(3–4), 263–270, doi:10.1016/0031-0182(94)00101-D.
- Coordinating Committee for Geoscience Programmes in East and Southeast Asia (2004), Digital geologic map of East and Southeast Asia, version 2, *Dig. Geosci. Map, G-2*, scale 1:2,000,000, Geol. Surv. of Jpn., Natl. Inst. of Adv. Ind. Sci. and Technol., Tsukuba.
- Copard, Y., P. Amiotte-Suchet, and C. Di-Giovanni (2007), Storage and release of fossil organic carbon related to weathering of sedimentary rocks, *Earth Planet. Sci. Lett.*, 258(1–2), 345–357, doi:10.1016/j.epsl.2007.03.048.
- Cope, J. C. W., J. K. Ingham, and P. F. Rawson (Eds.) (1992), *Atlas of Palaeogeography and Lithofacies*, 153 pp., Geol. Soc., London.
- Corrie, S. L., and M. J. Kohn (2011), Metamorphic history of the central Himalaya, Annapurna region, Nepal, and implications for tectonic models, *Geol. Soc. Am. Bull.*, 123(9–10), 1863–1879, doi:10.1130/B30376.1.
- Corvinus, G., and L. N. Rimal (2001), Biostratigraphy and geology of the neogene Siwalik group of the Surai Khola and Rato Khola areas in Nepal, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 165(3–4), 251–279, doi:10.1016/S0031-0182(00)00163-2.
- Coulthard, T. J. (2001), Landscape evolution models: A software review, *Hydrol. Processes*, 15(1), 165–173, doi:10.1002/hyp.426.
- Cox, B. M., J. D. Hudson, and D. M. Martill (1992), Lithostratigraphic nomenclature of the Oxford Clay (Jurassic), *Proc. Geol. Assoc.*, 103(4), 343–345.
- Crimes, T. R., J. F. G. Hidalgo, and D. G. Poire (1992), Trace fossils from arenig flysch sediments of eire and their bearing on the early colonisation of the deep seas, *Ichnos*, 2(1), 61–77, doi:10.1080/10420949209380076.
- Dahlgren, R. A., F. C. Ugolini, and W. H. Casey (1999), Field weathering rates of Mt. St. Helens tephra, *Geochim. Cosmochim. Acta*, 63(5), 587–598, doi:10.1016/S0016-7037(99)00067-8.
- Dasgupta, A. K., and K. K. Chakravorty (1998), Geological map of India, 4 sheets, 7th ed., Geol. Surv. of India, Calcutta, India.
- Dashzeveg, D., M. J. Novacek, M. A. Norell, J. M. Clark, L. M. Chiappe, A. Davidson, M. C. Mckenna, L. Dingus, C. Swisher, and P. Altangerel (1995), Extraordinary preservation in a new vertebrate assemblage from the Late Cretaceous of Mongolia, *Nature*, 374(6521), 446–449, doi:10.1038/374446a0.

- Davies, D. K. (1967), Origin of friable sandstone: Calcareous sandstone rhythms in the upper Lias of England, *J. Sediment. Petrol.*, 37(4), 1179–1188.
- Deb, M., D. M. Banerjee, and A. K. Bhattacharya (1978), Precambrian stromatolite and other structures in the Rajpura-Dariba polymetallic ore deposit, Rajasthan, India, *Miner. Deposita*, 13(1), 1–9, doi:10.1007/BF00202904.
- Deb, M., R. Thorpe, and D. Krstic (2002), Hindoli group of rocks in the eastern fringe of the Aravalli-Delhi orogenic belt—Archean secondary greenstone belt or Proterozoic supracrustals?, *Gondwana Res.*, 5(4), 879–883, doi:10.1016/S1342-937X(05)70922-9.
- de Caritat, P., C. Reimann, NGS Project Team, and GEMAS Project Team (2012), Comparing results from two continental geochemical surveys to world soil composition and deriving Predicted Empirical Global Soil (PEGS2) reference values, *Earth Planet. Sci. Lett.*, 319–320, 269–276, doi:10.1016/j.epsl.2011.12.033.
- DeVries, T. J. (1988), The geology of late Cenozoic marine terraces (tablazos) in northwestern Peru, *J. South Am. Earth Sci.*, 1(2), 121–136, doi:10.1016/0895-9811(88)90030-2.
- Dheeradilok, P., T. Wongwanich, W. Tansathien, and P. Chaodumrong (1992), An introduction to geology of Thailand, in *National Conference on Geologic Resources of Thailand: Potential for Future Development*, pp. 737–752, Dep. of Miner. Resour., Bangkok.
- Dicken, C. L., S. W. Nicholson, J. D. Horton, S. A. Kinney, G. Gunther, M. P. Foose, and J. A. L. Mueller (2005), Integrated geologic map databases for the United States: Delaware, Maryland, New York, Pennsylvania, and Virginia, U.S. Geol. Surv., Reston, Va.
- Dicken, C. L., S. W. Nicholson, J. D. Horton, M. P. Foose, and J. A. L. Mueller (2007), Preliminary integrated geologic map databases for the United States: Alabama, Florida, Georgia, Mississippi, North Carolina, and South Carolina, U.S. Geol. Surv., Reston, Va.
- Dietrich, V., and A. Gansser (1981), The leucogranites of the Bhutan Himalaya (crustal anatexis versus mantle melting), *Schweiz. Mineral. Petrogr. Mitt.*, 61(2–3), 177–202.
- Direction de l'Industrie, des Mines et de l'Energie (DIMENC) (1981), Carte de la géologie de la Nouvelle-Calédonie, scale 1:200,000, Noumea.
- Dinter, D. A., and L. Royden (1993), Late Cenozoic extension in northeastern Greece: Strymon Valley detachment system and Rhodope metamorphic core complex, *Geology*, 21(1), 45–48, doi:10.1130/0091-7613(1993)021<0045:LCEING>2.3.CO;2.
- Dirección Nacional de Minería y Geología (1985), Carta geológica, scale 1:500,000, Minist. de Ind., Energía y Miner., Montevideo.
- Doebrich, J. L., and R. R. Wahl (2006), Geologic and mineral resource map of Afghanistan, version 2, scale 1:850,000, U.S. Geol. Surv., Reston, Va.
- Dolginow, J., S. Kropatschjow, and E. Klitzsch (1994), *Abriß der Geologie Rußlands und angrenzender Staaten*, 87 pp., Schweizerbart, Stuttgart, Germany.
- Dottin, O. (1990), Geological map of the world, 1:25,000,000, Comm. for the Geol. Map of the World, U. N. Educ., Sci. and Cult. Organ., Paris.
- Douglas, J. A., and W. J. Arkell (1928), The stratigraphical distribution of the Cornbrash: I. The south-western area, *Q. J. Geol. Soc.*, 84(1–4), 117–178, doi:10.1144/GSL.JGS.1928.084.01-04.05.
- Douglas, R. J. W., H. Gabrielse, J. O. Wheeler, D. F. Stott, and H. R. Belyea (1970), Geology of western Canada, in *Geology and Economic Minerals of Canada*, pp. 364–488, Geol. Surv. of Can., Ottawa, Ont.
- Douglas, R. J. W., A. W. Norris, and D. K. Norris (1974), *Geology, Great Slave, District of Mackenzie*, Geol. Surv. of Can., Ottawa, Ont., doi:10.4095/109149.
- Draper, G., P. Mann, and J. F. Lewis (1994), Hispaniola, in *Caribbean Geology: An Introduction*, edited by S. Donovan and T. A. Jackson, pp. 129–150, Univ. of the West Indies Publ. Assoc., Kingston.
- Draper, G., G. Gutierrez, and J. F. Lewis (1996), Thrust emplacement of the Hispaniola peridotite belt: Orogenic expression of the mid-Cretaceous Caribbean arc polarity reversal?, *Geology*, 24(12), 1143–1146, doi:10.1130/0091-7613(1996)024<1143:TEOTHP>2.3.CO;2.
- Dunlap, W. J., and R. Wysoczanski (2002), Thermal evidence for early Cretaceous metamorphism in the Shyok suture zone and age of the Khardung volcanic rocks, Ladakh, India, *J. Asian Earth Sci.*, 20(5), 481–490, doi:10.1016/S1367-9120(01)00042-6.
- Dürr, H. H., M. Meybeck, and S. H. Dürr (2005), Lithologic composition of the Earth's continental surfaces derived from a new digital map emphasizing riverine material transfer, *Global Biogeochem. Cycles*, 19, GB4S10, doi:10.1029/2005GB002515.
- Dürr, H. H., G. G. Laruelle, C. M. van Kempen, C. P. Slomp, M. Meybeck, and H. Middelkoop (2011), Worldwide typology of nearshore coastal systems: Defining the estuarine filter of river inputs to the oceans, *Estuaries Coasts*, 34(3), 441–458, doi:10.1007/s12237-011-9381-y.
- Earth Sciences Research Division (1977), Geological map of the Socialist Republic of the Union of Burma, scale: 1:1,000,000, Burma.
- Economic and Social Commission for Asia and the Pacific (1989), Atlas of mineral resources of the ESCAP region: Sri Lanka, comprising the geological, metamorphic and mineral resources maps of Sri Lanka, scale 1:1,013,760, U. N. Publ., Blue Ridge Summit, Pa. [Available at <http://www.orrbodies.com/>]
- Economic and Social Research Institute (ESRI) (2008), ArcWorld supplement data distributed on CD-ROM with ArcGIS 9.3, Redlands, Calif.
- Edwards, R. A. (1976), Tertiary sediments and structure of the Bovey Basin, south Devon, *Proc. Geol. Assoc.*, 87(1), 1–26.
- Egawa, K., and Y. I. Lee (2009), Jurassic synorogenic basin filling in western Korea: Sedimentary response to inception of the western Circum-Pacific orogeny, *Basin Res.*, 21(4), 407–431, doi:10.1111/j.1365-2117.2009.00408.x.
- Egger, H., H. G. Krenmayer, G. W. Mandl, A. Matura, A. Nowotny, G. Pascher, G. Pestal, J. Pistontnik, M. Rockenschaub, and W. Schnabel (1999), Geologische Übersichtskarte der Republik Österreich, scale 1:1,500,000, Geol. Bundesanstalt, Vienna.
- Egyptian Geological Survey (1981), Geologic map of Egypt, scale 1:2,000,000, Minist. of Ind. and Miner. Resour., Cairo.
- Ernst, R. E., and K. L. Buchan (2004), Igneous rock associations in Canada 3. Large Igneous Provinces (LIPs) in Canada and adjacent regions: 3 Ga to present, *Geosci. Can.*, 31(3), 103–126.
- Escher, J. C., and T. C. R. Pulvertaft (1995), Geological map of Greenland, Geol. Survey of Greenl., Copenhagen.
- Farhoudi, G., and D. E. Karig (1977), Makran of Iran and Pakistan as an active arc system, *Geology*, 5(11), 664–668, doi:10.1130/0091-7613(1977)5<664:MOIAPA>2.0.CO;2.
- Fildani, A., A. D. Hanson, Z. Z. Chen, J. M. Moldovan, S. A. Graham, and P. R. Arriola (2005), Geochemical

- characteristics of oil and source rocks and implications for petroleum systems, Talara basin, northwest Peru, *AAPG Bull.*, 89(11), 1519–1545, doi:10.1306/06300504094.
- Fisher, B. E., and J. C. Poole (2006), Digital version of Nova Scotia Department of Natural Resources map ME 2000-1, geological map of the province of Nova Scotia, version 2, compiled by J. D. Keppie, 2000, *DP ME 43*, scale 1:500 000, Miner. Resour. Branch, Dep. of Nat. Resour., Halifax, N. S., Canada.
- Frank, E. F., C. Wicks, J. Mylroie, J. Troester, E. C. Alexander, and J. L. Carew (1998), Geology of Isla de Mona, Puerto Rico, *J. Cave Karst Stud.*, 60(2), 69–72.
- Frazier, W. J., and D. R. Schwimmer (1987), *Regional Stratigraphy of North America*, 719 pp., Plenum, New York, doi:10.1007/978-1-4613-1795-1.
- Frizzell, D. L. (1943), Upper Cretaceous foraminifera from northwestern Peru, *J. Paleontol.*, 17(4), 331–353.
- Fromaget, J., and E. Saurin (1952), Geological map of Viet-Nam, Cambodia and Laos, 3rd edition, scale 1:2,000,000, Natl. Geogr. Direct. of Vietnam, Đà Lạt.
- Fuchs, G. (1975), *Contributions to the Geology of the North-Western Himalayas*, 64 pp., Geol. Bundesanst., Vienna.
- Fuchs, G., and A. K. Sinha (1978), The tectonics of the Garhwal—Kumaun Lesser Himalaya, *Jahrb. Geol. Bundesanst.*, 121(2), 219–241.
- Furman, T., and D. Graham (1994), Chemical and isotopic variations in volcanic rocks from the Rungwe Province: Constraints on the development and scales of source heterogeneity beneath the African Western Rift, *Mineral. Mag.*, 58A, 297–298, doi:10.1180/minmag.1994.58A.1.156.
- Fürsich, F. T., M. Wilmsen, K. Seyed-Emami, and M. R. Majidifard (2009), Lithostratigraphy of the Upper Triassic–Middle Jurassic Shemshak Group of northern Iran, *Geol. Soc. Spec. Publ.*, 312(1), 129–160, doi:10.1144/SP312.6.
- Fuzesy, L. M. (1979), Geology of Paleozoic strata in west-central Saskatchewan, *AAPG Bull.*, 63(5), 826–827.
- Fyffe, L. R., and D. M. Richard (2007), Carte lithologique du Nouveau-Brunswick, Planche 2007–18, Div. des Polit. et de la Planification, Direction des Ressources Minérales, Minist. des Ressources Nat. du N.-B., Fredericton, N. B., Canada.
- Gallienne, J. (1975), Carte géologique: Martinique: Carte III. (IN) Les ressources en eau de surface de la Martinique, Off. de la Rech. Sci. et Tech. Outre-Mer, Paris.
- Gallois, R. W. (1976), Coccolith blooms in the Kimmeridge Clay and origin of North Sea oil, *Nature*, 259(5543), 473–475, doi:10.1038/259473b0.
- Gallois, R., and S. Etches (2001), The stratigraphy of the youngest part of the Kimmeridge Clay Formation (Upper Jurassic) of the Dorset type area, *Proc. Geol. Assoc.*, 112, 169–182, doi:10.1016/S0016-7878(01)80025-0.
- Ganeshan, T. M., R. K. Chaturvedi, and K. P. Reddy (1982), Some mesozoic plants from the higher Himalayas of Bhutan, *Curr. Sci.*, 51(4), 194–195.
- Garrity, C. P., and D. R. Soller (2009), Database of the geologic map of North America; adapted from the map by J.C. Reed, Jr. and others (2005), U.S. Geol. Surv., Reston, Va.
- Garrity, C. P., P. C. Hackley, and F. Urbani (2006), Digital geologic map and GIS database of Venezuela, U.S. Geol. Surv., Reston, Va.
- Gendzwil, D. J., and S. D. Metieshin (1996), Seismic reflectivity survey of a kimberlite intrusion in the Fort à la Corne district, Saskatchewan, in *Searching for Diamonds in Canada*, edited by A. N. LeCheminant et al., *Geol. Surv. Can. Open File*, 3228, 251–253.
- Geological and Mineral Resources Department (1981), Geological map of the Sudan, Minist. of Energy and Mines, Khartoum.
- Geological Survey and Mines Department (1973), Geological map of Botswana, Geological Survey and Mines Department, Lobatse, Botswana.
- Geological Survey Department Jamaica (1958), Provisional geological map of Jamaica, Kingston.
- Geological Survey Institute of Indonesia (1993), INA GRDC 1:1M combined bedrock and superficial geology and age, Bandung, Indonesia. [Available at portal.onegeology.org]
- Geological Survey of India (1979), *Himalayan Geology Seminar, New Delhi, 1976: A Collection of Papers Presented at the Seminar Held on 13–17 September, 1976 in New Delhi*, Calcutta, India.
- Geological Survey of India (2005), Geological map of the Himalaya, Calcutta, India.
- Geological Survey of India (2009), *Geology and Mineral Resources of India*, 152 pp., Calcutta, India.
- Geological Survey of Ireland (2006), 1:500 000 geological map of Ireland, Dublin.
- Geological Survey of Japan (2003), Geological map of Japan 1:1,000,000, 3rd edition, 2nd CD-ROM version, Tsukuba, Japan.
- Geological Survey of Namibia (1980), Geological map of Namibia, digital version, Windhoek.
- Ghose, N. C., O. P. Agrawal, and N. Chatterjee (2010), Geological and mineralogical study of eclogite and glaucophane schists in the Naga Hills Ophiolite, northeast India, *Isl. Arc*, 19(2), 336–356, doi:10.1111/j.1440-1738.2010.00710.x.
- Gibbs, M. T., and L. R. Kump (1994), Global chemical erosion during the Last Glacial Maximum and the present: Sensitivity to changes in lithology and hydrology, *Paleoceanography*, 9(4), 529–543, doi:10.1029/94PA01009.
- Gibbs, R. J. (1967), Geochemistry of Amazon River system: Part I. Factors that control salinity and composition and concentration of suspended solids, *Geol. Soc. Am. Bull.*, 78(10), 1203–1232, doi:10.1130/0016-7606(1967)78[1203:TGTAR]2.0.CO;2.
- Gleeson, T., L. Smith, N. Moosdorf, J. Hartmann, H. H. Dürr, A. H. Manning, L. P. H. van Beek, and A. M. Jellinek (2011), Mapping permeability over the surface of the Earth, *Geophys. Res. Lett.*, 38, L02401, doi:10.1029/2010GL045565.
- Godderis, Y., C. Roelandt, J. Schott, M. C. Pierret, and L. M. Francois (2009), Towards an integrated model of weathering, climate, and biospheric processes, in *Thermodynamics and Kinetics of Water-Rock Interaction, Rev. in Mineral. and Geochem.*, vol. 70, pp. 411–434, Mineral. Soc. of Am., Washington, D. C.
- Godfrey, L. V., B. Zimmermann, D. C. Lee, R. L. King, J. D. Vervoort, R. M. Sherrell, and A. N. Halliday (2009), Hafnium and neodymium isotope variations in NE Atlantic seawater, *Geochem. Geophys. Geosyst.*, 10, Q08015, doi:10.1029/2009GC002508.
- Gomes, C. B., P. Comin-Chiaromonti, V. F. Velázquez, and D. Orué (1996), Alkaline magmatism in Paraguay: A review, in *Alkaline Magmatism in Central-Eastern Paraguay: Relationships With Coeval Magmatism in Brazil*, edited by P. Comin-Chiaromonti and C. B. Gomes, pp. 31–56, State of São Paulo Res. Found., Sao Paulo.
- Gómez, J., et al. (2007), Geological map of Colombia, scale 1:2,800,000, Inst. Colombiano de Geol. et Miner., Bogotá.
- González, M. E. (2000), Mapa geológico de Paraguay, Viceministerio de Minas y Energía Paraguay, San Lorenzo, Paraguay.

- Goodey, S. P., and A. J. Makepeace (2000), Yukon digital geology, *Open File, D3826*, Explor. and Geol. Serv. Div., Yukon Reg., Indian and Northern Affairs Can., Whitehorse, Yukon Territory, Canada.
- Greensmith, J. T. (1957), Lithology, with particular reference to cementation, of upper Carboniferous sandstones in northern Derbyshire, England, *J. Sediment. Res.*, 27(4), 405–416.
- Grubić, A. (1980), *Yugoslavia: An Outline of Geology of Yugoslavia*, 98 pp., Koeltz Sci., Koenigstein, Germany.
- Gunawan, H., Micheldiament, and V. Mikhailov (2008), Estimation of Bougues density precision: Development of method for analysis of La Soufriere volcano gravity data, *J. Geol. Indonesia*, 3(3), 151–159.
- Haase, D., J. Fink, G. Haase, R. Ruske, M. Pecs, H. Richter, M. Altermann, and K. D. Jager (2007), Loess in Europe—Its spatial distribution based on a European loess map, scale 1:2,500,000, *Quat. Sci. Rev.*, 26(9–10), 1301–1312.
- Hadlari, T., and R. H. Rainbird (2000), Sequence stratigraphy and sedimentology of the Paleoproterozoic Baker Lake Group in the Baker Lake Basin, Thirty Mile Lake, Nunavut, *Rep. 2000-C9*, 10 pp., Geol. Surv. of Can., Ottawa, Ont.
- Hadlari, T., R. H. Rainbird, and J. A. Donaldson (2006), Alluvial, eolian and lacustrine sedimentology of a Paleoproterozoic half-graben, Baker Lake Basin, Nunavut, Canada, *Sediment. Geol.*, 190(1–4), 47–70, doi:10.1016/j.sedgeo.2006.05.005.
- Haeberlin, Y., R. Moritz, L. Fontbote, and M. Cosca (2004), Carboniferous orogenic gold deposits at Pataz, eastern Andean Cordillera, Peru: Geological and structural framework, paragenesis, alteration, and $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology, *Econ. Geol.*, 99(1), 73–112.
- Haghipour, A., and A. Saidi (2010), International geological map of the Middle East, second edition, scale 1:5,000,000, Comm. for the Geol. World Map, Paris.
- Hallam, A., and B. W. Sellwood (1976), Middle Mesozoic sedimentation in relation to tectonics in the British area, *J. Geol.*, 84(3), 301–321, doi:10.1086/628197.
- Hamblin, A. P. (1997), Stratigraphic architecture of the Oldman Formation, Belly River Group, surface and subsurface of southern Alberta, *Bull. Can. Pet. Geol.*, 45(2), 155–177.
- Hambrey, M. J., W. B. Harland, and the International Geological Correlation Programme Project 38: Pre-Pleistocene Tillites (1981), *Earth's Pre-Pleistocene Glacial Record*, 1004 pp., Cambridge Univ. Press, Cambridge, U. K.
- Hamilton, W. N., C. W. Langenberg, and M. Price (2004), Bedrock geology of Alberta, Alberta Geol. Surv., Edmonton, Alberta, Canada.
- Hanzl, P., and Z. Krejci (2005), Geological map of the Trans-Alтай Gobi, scale 1:500,000, Czech Geol. Surv., Prague.
- Hartlaub, R. P., L. M. Heaman, K. E. Ashton, and T. Chacko (2004), The Archean Murmac Bay Group: Evidence for a giant archaic rift in the Rae Province, Canada, *Precambrian Res.*, 131(3–4), 345–372, doi:10.1016/j.precamres.2004.01.001.
- Hartmann, J. (2009), Bicarbonate-fluxes and CO_2 -consumption by chemical weathering on the Japanese Archipelago: Application of a multi-lithological model framework, *Chem. Geol.*, 265(3–4), 237–271, doi:10.1016/j.chemgeo.2009.03.024.
- Hartmann, J., and N. Moosdorf (2011), Chemical weathering rates of silicate-dominated lithological classes and associated liberation rates of phosphorus on the Japanese Archipelago: Implications for global scale analysis, *Chem. Geol.*, 287(3–4), 125–157, doi:10.1016/j.chemgeo.2010.12.004.
- Hartmann, J., N. Jansen, H. H. Dürr, A. Harashima, K. Okubo, and S. Kempe (2010), Predicting riverine dissolved silica fluxes to coastal zones from a hyperactive region and analysis of their first order controls, *Int. J. Earth Sci.*, 99(1), 207–230, doi:10.1007/s00531-008-0381-5.
- Hayes, B. J. R., J. E. Christopher, L. Rosenthal, G. Los, B. McKercher, D. Minken, Y. M. Tremblay, and J. Fennell (1994), Cretaceous Mannville Group of the Western Canada Sedimentary Basin [online], in *Geological Atlas of the Western Canada Sedimentary Basin*, compiled by G. D. Mossop and I. Shetsen, chap. 19, Can. Soc. of Pet. Geol., Calgary, Alberta. [Available at http://www.ags.gov.ab.ca/publications/wcsb_atlas/atlas.htm]
- Hegner, E., and J. S. Pallister (1989), Pb, Sr, and Nd isotopic characteristics of Tertiary Red Sea rift volcanics from the central Saudi Arabian coastal plain, *J. Geophys. Res.*, 94(B6), 7749–7755, doi:10.1029/JB094iB06p07749.
- Helmcke, D. (1985), The Permo-Triassic Paleotethys in mainland Southeast-Asia and adjacent parts of China, *Geol. Rundsch.*, 74(2), 215–228, doi:10.1007/BF01824893.
- Heng, Y. E. (1992), Geological map of Sarawak, second edition, scale 1:500,000, Geol. Surv. of Malaysia, Kuala Lumpur.
- Henriksen, N., A. K. Higgins, F. Kalsbeek, and T. C. R. Pulvertaft (2009), Greenland from Archaean to Quaternary: Descriptive text to the 1995 geological map of Greenland, 1:2,500,000, 2nd edition, report, 126 pp., Geol. Surv. of Denm. and Greenl., Copenhagen.
- Hillebrandt, A. (1970), Die Kreide in der Zentralkordillere östlich von Lima (Peru, Südamerika), *Int. J. Earth Sci.*, 59(3), 1180–1203.
- Hottin, G., and O. F. Ouedraogo (1976), Carte géologique de la République de Haute-Volta, Direct. de la Géol. et des Mines, Ouagadougou.
- Hsü, K. J., and U. Briegel (1991), *Geologie der Schweiz: Ein Lehrbuch für den Einstieg und eine Auseinandersetzung mit den Experten*, 219 pp., Birkhäuser, Basel, Switzerland.
- Iancu, V., T. Berza, A. Seghedi, and M. Marunțiu (2005), Paleozoic rock assemblages incorporated in the South Carpathian Alpine Thrust Belt (Romania and Serbia): A review, *Geol. Belg.*, 8(4), 48–68.
- Institute of Mineral Research and Exploration (1961), Geology map of Turkey, scale 1:500,000, 18 sheets, Ankara.
- Instituto de Geología y Minería (1975), Mapa geológico del Peru, scale 1:1,000,000, Minist. de Energía y Minas, San Borja, Peru.
- Instituto Geológico y Minero de España (1994), Mapa geológico de España, scale 1:1,000,000, Madrid.
- Irish, E. J. W. (1971), *Geology, Southern Plains of Alberta, West of Fourth Meridian*, Geol. Surv. of Can., Ottawa, Ont., doi:10.4095/108960.
- Irwin, D. (2005), *Bedrock Geology of the NWT*, Northwest Territories Geosci. Off., Yellowknife, Northwest Territories, Canada.
- Jacay, J., and T. Sempere (2005), Emplacement levels of the Coastal Batholith in central Peru, paper presented at 6th International Symposium on Andean Geodynamics, Univ. de Barcelona, Barcelona, Spain.
- Jacay, J., V. Alejandro, A. Pino, and T. Sempere (2008), Mesozoic backarc basins in western Peru: A brief summary, paper presented at 7th International Symposium on Andean Geodynamics, Inst. de Rech. pour le Environ., Nice, France.
- Jahn, B.-M., F. Wu, and B. Chen (2000), Granitoids of the Central Asian Orogenic Belt and continental growth in the Phanerozoic, *Spec. Pap. Geol. Soc. Am.*, 350, 181–193.
- Jaillard, E., P. Bengtson, and A. V. Dhondt (2005), Late Cretaceous marine transgressions in Ecuador and northern Peru: A refined stratigraphic framework, *J. South Am. Earth Sci.*, 19(3), 307–323, doi:10.1016/j.jsames.2005.01.006.

- James, N. P., and R. N. Ginsburg (1979), *The Seaward margin Of Belize Barrier and Atoll Reefs: Morphology, Sedimentology, Organism Distribution, and Late Quaternary History*, 191 pp., Blackwell Sci., Oxford, U. K.
- Jansen, N., J. Hartmann, R. Lauerwald, H. H. Dürr, S. Kempe, S. Loos, and H. Middelkoop (2010), Dissolved Silica mobilization in the conterminous USA, *Chem. Geol.*, 270(1–4), 90–109, doi:10.1016/j.chemgeo.2009.11.008.
- Jassim, S. Z., and J. C. Goff (2006), *Geology of Iraq*, 341 pp., Dolin, Prague.
- John, D. T., and P. F. Fisher (1984), The stratigraphical and geomorphological significance of the Red Crag fossils at Netley Heath, Surrey: A review and re-appraisal, *Proc. Geol. Assoc.*, 95(3), 235–247.
- Johns, S. M., and M. D. Young (2006), Bedrock geology and economic potential of the Archean Mary River Group, northern Baffin Island, Nunavut, *Rep. 2006-C5*, 13 pp., Geol. Surv. of Can., Ottawa, Ont.
- Joshi, M. P., A. C. Fanger, and C. W. Brown (1990), *Himalaya, Past and Present*, 399 pp., Shree Almora Book Depot, Almora, India.
- Karpinsky, A. P. (1983), Geological map of the USSR and adjoining water-covered areas, Minist. of Geol. of the USSR, St. Petersburg, Russia.
- Kazmin, V. (1972), Geological map of Ethiopia, Geol. Surv. of Ethiopia, Addis Ababa.
- Kesler, S. E., J. F. Sutter, L. M. Jones, and R. L. Walker (1977), Early Cretaceous basement rocks in Hispaniola, *Geology*, 5(4), 245–247, doi:10.1130/0091-7613(1977)5<245:ECBRIH>2.0.CO;2.
- Keyser, N. (1998), Polygon data of the geological map of the republik of South Africa, Counc. for Geosci. South Africa, Johannesburg.
- Khain, V. Y., A. B. Ronov, and A. N. Balukhovskiy (1981), Neogene lithologic associations of the continents, *Int. Geol. Rev.*, 23(4), 426–454, doi:10.1080/00206818109455078.
- Khan, N. M., M. Abu Bakr, and R. O. Jackson (1964), Geological map of Pakistan, scale 1:2,000,000, Geol. Surv. of Pakistan, Rawalpindi.
- Khin, K., and Myitta (1999), Marine transgression and regression in Miocene sequences of northern Pegu (Bago) Yoma, central Myanmar, *J. Asian Earth Sci.*, 17(3), 369–393, doi:10.1016/S0743-9547(98)00065-8.
- Kim, J. C., and Y. I. Lee (1996), Marine diagenesis of lower Ordovician carbonate sediments (Dumugol Formation), Korea: Cementation in a calcite sea, *Sediment. Geol.*, 105(3–4), 241–257, doi:10.1016/0037-0738(95)00141-7.
- Krishnan, M. S. (1968), *Geology of India and Burma*, 5th ed., 536 pp., Higginbothams, Madras, India.
- Kumar, G., and G. Singh (1983), Middle Cambrian trilobites from Karihul, Liddar valley, Anantnag District, Kashmir and its significance, *Curr. Sci.*, 52(11), 548–549.
- Kumar, G., B. K. Raina, O. N. Bhargava, P. K. Maithy, and R. Babu (1984), The Precambrian-Cambrian boundary problem and its prospects, northwest Himalaya, India, *Geol. Mag.*, 121(3), 211–219, doi:10.1017/S0016756800028272.
- Kumar, P., and A. Pankaj (2009), Role of rainfall in inducing landslides—A case study of Vaishnodevi Area, Jammu and Kashmir, paper presented at Second India Disaster Management Congress, Natl. Inst. of Disast. Manage., New Delhi.
- Kumar, R. (1985), *Fundamentals of Historical Geology and Stratigraphy of India*, 254 pp., John Wiley, New York.
- Lacan, F., and C. Jeandel (2005), Neodymium isotopes as a new tool for quantifying exchange fluxes at the continent-ocean interface, *Earth Planet. Sci. Lett.*, 232(3–4), 245–257, doi:10.1016/j.epsl.2005.01.004.
- Lakhera, R., A. Bhattacharya, and A. Roy (1980), Geology in parts of Sainj Valley, Kulu district, Himachal Pradesh, *J. Indian Soc. Remote Sens.*, 8(1), 53–64.
- Latt, T. T., Z. Win, and K. Ueno (2008), Permian fusuline fauna from the plateau limestone of the Lebyin area, eastern Myanmar: Biochronologic and paleobiogeographic assessments, paper presented at International Symposia on Geoscience Resources and Environments of Asian Terranes, Dep. of Geol., Fac. of Sci., Chulalongkorn Univ., Bangkok, Thailand.
- Ledru, P., V. Johan, J. P. Milési, and M. Tegye (1994), Markers of the last stages of the Palaeoproterozoic collision: Evidence for a 2 Ga continent involving circum-South Atlantic provinces, *Precambrian Res.*, 69(1–4), 169–191, doi:10.1016/0301-9268(94)90085-X.
- Lee, H., and S. Chough (2006), Sequence stratigraphy of Pyeongan Supergroup (Carboniferous-Permian), Taebaek area, mid-east Korea, *Geosci. J.*, 10(4), 369–389, doi:10.1007/BF02910433.
- Lehnert, K., Y. Su, C. H. Langmuir, B. Sarbas, and U. Nohl (2000), A global geochemical database structure for rocks, *Geochim. Geophys. Geosyst.*, 1(5), 1012, doi:10.1029/1999GC000026.
- Le Maitre, R. W. (2004), *Igneous Rocks: A Classification and Glossary of Terms: Recommendations of the International Union of Geological Sciences, Subcommittee on the Systematics of Igneous Rocks*, 2nd ed., 236 pp., Cambridge Univ. Press, Cambridge, U. K.
- Lepvrier, C., M. Faure, V. N. Van, T. Van Vu, W. Lin, T. T. Trong, and P. T. Hoa (2011), North-directed Triassic nappes in northeastern Vietnam (East Bac Bo), *J. Asian Earth Sci.*, 41, 56–68, doi:10.1016/j.jseas.2011.01.002.
- Ligarda, R. C., G. Carlier, and V. C. Carlotto (1993), Petrogenesis and occurrences of gabbroic rocks in the limit eastern cordillera-high plateau in the Abancay deflection area (Curahuasi, south Peru), paper presented at International Symposium on Andean Geodynamics, Univ. of Oxford, Oxford, U. K.
- Lindell, L., M. Astrom, and T. Oberg (2010), Land-use change versus natural controls on stream water chemistry in the Subandean Amazon, Peru, *Appl. Geochem.*, 25(3), 485–495, doi:10.1016/j.apgeochem.2009.12.013.
- Liu, S. F., O. L. Raymond, A. J. Stewart, I. P. Sweet, M. B. Duggan, C. Charlick, D. Phillips, and A. J. Retter (2006), Surface geology of Australia 1:1,000,000 scale, Northern Territory [digital dataset], Geosci. Aust., Canberra, A. C. T., Australia.
- Llinas, R. A. (2005), Síntesis sobre la situación geológica de la Hispaniola, report, Secr. de Estado de Ind. y Comer., Santo Domingo.
- Long, S., N. McQuarrie, T. Tobgay, D. Grujic, and L. Hollister (2011), Geologic map of Bhutan, *J. Maps*, 2011, 184–192.
- Loveman, M. H. (1919), A connecting link between the geology of the northern Shan states and Yunnan, *J. Geol.*, 27(3), 204–211, doi:10.1086/622654.
- Ludington, S., B. C. Moring, R. J. Miller, K. S. Flynn, M. J. Hopkins, and G. A. Haxel (2005), Preliminary integrated geologic map databases for the United States—Western States: California, Nevada, Arizona, Washington, Oregon, Idaho, and Utah, U.S. Geol. Surv., Reston, Va.
- Lustrino, M., and E. Sharkov (2006), Neogene volcanic activity of western Syria and its relationship with Arabian plate kinematics, *J. Geodyn.*, 42(4–5), 115–139, doi:10.1016/j.jog.2006.06.003.

- Lustrino, M., L. Melluso, P. Brotzu, C. B. Gomes, L. Morbidelli, R. Muzio, E. Ruberti, and C. C. G. Tassinari (2005), Petrogenesis of the early Cretaceous Valle Chico igneous complex (SE Uruguay): Relationships with Parand-Etendeka magmatism, *Lithos*, 82(3–4), 407–434, doi:10.1016/j.lithos.2004.07.004.
- Macdonald, R., and W. L. Slimmon (1999), Geological map of Saskatchewan [digital], Sask. Indust. and Resour., Sask. Geol. Surv., Regina, Sask., Canada.
- Macdougall, J. D. S., and J. E. Prentice (1964), Sedimentary environments of the Weald Clay of South Eastern England, in *Deltaic and Shallow Marine Deposits, Proceedings of the Sixth International Congress the Netherlands and Belgium—1963*, edited by L. M. J. U. van Straaten, pp. 257–264, Elsevier, Amsterdam.
- Macfarlane, A., R. B. Sorkhabi, and J. Quade (1999), *Himalaya and Tibet: Mountain Roots to Mountain Tops*, 330 pp., Geol. Soc. of Am., Boulder, Colo.
- Manganelli, A., C. Goso, R. Guerequiz, J. L. F. Turiel, M. G. Valles, D. Gimeno, and C. Perez (2007), Groundwater arsenic distribution in south-western Uruguay, *Environ. Geol.*, 53(4), 827–834, doi:10.1007/s00254-007-0695-9.
- Manten, A. A. (1971), *Silurian Reefs of Gotland*, 593 pp., Elsevier, Amsterdam.
- Marsh, A., and K. Heinemann (2006), Report on the Regional Stratigraphic Framework of Western Saskatchewan—Phase 1, Pet. Technol. Res. Cent., Regina, Sask., Canada.
- Martínez, S., and A. Rojas (2004), Quaternary continental molluscs from northern Uruguay: Distribution and paleoecology, *Quat. Int.*, 114, 123–128, doi:10.1016/S1040-6182(03)00047-8.
- Marzoli, A., E. M. Piccirillo, P. R. Renne, G. Bellieni, M. Iacumin, J. B. Nyobe, and A. T. Tongwa (2000), The Cameroon Volcanic Line revisited: Petrogenesis of continental basaltic magmas from lithospheric and asthenospheric mantle sources, *J. Petrol.*, 41(1), 87–109, doi:10.1093/petrology/41.1.87.
- Masquelin, H., T. Aifa, R. Muzio, E. Hallot, G. Veroslavsky, and L. Bonneville (2009), The Cuaro Mesozoic doleritic dyke swarm, southern Parana basin, Uruguay: Examples of superimposed magnetic fabrics?, *C. R. Geosci.*, 341(12), 1003–1015, doi:10.1016/j.crte.2009.07.004.
- Massey, N. W. D., D. G. MacIntyre, P. J. Desjardins, and R. T. Cooney (2005), Digital map of British Columbia: Whole province, B. C. Minist. of Energy and Mines, Victoria, B. C., Canada.
- Mast, M. A., J. I. Drever, and J. Baron (1990), Chemical weathering in the Loch Vale watershed, Rocky Mountain National Park, Colorado, *Water Resour. Res.*, 26(12), 2971–2978, doi:10.1029/WR026i012p02971.
- Mathalone, J. M. P., and M. R. Montoya (1995), Petroleum geology of the Sub-Andean basins of Peru, in *Petroleum Basins of South America*, edited by A. J. Tankard, R. Suarez Soruco, and H. J. Welsink, pp. 423–444, Geol. Soc., London, doi:10.1306/D9CB6E11-1715-11D7-8645000102C1865D.
- Mazumdar, A., and S. K. Bhattacharya (2004), Stable isotopic study of late neoproterozoic-early Cambrian (?) sediments from Nagaur-Ganganagar basin, western India: Possible signatures of global and regional C-isotopic events, *Geochem. J.*, 38(2), 163–175, doi:10.2343/geochemj.38.163.
- McAuliffe, J. R. (1994), Landscape evolution, soil formation, and ecological patterns and processes in Sonoran Desert Bajadas, *Ecol. Monogr.*, 64(2), 111–148, doi:10.2307/2937038.
- McClure, H. A. (1978), Early Paleozoic glaciation in Arabia, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 25(4), 315–326, doi:10.1016/0031-0182(78)90047-0.
- McMechan, M. E., and F. M. Dawson (1995), Geology, wapiti, west of Sixth Meridian, Geol. Surv. of Can., Edmonton, Alberta, Canada.
- Mégard, F., and H. Philip (1976), Plio-Quaternary tectonomagmatic zonation and plate tectonics in central Andes, *Earth Planet. Sci. Lett.*, 33(2), 231–238, doi:10.1016/0012-821X(76)90229-6.
- Mégard, F., D. C. Noble, E. H. McKee, and H. Bellon (1984), Multiple pulses of Neogene compressive deformation in the Ayacucho intermontane basin, Andes of central Peru, *Geol. Soc. Am. Bull.*, 95(9), 1108–1117, doi:10.1130/0016-7606(1984)95<1108:MPONCD>2.0.CO;2.
- Mehrabi, B., B. W. D. Yardley, and J. R. Cann (1999), Sediment-hosted disseminated gold mineralisation at Zarshuran, NW Iran, *Miner. Deposita*, 34(7), 673–696, doi:10.1007/s001260050227.
- Mesolella, K. J., R. K. Matthews, W. S. Broecker, and D. L. Thurber (1969), The astronomical theory of climatic change: Barbados data, *J. Geol.*, 77(3), 250–274, doi:10.1086/627434.
- Metcalfe, I. (1990), Stratigraphic and tectonic implications of Triassic conodonts from northwest peninsular Malaysia, *Geol. Mag.*, 127(6), 567–578, doi:10.1017/S0016756800015454.
- Meybeck, M. (1987), Global chemical weathering of surficial rocks estimated from river dissolved loads, *Am. J. Sci.*, 287(5), 401–428, doi:10.2475/ajs.287.5.401.
- Michel, P. (1973), *Géologie des régions traversées par les fleuves Sénégal et Gambie*, Off. de la Rech. Sci. et Tech. d’Outre-Mer, Paris.
- Millard, M. J. (1996), Geology and groundwater resources of the Hudson Bay area, Saskatchewan, *Rep. 63C-D*, Sask. Res. Council, Saskatoon, Sask., Canada.
- Milodowski, A. E., and J. A. Zalasiewicz (1991), Redistribution of rare earth elements during diagenesis of turbidite/hemipelagite mudrock sequences of Llandovery age from central Wales, *Geol. Soc. Spec. Publ.*, 57(1), 101–124, doi:10.1144/GSL.SP.1991.057.01.10.
- Mineral Resources Authority of Mongolia (1998), Geological map of Mongolia, scale 1:1,000,000, Ulanbaatar.
- Ministerio de Medio Ambiente y Recursos Naturales (2002), Mapa geológico de la Republica de El Salvador, San Salvador.
- Ministry of Geology and Mineral Resources of the People’s Republic of China (1991), Geological map of Nei Mongol Autonomous Region, People’s Republic of China, scale 1:1,500,000, Geol. Publ. House, Beijing, doi:10.1016/S0012-1821X(1002)00774-00774.
- Mishra, D. (2009), High energy transgressive deposits from the Late Jurassic of Wagad, eastern Kachchh, India, *J. Asian Earth Sci.*, 34(3), 310–316, doi:10.1016/j.jseae.2008.05.010.
- Mitchell, A. H. G. (1992), Late Permian-Mesozoic events and the Mergui group Nappe in Myanmar and Thailand, *J. Southeast Asian Earth Sci.*, 7(2–3), 165–178, doi:10.1016/0743-9547(92)90051-C.
- Mitchell, A. H. G., F. Hernandez, and A. P. dela Cruz (1986), Cenozoic evolution of the Philippine archipelago, *J. Southeast Asian Earth Sci.*, 1(1), 3–22, doi:10.1016/0743-9547(86)90003-6.
- Moll, S. J., S. Bie, D. Peterson, D. C. Pray, F. H. Wilson, J. M. Schmidt, J. R. Riehele, and T. P. Miller (1997), Polygon data from 1980 “Geologic map of Alaska,” U.S. Geol. Surv., Reston, Va.

- Mollock, D. I. J. (1974), Geological map of the New Hebrides condominium, scale 1:1,000,000, Geol. Surv. of the New Hebrides, Port Vila.
- Moores, E. M., and R. W. Fairbridge (1997), *Encyclopedia of European and Asian Regional Geology*, 804 pp., Chapman and Hall, London.
- Moosdorf, N., J. Hartmann, and H. H. Durr (2010), Lithological composition of the North American continent and implications of lithological map resolution for dissolved silica flux modeling, *Geochem. Geophys. Geosyst.*, *11*, Q11003, doi:10.1029/2010GC003259.
- Moosdorf, N., J. Hartmann, R. Lauerwald, B. Hagedorn, and S. Kempe (2011), Atmospheric CO₂ consumption by chemical weathering in North America, *Geochim. Cosmochim. Acta*, *75*(24), 7829–7854, doi:10.1016/j.gca.2011.10.007.
- Moshrif, M. A. (1984), Sequential development of Hanifa formation (Upper Jurassic) paleoenvironments and paleogeography, central Saudi Arabia, *J. Pet. Geol.*, *7*(4), 451–460, doi:10.1111/j.1747-5457.1984.tb00889.x.
- Ministère des Ressources Naturelles (2002), Carte géologique du Québec, scale 1:2,000,000, Quebec, Que., Canada.
- Mukherjee, A. K., M. M. Alam, S. K. Mazumdar, R. Haque, and S. Gowrisankaran (1992), Physico-chemical properties and petrographic characteristics of the Kapurdi lignite deposit, Barmer Basin, Rajasthan, India, *Int. J. Coal Geol.*, *21*(1–2), 31–44, doi:10.1016/0166-5162(92)90034-T.
- Muzio, R., E. Peel, E. Morales, G. Veroslavsky, and B. Conti (2009), Mesozoic magmatism in east Uruguay: Petrological constraints related to the Sierra San Miguel Region, *Earth Sci. Res. J.*, *13*(1), 16–29.
- Myroie, J. E., and J. L. Carew (1995), Geology and karst geomorphology of San Salvador Island, Bahamas, *Carbonate Evaporite*, *10*(2), 193–206, doi:10.1007/BF03175404.
- Myrow, P. M., K. R. Thompson, N. C. Hughes, T. S. Paulsen, B. K. Sell, and S. K. Parcha (2006), Cambrian stratigraphy and depositional history of the northern Indian Himalaya, Spiti Valley, north-central India, *Geol. Soc. Am. Bull.*, *118*(3–4), 491–510, doi:10.1130/B25828.1.
- Nag, S., S. K. Sengupta, R. K. Gaur, and A. Absar (1999), Alkaline rocks of Samchampi-Samteran, District Karbi-Anglong, Assam, India, *J. Earth Syst. Sci.*, *108*(1), 33–48.
- Natural Resources Project (1990), Geological map, Republic of Yemen, scale 1:1,000,000, Minist. of Oil and Miner. Resour., Sana.
- Nayak, B., A. K. Singh, A. K. Upadhyay, and K. K. Bhattacharyya (2009), A note on the characters of some Lower Gondwana coals of West Siang District in the Arunachal Himalaya and their trace element content, *J. Geol. Soc. India*, *74*(3), 395–401, doi:10.1007/s12594-009-0138-1.
- Nehlig, P., J. P. Quinquis, M. Bucelle, and O. Odon (2006), Carte géologique de la Réunion, Bur. de Rech. Geol. et Min., Orleans, France.
- Németh, K., K. Suwesi, Z. Peregi, Z. Gulácsi, and J. Ujszászi (2003), Plio/Pleistocene flood basalt related scoria and spatter cones, rootless lava flows, and pit craters, Al Haruj Al Abiyad, Libya, *Geolines*, *15*, 98–103.
- New Zealand Geological Survey (1972), Geological map of New Zealand, scale 1:1,000,000, Dep. of Sci. and Ind. Res., Wellington, New Zealand.
- Nicholson, S. W., C. L. Dicken, J. D. Horton, K. A. Labay, M. P. Foote, and J. A. L. Mueller (2005), Preliminary integrated geologic map databases for the United States: Kentucky, Ohio, Tennessee, and West Virginia, U.S. Geol. Surv., Reston, Va.
- Nicholson, S. W., C. L. Dicken, J. D. Horton, M. P. Foote, J. A. L. Mueller, and R. Hon (2006), Preliminary integrated geologic map databases for the United States: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, Rhode Island, Vermont, U.S. Geol. Surv., Reston, Va.
- Nicholson, S. W., C. L. Dicken, M. P. Foote, and J. A. L. Mueller (2007), Preliminary integrated geologic map databases for the United States: The upper Midwest states: Minnesota, Wisconsin, Michigan, Illinois and Indiana, U.S. Geol. Surv., Reston, Va.
- Noble, D. C., E. H. McKee, E. Farrar, and U. Petersen (1974), Episodic Cenozoic volcanism and tectonism in Andes of Peru, *Earth Planet. Sci. Lett.*, *21*(2), 213–220, doi:10.1016/0012-821X(74)90057-0.
- Noble, D. C., E. Farrar, and E. J. Cobbing (1979), Nazca group of south-central Peru—Age, source, and regional volcanic and tectonic significance, *Earth Planet. Sci. Lett.*, *45*(1), 80–86, doi:10.1016/0012-821X(79)90109-2.
- Noble, D. C., E. H. McKee, T. Mourier, and F. Megard (1990), Cenozoic stratigraphy, magmatic activity, compressive deformation, and uplift in northern Peru, *Geol. Soc. Am. Bull.*, *102*(8), 1105–1113, doi:10.1130/0016-7606(1990)102<1105:CSMACD>2.3.CO;2.
- Nyagah, K. (1995), Stratigraphy, depositional history and environments of deposition of Cretaceous through Tertiary strata in the Lamu Basin, southeast Kenya and implications for reservoirs for hydrocarbon exploration, *Sediment. Geol.*, *96*(1–2), 43–71, doi:10.1016/0037-0738(94)00126-F.
- Okay, A. I., O. Monod, and P. Monie (2002), Triassic blueschists and eclogites from northwest Turkey: Vestiges of the Paleo-Tethyan subduction, *Lithos*, *64*(3–4), 155–178, doi:10.1016/S0024-4937(02)00200-1.
- Okay, A. I., E. Bozkurt, M. Satir, E. Yigitbas, Q. G. Crowley, and C. K. Shang (2008), Defining the southern margin of Avalonia in the Pontides: Geochronological data from the Late Proterozoic and Ordovician granitoids from NW Turkey, *Tectonophysics*, *461*(1–4), 252–264, doi:10.1016/j.tecto.2008.02.004.
- Okulitch, A. V. (2006), Phanerozoic bedrock geology, Lake Athabasca, Alberta, Sask. Geol. Surv., Regina, Sask., Canada.
- Ollerenshaw, N. C. (1970), Marble Mountain, Alberta, Geol. Surv. of Can., Ottawa, Ont.
- Ontario Geological Survey (1993), Bedrock geology, seamless coverage of the province of Ontario, *Data Set*, *6*, Sudbury, Ont., Canada.
- Ortega, E., P. Freile, R. Longo, and J. Baldock (1982), National Geological Map of the Republic of Ecuador, scale 1:1,000,000, Minist. de Recursos Nat. y Energéticos, Quito.
- Palmer, R. H. (1945), Outline of the geology of Cuba, *J. Geol.*, *53*(1), 1–34, doi:10.1086/625242.
- Pande, I. C., and R. Kumar (1965), Origin of Katni bauxite by alteration of Semri series, *Curr. Sci.*, *34*(7), 214–215.
- Paraeija, J., and R. Ballon (1978), Mapa geológico de Bolivia, scale 1:1,000,000, Serv. Geol. de Bolivia, La Paz.
- Patwardhan, A. M. (2010), *The Dynamic Earth System*, 433 pp., Prentice-Hall of India, New Delhi.
- Paul, D., S. Hanmer, S. Tella, T. D. Peterson, and A. N. LeCheminant (2002), Compilation, bedrock geology of part of the Western Churchill Province, *Open File*, *4236*, Geol. Surv. of Can., Ottawa, Ont., doi:10.4095/213530.
- Pawlewicz, M. J., D. W. Steinshouer, and D. L. Gautier (1997), Map showing geology, oil and gas fields, and geologic provinces of Europe including Turkey, *U.S. Geol. Surv. Open File Rep.*, *97-470I*, 14 pp.

- Pécsi, M. (1990), Loess is not just the accumulation of dust, *Quat. Int.*, 14(7–8), 1–21, doi:10.1016/1040-6182(90)90034-2.
- Persits, F. M., C. J. Wandrey, R. C. Milici, and A. Manwar (2001), Digital geologic and geophysical data of Bangladesh, *U.S. Geol. Surv. Open File Rep.*, OFR-97-470-H.
- Persits, F. M., T. S. Ahlbrandt, M. L. Tuttle, R. R. Charpentier, M. E. Brownfield, and K. I. Takahashi (2002), Surficial geology of Africa (geo7_2ag), U.S. Geol. Surv., Denver, Colo.
- Peter, H. (1986), The Lower Lias (Lower Jurassic) of the Bridgend area, South Wales, *Proc. Geol. Assoc.*, 97(3), 237–242.
- Petersen, U., D. C. Noble, M. J. Arenas, and P. C. Goodell (1977), Geology of Julcani mining district, Peru, *Econ. Geol.*, 72(6), 931–949, doi:10.2113/gsecongeo.72.6.931.
- Petterson, M. G., et al. (1999), Geological-tectonic framework of Solomon Islands, SW Pacific: Crustal accretion and growth within an intra-oceanic setting, *Tectonophysics*, 301(1–2), 35–60, doi:10.1016/S0040-1951(98)00214-5.
- Pharaoh, T. C., and J. N. Carney (2000), An introduction to the Precambrian rocks of England and Wales, in *Precambrian Rocks of England and Wales*, edited by J. N. Carney et al., pp. 1–17, Nat. Conserv. Comm., Peterborough, U. K.
- Pinna, P., P. Marteau, J.-F. Becq-Giraudon, and B. Manigault (1987), Carta geológica, República Popular de Moçambique, Bur. de Rech. Géol. et Min., Orleans, France.
- Pollastro, R. M., A. S. Karshbaum, and R. J. Viger (1997a), Maps showing geology, oil and gas fields and geologic provinces of the Arabian Peninsula, scale 1:4,500,000, U.S. Geol. Surv., Reston, Va.
- Pollastro, R. M., F. M. Persits, and D. W. Steinshouer (1997b), Map showing geology, oil and gas fields, and geologic provinces of Iran, 10 pp., U.S. Geol. Surv., Reston, Va.
- Pollock, S. M., F. Goodarzi, and C. L. Riediger (2000), Mineralogical and elemental variation of coal from Alberta, Canada: An example from the No. 2 seam, Genesee Mine, *Int. J. Coal Geol.*, 43(1–4), 259–286, doi:10.1016/S0166-5162(99)00063-4.
- Porder, S., P. M. Vitousek, O. A. Chadwick, C. P. Chamberlain, and G. E. Hilley (2007), Uplift, erosion, and phosphorus limitation in terrestrial ecosystems, *Ecosystems*, 10(1), 159–171, doi:10.1007/s10021-006-9011-x.
- Portugal, J. A. (1974), Mesozoic and Cenozoic stratigraphy and tectonic events of Puno-Santa Lucia area, Dept. of Puno, Peru, *AAPG Bull.*, 58(6), 982–999.
- Powers, D. L. (1931), Subsurface study of pale beds and foremost formation in Lethbridge-Brooks area of southern Alberta, *AAPG Bull.*, 15, 1197–1213.
- Prasad, G. (1983), A review of the early Tertiary bauxite event in South America, Africa and India, *J. Afr. Earth Sci.*, 1(3–4), 305–313.
- Preciozzi Porta, F. (1993), Petrography and geochemistry of five granitic plutons from South-central Uruguay: Contribution to the knowledge of the Piedra Alta terrane = (Petrographie et geochemie de cinq plutons granitiques du Centre-Sud de l'Uruguay: Contribution a la connaissance du terrain Piedra Alta), thèse de doctorat, 189 pp., Univ. du Québec à Chicoutimi, Chicoutimi, Que., Canada.
- Pruett, R. J., and H. H. Murray (1991), Clay mineralogy, alteration history, and economic geology of the whitemud formation, southern Saskatchewan, Canada, *Clays Clay Miner.*, 39(6), 586–596, doi:10.1346/CCMN.1991.0390604.
- Purser, B. H. (1998), *Sedimentation and Tectonics in Rift Basins: Red Sea–Gulf of Aden*, 1st ed., 663 pp., Chapman and Hall, London.
- Qari, M. Y. H. T. (1989), Lithological mapping and structural analysis of Proterozoic rocks in part of the southern Arabian shield using Landsat images, *Int. J. Remote Sens.*, 10(3), 499–503, doi:10.1080/01431168908903887.
- Radhakrishna, T., V. D. Rao, and A. V. Murali (1984), Geochemistry of Dras Volcanics and the evolution of the Indus Suture Ophiolites, *Tectonophysics*, 108(1–2), 135–153, doi:10.1016/0040-1951(84)90157-4.
- Rai, S., M. Yoshida, B. Upreti, and P. Ulakm (2007), Geology of the Lesser Himalayan and Higher Himalayan Crystalline sequences of the Everest area along the Dudh Koshi valley, eastern Nepal Himalaya, *J. Nepal Geol. Soc.*, 36, 11.
- Ramirez, M. C., and H. Cisneros (2007), *Andean System of Basins: Watershed Profiles*, 105 pp., Int. Potato Cent., Lima.
- Rana, R. S., K. Kumar, H. Singh, and K. D. Rose (2005), Lower vertebrates from the Late Palaeocene–Earliest Eocene Akli Formation, Giral Lignite Mine, Barmer District, western India, *Curr. Sci.*, 89(9), 1606–1613.
- Raymond, O. L., S. F. Liu, and P. Kilgour (2007a), Surface geology of Australia 1:1,000,000 scale, Tasmania—3rd edition [digital dataset], Geosci. Australia, Canberra, A. C. T., Australia.
- Raymond, O. L., S. F. Liu, P. Kilgour, A. J. Retter, and D. P. Connolly (2007b), Surface geology of Australia 1:1,000,000 scale, Victoria—3rd edition [digital dataset], Geosci. Aust., Canberra, A. C. T., Australia.
- Raymond, O. L., S. F. Liu, P. Kilgour, A. J. Retter, A. J. Stewart, and G. Stewart (2007c), Surface geology of Australia 1:1,000,000 scale, New South Wales—2nd edition [digital dataset], Geosci. Aust., Canberra, A. C. T., Australia.
- Riccardi, A. C., C. A. Gulisano, J. Mojica, O. Palacio, C. Schubert, and M. R. A. Thomson (1992), Western South America and Antarctica, in *The Jurassic of the Circum-Pacific*, edited by G. E. G. Westermann, pp. 122–161, Cambridge Univ. Press, New York.
- Richards, H. C. (1974), *Annotated Bibliography of Quaternary Shorelines, Second Supplement of 1970–1973*, 214 pp., Acad. of Nat. Sci., Philadelphia, Pa.
- Richardson, R. J. H., C. W. Langenberg, D. Chao, and D. W. Fietz (1990), Coal compilation project: Entrance, NTS 83F/5, *Open File Rep.*, 1990–02, 1 p., Alberta Geol. Surv., Edmonton, Alberta, Canada.
- Robertson, A., and M. Shallo (2000), Mesozoic–Tertiary tectonic evolution of Albania in its regional eastern Mediterranean context, *Tectonophysics*, 316(3–4), 197–254, doi:10.1016/S0040-1951(99)00262-0.
- Rod, E., and W. Maync (1954), Revision of Lower Cretaceous stratigraphy of Venezuela, *AAPG Bull.*, 38(2), 193–283.
- Rodda, P. (1967), Outline of the geology of Viti Levu, *N. Z. J. Geol. Geophys.*, 10(5), 1260–1273, doi:10.1080/00288306.1967.10420217.
- Rushton, A. W. A. (1979), A review of the Middle Cambrian Agnostida from the Abbey Shales, England, *Alcheringa Australasian J. Palaeontol.*, 3(1), 43–61, doi:10.1080/03115517908565439.
- Saadi, M. (1982), Carte structurale du Maroc, Ministère de l'Énergie et des Mines, Rabat.
- Sabir Khan, M. (1994), Petrology and geochemistry of the Panjal Volcanics in the Azad Kashmir and Kaghan areas of the NW Himalaya, Univ. of the Punjab, Lahore, Pakistan.
- Sandeman, H. A., A. H. Clark, E. Farrar, and G. Arroyo-Pauca (1996), A critical appraisal of the Cayconi Formation, Crucero Basin, southeastern Peru, *J. South Am. Earth Sci.*, 9(5–6), 381–392, doi:10.1016/S0895-9811(96)00021-1.

- Schau, M. (1993), Geology, northern Melville Peninsula, District of Franklin, Northwest Territories, Geol. Surv. of Can., Ottawa, Ont., doi:10.4095/183869.
- Schefer, S., V. Cvetkovic, B. Fugenschuh, A. Kounov, M. Ovtcharova, U. Schaltegger, and S. M. Schmid (2011), Cenozoic granitoids in the Dinarides of southern Serbia: Age of intrusion, isotope geochemistry, exhumation history and significance for the geodynamic evolution of the Balkan Peninsula, *Int. J. Earth Sci.*, 100(5), 1181–1206, doi:10.1007/s00531-010-0599-x.
- Schledewitz, D. C. P., and D. Lindal (2005), Bedrock geology compilation map series, NTS 54M, scale 1:250,000, Manitoba Geol. Surv., Winnipeg, Manitoba, Canada.
- Schlüter, T. (2005), *Geological Atlas of Africa*, 1st ed., 272 pp., Springer, Berlin.
- Schobbenhaus, C., and A. Bellizia (2001), Geological map of South America, scale 1:5,000,000, Comm. for the Geol. Map of the World, Brasília.
- Schruben, P. G. (1996), Geology and resource assessment of Costa Rica at 1:500,000 scale—A digital representation of maps of the U.S. Geological Survey's 1987 Folio I-1865, U.S. Geol. Surv., Reston, Va.
- Scott, D. J., and E. A. de Kemp (1998), Bedrock geology compilation, northern Baffin Island and northern Melville Peninsula, Northwest Territories, Geol. Surv. of Can., Ottawa, Ont., doi:10.4095/210024.
- Serviço Geológico do Brasil (2004), Carta geológica do Brasil ao Milionésimo: Sistema de informações geográficas-SIG, Brasília.
- Servicio Geológico Mexicano (2007), Carta Geológica-Minera Estatal Distrito Federal, scale 1:500,000, Pachuca, Mexico.
- Servicio Geológico Minero Argentino (1997), Mapa geológico de la República Argentina, scale 1:2,500,000, Buenos Aires.
- Servicio Nacional de Geología y Minería (2002), Mapa geológico de Chile, Santiago.
- Servicio Nacional de Geología y Minería (2004), Mapa geológico de Chile: Versión digital, Santiago.
- Seyed-Emami, K. (2003), Triassic in Iran, *Facies*, 48, 91–106, doi:10.1007/BF02667532.
- Sharief, F. A. (1982), Lithofacies distribution of the Permian-Triassic rocks in the Middle East, *J. Pet. Geol.*, 4(3), 299–310, doi:10.1111/j.1747-5457.1982.tb00541.x.
- Sharma, B. K., and A. M. Bhola (2005), Kink bands in the Chamba region, western Himalaya, India, *J. Asian Earth Sci.*, 25(3), 513–528, doi:10.1016/j.jseae.2004.05.003.
- Sharma, K. K. (2007), K-T magmatism and basin tectonism in western Rajasthan, India, results from extensional tectonics and not from Réunion plume activity, *Spec. Pap. Geol. Soc. Am.*, 430, 775–784.
- Sharma, R. S., and H. Thomas (2005), *Metamorphism and Crustal Evolution: Papers in Honour of Prof. R. S. Sharma*, Atlantic, New Delhi.
- Shekhawat, M. S., and G. Prabhulingaiah (2010), Mineralogical characteristics and mineral economics of Sanu and Gotan SMS grade limestone deposits of Rajasthan, *J. Geol. Soc. India*, 75(5), 739–748, doi:10.1007/s12594-010-0060-6.
- Sherrod, D. R., J. M. Sinton, S. E. Watkins, and K. M. Brunt (2007), Geologic map of the state of Hawaii, *U.S. Geol. Surv. Open File Rep.*, 2007-1089.
- Sigmond, E. M. O. (2002), Land and sea areas of northern Europe, digital version, scale 1:4,000,000, Geol. Surv. of Norway, Trondheim.
- Sim, M. S., and Y. I. Lee (2006), Sequence stratigraphy of the Middle Cambrian Daegi Formation (Korea), and its bearing on the regional stratigraphic correlation, *Sediment. Geol.*, 191(3–4), 151–169, doi:10.1016/j.sedgeo.2006.03.016.
- Singh, A. K. (2010), Mafic dykes/sills in Siang window of eastern syntaxial bend, eastern Himalaya, India: Geochemistry and petrogenetic aspects, paper presented at 6th International Dyke Conference, Banaras Hindu Univ., Varanasi, India.
- Singh, M. K. (2011), Assessing the relative packing behaviour of Chandarpur and Khairagarh sandstones of Proterozoic Chhattisgarh Basin, *J. Geol. Soc. India*, 77(1), 35–41, doi:10.1007/s12594-011-0006-7.
- Singh, S., and A. K. Jain (2003), Himalayan granitoids, *J. Virtual Explor.*, 11, paper 01.
- Sinha, A. K. (1989), *Geology of the Higher Central Himalaya*, 219 pp., John Wiley, Chichester, U. K.
- Sinha, A. K., and S. P. Nautiyal (1981), *Contemporary Geoscientific Researches in Himalaya: A Commemorative Volume in Honour of S.P. Nautiyal*, 168 pp., Bishen Singh Mahendra Pal Singh, Dehra Dun, India.
- Sinha, A. K., H. Rai, R. Upadhyay, and R. Chandra (1999), Contribution to the geology of the eastern Karakorum, India, in *Himalaya and Tibet: Mountain Roots to Mountain Tops*, edited by A. Macfarlane, R. B. Sorkhabi, and J. Quade, pp. 33–46, Geol. Soc. of Am., Boulder, Colo., doi:10.1130/0-8137-2328-0.33.
- Smith, S. A., and R. A. Edwards (1991), Regional sedimentological variations in Lower Triassic fluvial conglomerates (Budleigh Salterton pebble beds), southwest England—Some implications for paleogeography and basin evolution, *Geol. J.*, 26(1), 65–83, doi:10.1002/gj.3350260105.
- Snoke, A. W. (2001), *Petrologic and Structural History of Tobago, West Indies: A Fragment of the Accreted Mesozoic Oceanic Arc of the Southern Caribbean*, 54 pp., Geol. Soc. of Am., Boulder, Colo., doi:10.1130/0-8137-2354-X.1.
- Soliman, S. M., and M. A. Elfetouh (1970), Carboniferous of Egypt—Isopach and lithofacies maps, *AAPG Bull.*, 54(10), 1918–1930.
- Sotiropoulos, S., M. V. Triantaphyllou, E. Kamberis, and S. Tsaila-Monopolis (2008), Paleogene terrigenous (flysch) sequences in Etoloakarnania region (W. Greece). Plankton stratigraphy and paleoenvironmental implications, *Geobios*, 41(3), 415–433, doi:10.1016/j.geobios.2007.10.007.
- Srimal, N. (2005), Abor volcanics: Slab window volcanism at the India-Asia collision zone, paper presented at Annual Meeting, Geol. Soc. of Am., Salt Lake City, Utah.
- Stead, D. (1990), Engineering geology in Papua New Guinea—A review, *Eng. Geol.*, 29(1), 1–29, doi:10.1016/0013-7952(90)90079-G.
- Steinsouer, D. W., J. Qiang, P. J. McCabe, and R. T. Ryder (1999), Maps Showing geology, oil and gas fields, and geologic provinces of the Asia Pacific region; Plate 2: Southeast Asia, Cent. Energy Resour. Team, U.S. Geol. Surv., Denver, Colo.
- Stewart, A. J., I. P. Sweet, R. S. Needham, O. L. Raymond, A. J. Whitaker, S. F. Liu, D. Phillips, A. J. Retter, D. P. Connolly, and G. Stewart (2008), Surface geology of Australia 1:1,000,000 scale, Western Australia [digital dataset], Geosci. Aust., Canberra, A. C. T., Australia.
- Stoeser, D. B., G. N. Green, L. C. Morath, W. D. Heran, A. B. Wilson, D. W. Moore, and B. S. Van Gosen (2005), Preliminary integrated geologic map databases for the United States central States: Montana, Wyoming, Colorado, New Mexico, Kansas, Oklahoma, Texas, Missouri, Arkansas, and Louisiana, North Dakota, South Dakota, Nebraska, and Iowa, U.S. Geol. Surv., Reston, Va.

- Streule, M. J., R. J. Phillips, M. P. Searle, D. J. Waters, and M. S. A. Horstwood (2009), Evolution and chronology of the Pangong Metamorphic Complex adjacent to the Karakoram Fault, Ladakh: Constraints from thermobarometry, metamorphic modelling and U-Pb geochronology, *J. Geol. Soc.*, *166*(5), 919–932, doi:10.1144/0016-76492008-117.
- Stubbley, M. (2005), Bedrock geology of the Slave Craton, Northwest Territories Geosci. Off., Yellowknife, Northwest Territories, Canada.
- Sundaram, R., R. A. Henderson, K. Ayyasami, and J. D. Stilwell (2001), A lithostratigraphic revision and palaeoenvironmental assessment of the Cretaceous System exposed in the onshore Cauvery Basin, southern India, *Cretaceous Res.*, *22*(6), 743–762, doi:10.1006/cres.2001.0287.
- Sur, S., J. Schieber, and S. Banerjee (2006), Petrographic observations suggestive of microbial mats from Rampur Shale and Bijaigarh Shale, Vindhyan basin, India, *J. Earth Syst. Sci.*, *115*(1), 61–66, doi:10.1007/BF02703026.
- Szabó, C., S. Harangi, and L. Csontos (1992), Review of Neogene and Quaternary volcanism of the Carpathian-Pannonian region, *Tectonophysics*, *208*(1–3), 243–256, doi:10.1016/0040-1951(92)90347-9.
- Szekely, T. S. (1966), Correlation of Mesozoic formations of southern Peru and northern Chile: Geological notes, *AAPG Bull.*, *50*, 1994–1998.
- Takai, F., T. Matsumoto, and R. Toriyama (Eds.) (1963), *Geology of Japan*, 279 pp., Univ. of Calif. Press, Berkeley.
- Taylor, G. K., J. Grocott, A. Pope, and D. E. Randall (1998), Mesozoic fault systems, deformation and fault block rotation in the Andean forearc: A crustal scale strike-slip duplex in the Coastal Cordillera of northern Chile, *Tectonophysics*, *299*(1–3), 93–109, doi:10.1016/S0040-1951(98)00200-5.
- Taylor, P. J., and W. A. Mitchell (2000), The Quaternary glacial history of the Zaskar Range, north-west Indian Himalaya, *Quat. Int.*, *65–66*, 81–99, doi:10.1016/S1040-6182(99)00038-5.
- Tella, S., D. Paul, R. G. Berman, W. J. Davis, T. D. Peterson, S. J. Pehrsson, and J. A. Kerswill (2007), Bedrock geology compilation and regional synthesis of parts of Hearne and Rae domains, western Churchill Province, Nunavut-Manitoba, Geol. Surv. of Can., Ottawa, Ont., Canada, doi:10.4095/224573.
- Tewari, V. C., and J. Seckbach (2011), *Stromatolites: Interaction of Microbes with Sediments*, 751 pp., Springer, Dordrecht, Netherlands, doi:10.1007/978-94-007-0397-1.
- Thakur, V. C. (1998), Structure of the Chamba nappe and position of the main central thrust in Kashmir Himalaya, *J. Asian Earth Sci.*, *16*(2–3), 269–282, doi:10.1016/S0743-9547(98)00011-7.
- Tingey, R. J. (1991a), Schematic geological map of Antarctica, scale 1:10,000,000, Bur. of Miner. Resour., Geol. and Geophys., Canberra, A. C. T., Australia.
- Tingey, R. J. (1991b), Commentary on schematic geological map of Antarctica scale 1:10,000,000, Bur. of Miner. Resour., Geol. and Geophys., Canberra, A. C. T., Australia.
- Tobgay, T., N. McQuarrie, and S. P. Long (2009), Metamorphic grade of Paro Formation, western Bhutan and its implications, *Eos Trans. AGU*, *90*(52), Fall Meet. Suppl., Abstract T43C-2126.
- Tófaló, O. R., and H. J. M. Morrás (2009), Evidencias paleoclimáticas en duricostras, paleosuelos y sedimentitas silicoclásticas del Cenozoico de Uruguay, *Asoc. Geol. Argent. Rev.*, *65*(4), 674–686.
- Tófaló, O. R., and P. J. Pazos (2010), Paleoclimatic implications (Late Cretaceous-Paleogene) from micromorphology of calcretes, palustrine limestones and silcretes, southern Paraná Basin, Uruguay, *J. South Am. Earth Sci.*, *29*(3), 665–675, doi:10.1016/j.jsames.2009.09.002.
- Tosdal, R. M., E. Farrar, and A. H. Clark (1981), K-Ar geochronology of the Late Cenozoic volcanic rocks of the Cordillera Occidental, southernmost Peru, *J. Volcanol. Geotherm. Res.*, *10*(1–3), 157–173, doi:10.1016/0377-0273(81)90060-3.
- Traynor, J. J., and C. Sladen (1995), Tectonic and stratigraphic evolution of the Mongolian People's Republic and its influence on hydrocarbon geology and potential, *Mar. Pet. Geol.*, *12*(1), 35–52, doi:10.1016/0264-8172(95)90386-X.
- Turnit, P., A. Voges, and H. Wittekindt (2003), Geologische Karte der Bundesrepublik Deutschland, scale 1:1,000,000, Bundesanst. für Geowiss. und Rohstoffe, Hannover, Germany.
- Tsuchi, R. (1990), Neogene events in Japan and the Pacific, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, *77*(3–4), 355–365, doi:10.1016/0031-0182(90)90186-B.
- Tunbridge, I. P. (1981), Sandy high-energy flood sedimentation—Some criteria for recognition, with an example from the Devonian of S.W. England, *Sediment. Geol.*, *28*(2), 79–95, doi:10.1016/0037-0738(81)90058-0.
- Turner, C. C. (1978), Shortland Islands, Solomon Islands, 1:100,000 geological map, Solomon Islands Geol. Surv., Honiara.
- Ubilla, M., D. Perea, C. Goso Aguilar, and N. Lorenzo (2004), Late Pleistocene vertebrates from northern Uruguay: Tools for biostratigraphic, climatic and environmental reconstruction, *Quat. Int.*, *114*(1), 129–142, doi:10.1016/S1040-6182(03)00048-X.
- U.S. Geological Survey (2008), Geochemistry of rock samples from the National Geochemical Database, *U.S. Geol. Surv. Open File Rep.*, 97-0492.
- Valdiya, A. G., and K. S. Jhingran (1973), *Himalayan Geology*, vol. 3, 475 pp., Wadia Inst. of Himalayan Geol., Delhi, India.
- Valdiya, K. S. (1995), Proterozoic sedimentation and Pan-African geodynamic development in the Himalaya, *Precambrian Res.*, *74*(1–2), 35–55, doi:10.1016/0301-9268(95)00004-O.
- Valdiya, K. S., and S. B. Bhatia (1980), *Stratigraphy and Correlations of Lesser Himalayan Formations: Proceedings*, 330 pp., Hindustan Publ. Corp., Delhi, India.
- Verde, M., and S. Martinez (2004), A new ichnogenus for crustacean trace fossils from the Upper Miocene Camacho Formation of Uruguay, *Palaeontology*, *47*, 39–49, doi:10.1111/j.0031-0239.2004.00346.x.
- Veroslavsky, G., and M. Ubilla (2007), A ‘snapshot’ of the evolution of the Uruguay River (Del Plata Basin): The Salto depositional sequence (Pleistocene, Uruguay, South America), *Quat. Sci. Rev.*, *26*(22–24), 2913–2923, doi:10.1016/j.quascirev.2007.02.018.
- Wadia, D. N. (1975), *Geology of India*, 4th ed., 508 pp., McGraw-Hill, New Delhi.
- Walter, R. (1995), *Geologie von Mitteleuropa*, 6th ed., 566 pp., Schweizerbart, Stuttgart, Germany.
- Wandrey, C. J., and B. E. Law (1998), Geologic map of South Asia (geo8ag), Cent. Energy Resour. Team, U.S. Geol. Surv., Denver, Colo.
- Wang, X. D., and T. Sugiyama (2002), Permian coral faunas of the eastern Cimmerian Continent and their biogeographical implications, *J. Asian Earth Sci.*, *20*(6), 589–597, doi:10.1016/S1367-9120(01)00064-5.
- Watts, W. A. (1962), Early Tertiary pollen deposits in Ireland, *Nature*, *193*(4815), 600, doi:10.1038/193600a0.

- Weissert, H. J., and I. Stössel (2010), *Der Ozean im Gebirge: Eine geologische Zeitreise durch die Schweiz*, Hochschulverlag AG an der ETH Zürich, Zurich, Switzerland.
- Wendt, J., B. Kaufmann, Z. Belka, N. Farsan, and A. K. Bavandpur (2005), Devonian/Lower Carboniferous stratigraphy, facies patterns and palaeogeography of Iran—Part II. Northern and central Iran, *Acta Geol. Pol.*, 55(1), 31–97.
- Wessel, P., and W. H. F. Smith (1996), A global, self-consistent, hierarchical, high-resolution shoreline database, *J. Geophys. Res.*, 101(B4), 8741–8743, doi:10.1029/96JB00104.
- Wheeler, J. O., P. F. Hoffman, K. D. Card, A. Davidson, B. V. Sanford, A. V. Okulitch, and W. R. Roest (1997), Geological map of Canada [CD-ROM], *Map D1860A*, Geol. Surv. of Can., Ottawa, Ont.
- Whitaker, A. J., D. C. Champion, I. P. Sweet, P. Kilgour, and D. P. Connolly (2007), Surface geology of Australia 1:1,000,000 scale, Queensland—2nd edition [digital dataset], Geosci. Aust., Canberra, A. C. T., Australia.
- Whitaker, A. J., H. D. Glanville, P. M. English, A. J. Stewart, A. J. Retter, D. P. Connolly, G. A. Stewart, and C. L. Fisher (2008), Surface geology of Australia 1:1,000,000 scale, South Australia—1st edition [digital dataset], Geosci. Aust., Canberra, A. C. T., Australia.
- Wieczorek, G. F., W. L. Newell, P. G. Chirico, G. S. Gohn, R. Nardini, and T. Putbrese (1998), Preliminary digital map database for Honduras, U.S. Geol. Surv., Reston, Va.
- Williams, M., and J. S. Floyd (2000), Mid-Caradoc (Ordovician) ostracodes from the Craighead Limestone Formation, Girvan district, SW Scotland, *Scott. J. Geol.*, 36, 51–60, doi:10.1144/sjg36010051.
- Williams, P. R., C. R. Johnston, R. A. Almond, and W. H. Simamora (1988), Late Cretaceous to Early Tertiary structural elements of west Kalimantan, *Tectonophysics*, 148(3–4), 279–297, doi:10.1016/0040-1951(88)90135-7.
- Wilson, J. J. (1963), Cretaceous stratigraphy of central Andes of Peru, *AAPG Bull.*, 47(1), 1–34.
- Winkler, W., D. Villagomez, R. Spikings, P. Abegglen, S. Tobler, and A. Eguez (2005), The Chota basin and its significance for the inception and tectonic setting of the inter-Andean depression in Ecuador, *J. South Am. Earth Sci.*, 19(1), 5–19, doi:10.1016/j.jsames.2004.06.006.
- Wright, D. T., and I. Knight (1995), A revised chronostratigraphy for the lower Durness Group, *Scott. J. Geol.*, 31(1), 11–22, doi:10.1144/sjg31010011.
- Wright, J. B. (1965), Petrographic sub-provinces in the Tertiary to recent volcanics of Kenya, *Geol. Mag.*, 102(6), 541–557, doi:10.1017/S001675680000025X.
- Wright, T. (1856), On the palaeontological and stratigraphical relations of the so-called “Sands of the Inferior Oolite,” *Q. J. Geol. Soc.*, 12(1–2), 292–325, doi:10.1144/GSL.JGS.1856.012.01-02.37.
- Yemen Geological Survey and Mineral Resources Board (2009), Geology of Yemen, Sana. [Available at http://web.archive.org/web/20100329141304/http://www.ygsmrb.org.ye/geo_of_yemen.htm]
- Yeo, G., C. W. Jefferson, and P. Ramaekers (2002), A preliminary comparison of Manitou Falls Formation stratigraphy in four Athabasca Basin deposystems, 14 pp., Sask. Geol. Surv., Regina, Sask., Canada.
- Yilmaz, Y. (1993), New evidence and model on the evolution of the southeast Anatolian Orogen, *Geol. Soc. Am. Bull.*, 105(2), 251–271, doi:10.1130/0016-7606(1993)105<0251:NEAMOT>2.3.CO;2.
- Yin, E. H. (1985), Geological map of Sabah, 3rd edition, scale 1:500,000, Geol. Surv. of Malaysia, Kuala Lumpur.
- Zapata, A. M., A. F. Sánchez, S. V. Carrasco, A. Cordona, J. H. Galdos, F. Z. Cerrón, and T. Sempere (2005), The lower Carboniferous of the western edge of Gondwana in Peru and Bolivia: Distribution of sedimentary basins and associated magmatism, paper presented at 6th International Symposium on Andean Geodynamics, Univ. de Barcelona, Barcelona, Spain.
- Zárate, M. A. (2003), Loess of southern South America, *Quat. Sci. Rev.*, 22(18–19), 1987–2006, doi:10.1016/S0277-3791(03)00165-3.
- Zhamojda, A. L. (1968), *Geologicheskoe stroenie SSSR*, vol. 1, *Stratigraphiya*, 711 pp., Nedra, Moscow.