

# AI20MTECH14010

Anjana Vasudevan

**Abstract**—This document aims to plot a conic given its equation using matrix approach where,

Download all python codes from

[https://github.com/anjanavasudevan/  
grad\\_schoolwork/tree/master/EE5609/  
Assignment6/Code](https://github.com/anjanavasudevan/grad_schoolwork/tree/master/EE5609/Assignment6/Code)

and latex-tikz codes from

[https://github.com/anjanavasudevan/  
grad\\_schoolwork/tree/master/EE5609/  
Assignment6/Latex](https://github.com/anjanavasudevan/grad_schoolwork/tree/master/EE5609/Assignment6/Latex)

$$\mathbf{V} = \begin{pmatrix} 2 & \frac{3}{2} \\ \frac{3}{2} & -2 \end{pmatrix} \quad (2.0.6)$$

$$\mathbf{u} = \begin{pmatrix} -\frac{7}{2} \\ \frac{1}{2} \end{pmatrix} \quad (2.0.7)$$

$$f = -2 \quad (2.0.8)$$

$$\det(\mathbf{V}) = \begin{vmatrix} 2 & \frac{3}{2} \\ \frac{3}{2} & -2 \end{vmatrix} = -\frac{25}{4} \quad (2.0.9)$$

As  $\det(\mathbf{V}) < 0$ , the given conic represents a hyperbola.

The characteristic equation of  $\mathbf{V}$  is given by the determinant:

$$|\mathbf{V} - \lambda \mathbf{I}| = 0 \quad (2.0.10)$$

$$\begin{vmatrix} 2 - \lambda & \frac{3}{2} \\ \frac{3}{2} & -2 - \lambda \end{vmatrix} = 0 \quad (2.0.11)$$

$$\Rightarrow \lambda^2 - \frac{25}{4} = 0 \quad (2.0.12)$$

## 1 QUESTION

Trace the following central conic:

$$2x^2 + 3xy - 2y^2 - 7x + y - 2 = 0 \quad (1.0.1)$$

The roots of (2.0.12) (the eigenvalues) are:

$$\lambda_1 = \frac{5}{2}, \lambda_2 = -\frac{5}{2} \quad (2.0.13)$$

## 2 ANSWER

The eigenvector  $\mathbf{p}$  is defined as:

$$\mathbf{V}\mathbf{p} = \lambda\mathbf{p} \quad (2.0.14)$$

$$\Rightarrow (\mathbf{V} - \lambda\mathbf{I})\mathbf{p} = 0 \quad (2.0.15)$$

Any second degree equation of the form:

$$ax^2 + 2bxy + cy^2 + 2dx + 2ey + f = 0 \quad (2.0.1)$$

Evaluating (2.0.15) for  $\lambda_1 = \frac{5}{2}$ , we get:

Can be represented in matrix / vector form as:

$$\mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (2.0.2)$$

where,

$$\mathbf{V} = \mathbf{V}^T = \begin{pmatrix} a & b \\ b & c \end{pmatrix} \quad (2.0.3)$$

$$\mathbf{u} = \begin{pmatrix} d & e \end{pmatrix} \quad (2.0.4)$$

Rewriting (1.0.1) in matrix form, we get:

$$\mathbf{x}^T \begin{pmatrix} 2 & \frac{3}{2} \\ \frac{3}{2} & -2 \end{pmatrix} \mathbf{x} + 2 \begin{pmatrix} -\frac{7}{2} & \frac{1}{2} \end{pmatrix} \mathbf{x} - 2 = 0 \quad (2.0.5)$$