

Math Document Template

Pothukuchi Siddhartha

Abstract—This document solves determinant of a matrix using properties of determinants.

Download all python codes from

a

and latex-tikz codes from

a

1 PROBLEM

By using properties of determinants, show that:

$$\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^3 & b^3 & c^3 \end{vmatrix} = (a-b)(b-c)(c-a)(a+b+c) \quad (1.0.1)$$

2 CONSTRUCTION

3 SOLUTION

Using column operations to simplify the equation, we get:

$$\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^3 & b^3 & c^3 \end{vmatrix} \xrightarrow{C_1 \rightarrow C_1 - C_2} \begin{vmatrix} 0 & 1 & 1 \\ a-b & b & c \\ a^3-b^3 & b^3 & c^3 \end{vmatrix} \quad (3.0.1)$$

$$\begin{vmatrix} 0 & 1 & 1 \\ a-b & b & c \\ a^3-b^3 & b^3 & c^3 \end{vmatrix} \xrightarrow{C_2 \rightarrow C_2 - C_3} \begin{vmatrix} 0 & 0 & 1 \\ a-b & b-c & c \\ a^3-b^3 & b^3-c^3 & c^3 \end{vmatrix} \quad (3.0.2)$$

Taking out common factors $a-b$ and $b-c$ from C_1 and C_2 respectively, we get:

$$(a-b)(b-c) \begin{vmatrix} 0 & 0 & 1 \\ 1 & 1 & c \\ a^2+ab+b^2 & b^2+bc+c^2 & c^3 \end{vmatrix} \quad (3.0.3)$$

Further column operations give:

$$\xrightarrow{C_1 \rightarrow C_1 - C_2} (a-b)(b-c) \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & c \\ a^2+ab-bc-c^2 & b^2+bc+c^2 & c^3 \end{vmatrix} \quad (3.0.4)$$

$$\Rightarrow (a-b)(b-c) \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & c \\ (a^2-c^2)+b(a-c) & b^2+bc+c^2 & c^3 \end{vmatrix} \quad (3.0.5)$$

$$\Rightarrow (a-b)(b-c) \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & c \\ (a-c)(a+c)+b(a-c) & b^2+bc+c^2 & c^3 \end{vmatrix} \quad (3.0.6)$$

$$(3.0.7)$$

Taking $a-c$ from C_1 , we get

$$\Rightarrow (a-b)(b-c)(a-c) \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & c \\ a+b+c & b^2+bc+c^2 & c^3 \end{vmatrix} \quad (3.0.8)$$

$$= (a-b)(b-c)(a-c)(-1)(a+b+c) \quad (3.0.9)$$

$$= (a-b)(b-c)(c-a)(a+b+c) \quad (3.0.10)$$