

ASSIGNMENT 5

Anjana V
AI20MTECH14010

Abstract—This document aims to plot equation of circle with given points using matrices

Download all python codes from

[https://github.com/anjanavasudevan/
grad_schoolwork/tree/master/EE5609/
Assignment5/Code](https://github.com/anjanavasudevan/grad_schoolwork/tree/master/EE5609/Assignment5/Code)

and latex-tikz codes from

[https://github.com/anjanavasudevan/
grad_schoolwork/tree/master/EE5609/
Assignment5/Latex](https://github.com/anjanavasudevan/grad_schoolwork/tree/master/EE5609/Assignment5/Latex)

1 QUESTION

Find the equation to the circle that passes through the points:

$$\mathbf{x}_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \mathbf{x}_2 = \begin{pmatrix} 3 \\ -4 \end{pmatrix}, \mathbf{x}_3 = \begin{pmatrix} 5 \\ -6 \end{pmatrix} \quad (1.0.1)$$

2 ANSWER

The equation of circle in vector form is given by:

$$\mathbf{x}^T \mathbf{x} + 2\mathbf{x}^T \mathbf{u} + f = 0 \quad (2.0.1)$$

Using $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3$ in (??),

$$\mathbf{x}_1^T \mathbf{x}_1 + 2\mathbf{x}_1^T \mathbf{u} + f = 0 \quad (2.0.2)$$

$$\mathbf{x}_2^T \mathbf{x}_2 + 2\mathbf{x}_2^T \mathbf{u} + f = 0 \quad (2.0.3)$$

$$\mathbf{x}_3^T \mathbf{x}_3 + 2\mathbf{x}_3^T \mathbf{u} + f = 0 \quad (2.0.4)$$

The above system can be written in matrix form as:

$$\begin{pmatrix} 2\mathbf{x}_1^T & 1 \\ 2\mathbf{x}_2^T & 1 \\ 2\mathbf{x}_3^T & 1 \end{pmatrix} \begin{pmatrix} \mathbf{u} \\ f \end{pmatrix} = \begin{pmatrix} -\mathbf{x}_1^T \mathbf{x}_1 \\ -\mathbf{x}_2^T \mathbf{x}_2 \\ -\mathbf{x}_3^T \mathbf{x}_3 \end{pmatrix} \quad (2.0.5)$$

Substituting the values for $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3$ in (??),

$$\begin{pmatrix} 2 & 4 & 1 \\ 6 & -8 & 1 \\ 10 & -12 & 1 \end{pmatrix} \begin{pmatrix} \mathbf{u} \\ f \end{pmatrix} = \begin{pmatrix} -5 \\ -25 \\ -61 \end{pmatrix} \quad (2.0.6)$$

Using row echelon form to reduce (??), we get:

$$\begin{array}{l} \xleftrightarrow{R_2 \rightarrow R_2 - 3R_1} \\ \xleftrightarrow{R_3 \rightarrow R_3 - 5R_1} \end{array} \left(\begin{array}{ccc|c} 2 & 4 & 1 & -5 \\ 0 & -20 & -2 & -10 \\ 0 & -32 & -4 & -36 \end{array} \right) \quad (2.0.7)$$

$$\begin{array}{l} \xleftrightarrow{R_2 \rightarrow -\frac{1}{2}R_2} \\ \xleftrightarrow{R_3 \rightarrow -\frac{1}{4}R_3} \end{array} \left(\begin{array}{ccc|c} 2 & 4 & 1 & -5 \\ 0 & 10 & 1 & 5 \\ 0 & 8 & 1 & 9 \end{array} \right) \quad (2.0.8)$$

$$\xleftrightarrow{R_3 \rightarrow 5R_3 - 4R_2} \left(\begin{array}{ccc|c} 2 & 4 & 1 & -5 \\ 0 & 10 & 1 & 5 \\ 0 & 0 & 1 & 25 \end{array} \right) \quad (2.0.9)$$

$$\begin{array}{l} \xleftrightarrow{R_1 \rightarrow \frac{1}{2}R_1} \\ \xleftrightarrow{R_2 \rightarrow \frac{1}{10}R_2} \end{array} \left(\begin{array}{ccc|c} 1 & 2 & \frac{1}{2} & -\frac{5}{2} \\ 0 & 1 & \frac{1}{10} & \frac{2}{5} \\ 0 & 0 & 1 & 25 \end{array} \right) \quad (2.0.10)$$

Back solving the system using (??) and (??), we get:

$$\begin{pmatrix} \mathbf{u} \\ f \end{pmatrix} = \begin{pmatrix} -11 \\ -2 \\ 25 \end{pmatrix} \quad (2.0.11)$$

$$\Rightarrow \mathbf{u} = \begin{pmatrix} -11 \\ -2 \end{pmatrix}, f = 25 \quad (2.0.12)$$

The equation of the circle that passes through the points $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3$ is given by:

$$\mathbf{x}^T \mathbf{x} + 2 \begin{pmatrix} -11 \\ -2 \end{pmatrix}^T \mathbf{x} + f = 0 \quad (2.0.13)$$

$$\Rightarrow x^2 + y^2 - 22x - 4y + 25 = 0 \quad (2.0.14)$$

The plot of the circle is given below:




Fig. 1. Circle centered at $(11, 2)$ with radius 10.