

Contents

1	Abstract	1
2	Introduction	1
3	Physical theory	1

1 Abstract

2 Introduction

3 Physical theory

In this project, a solar system will be studied. By solar system, I mean a system where only gravitational forces effect the bodies, and where there is a large mass fixed in origo¹. Newtons gravitational law states that the gravitational force on a body with mass m from another body with mass M and relative position \vec{r} is given by

$$\vec{F}_G = -\frac{GmM}{\|\vec{r}\|^2}\vec{r}$$

where G is the gravitational constant, $6.67 \cdot 10^{-11} \text{ N m}^2/\text{s}^2$. The direction of the force is given by the fact that gravity is an attractive force. If one of the objects is the sun, the mass is denoted by M_\odot . With the sun placed in origo, r is simply the norm of the position vector of the planet with mass m .

If there are n planets in the solar system, in addition to the sun, the sum of the forces on planet i with mass m_i is

$$\sum \vec{F}_i = \sum_{\substack{j=0 \\ j \neq i}}^n \frac{Gm_i m_j}{\|\vec{r}_i - \vec{r}_j\|^3} (\vec{r}_i - \vec{r}_j)$$

with $m_0 = M_\odot$ and $\vec{r}_0 = \vec{0}$. From Newton's second law, we know that $\sum \vec{F}_i = m_i \vec{a}_i$, so the acceleration of planet i is given by

$$\vec{a}_i = \sum_{\substack{j=0 \\ j \neq i}}^n \frac{Gm_j}{\|\vec{r}_i - \vec{r}_j\|^3} (\vec{r}_i - \vec{r}_j) \quad (1)$$

¹This is a reasonable approximation, as the mass of the sun is much larger than the masses of the planets.

As the sun has been fixed to origo, \vec{a}_0 is set to $\vec{0}$.