

# FYS3150 Project 4

Anders Johansson

November 11, 2016

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Theory</b>	<b>1</b>
<b>3</b>	<b>Results</b>	<b>1</b>
3.1	Example: Analytical expression for the $2 \times 2$ case . . . . .	1
3.2	Comparison with numerical results . . . . .	5

## Abstract

## 1 Introduction

## 2 Theory

## 3 Results

### 3.1 Example: Analytical expression for the $2 \times 2$ case

For a  $2 \times 2$  lattice, the thermodynamic quantities can be found analytically without too much work. To find the partition function, we need to write out all possible microstates and calculate their energies. Using the periodic boundary conditions, we get

Microstate	Energy	Magnetization
$\begin{array}{cccc} & \downarrow & \downarrow & \\ \downarrow & \downarrow & \downarrow & \downarrow \\ \downarrow & \downarrow & \downarrow & \downarrow \\ & \downarrow & \downarrow & \end{array}$	$E = -8J$	$M = -4$
$\begin{array}{cccc} & \downarrow & \uparrow & \\ \downarrow & \downarrow & \downarrow & \downarrow \\ \uparrow & \downarrow & \uparrow & \downarrow \\ & \downarrow & \downarrow & \end{array}$	$E = 0$	$M = -2$
$\begin{array}{cccc} & \uparrow & \downarrow & \\ \downarrow & \downarrow & \downarrow & \downarrow \\ \downarrow & \uparrow & \downarrow & \uparrow \\ & \downarrow & \downarrow & \end{array}$	$E = 0$	$M = -2$
$\begin{array}{cccc} & \uparrow & \uparrow & \\ \downarrow & \downarrow & \downarrow & \downarrow \\ \uparrow & \uparrow & \uparrow & \uparrow \\ & \downarrow & \downarrow & \end{array}$	$E = 0$	$M = 0$
$\begin{array}{cccc} & \downarrow & \downarrow & \\ \uparrow & \downarrow & \uparrow & \downarrow \\ \downarrow & \downarrow & \downarrow & \downarrow \\ & \downarrow & \uparrow & \end{array}$	$E = 0$	$M = -2$
$\begin{array}{cccc} & \downarrow & \uparrow & \\ \uparrow & \downarrow & \uparrow & \downarrow \\ \uparrow & \downarrow & \uparrow & \downarrow \\ & \downarrow & \uparrow & \end{array}$	$E = 0$	$M = 0$
$\begin{array}{cccc} & \uparrow & \downarrow & \\ \uparrow & \downarrow & \uparrow & \downarrow \\ \downarrow & \uparrow & \downarrow & \uparrow \\ & \downarrow & \uparrow & \end{array}$	$E = 8J$	$M = 0$
$\begin{array}{cccc} & \uparrow & \uparrow & \\ \uparrow & \downarrow & \uparrow & \downarrow \\ \uparrow & \uparrow & \uparrow & \uparrow \\ & \downarrow & \uparrow & \end{array}$	$E = 0$	$M = 2$
$\begin{array}{cccc} & \downarrow & \downarrow & \\ \downarrow & \uparrow & \downarrow & \uparrow \\ \downarrow & \downarrow & \downarrow & \downarrow \\ & \uparrow & \downarrow & \end{array}$	$E = 0$	$M = -2$
$\begin{array}{cccc} & \downarrow & \uparrow & \\ \downarrow & \uparrow & \downarrow & \uparrow \\ \uparrow & \downarrow & \uparrow & \downarrow \\ & \uparrow & \downarrow & \end{array}$	$E = 8J$	$M = 0$

Microstate	Energy	Magnetization
$  \begin{array}{cccc}  & \uparrow & \downarrow & \\  \downarrow & \uparrow & \downarrow & \uparrow \\  \downarrow & \uparrow & \downarrow & \uparrow \\  & \uparrow & \downarrow &   \end{array}  $	$E = 0$	$M = 0$
$  \begin{array}{cccc}  & \uparrow & \uparrow & \\  \downarrow & \uparrow & \downarrow & \uparrow \\  \uparrow & \uparrow & \uparrow & \uparrow \\  & \uparrow & \downarrow &   \end{array}  $	$E = 0$	$M = 2$
$  \begin{array}{cccc}  & \downarrow & \downarrow & \\  \uparrow & \uparrow & \uparrow & \uparrow \\  \downarrow & \downarrow & \downarrow & \downarrow \\  & \uparrow & \uparrow &   \end{array}  $	$E = 0$	$M = 0$
$  \begin{array}{cccc}  & \downarrow & \uparrow & \\  \uparrow & \uparrow & \uparrow & \uparrow \\  \uparrow & \downarrow & \uparrow & \downarrow \\  & \uparrow & \uparrow &   \end{array}  $	$E = 0$	$M = 2$
$  \begin{array}{cccc}  & \uparrow & \downarrow & \\  \uparrow & \uparrow & \uparrow & \uparrow \\  \downarrow & \uparrow & \downarrow & \uparrow \\  & \uparrow & \uparrow &   \end{array}  $	$E = 0$	$M = 2$
$  \begin{array}{cccc}  & \uparrow & \uparrow & \\  \uparrow & \uparrow & \uparrow & \uparrow \\  \uparrow & \uparrow & \uparrow & \uparrow \\  & \uparrow & \uparrow &   \end{array}  $	$E = -8J$	$M = 4$

To summarise, we have

Number of $\uparrow$	Multiplicity	Energy	Magnetisation
4	1	$-8J$	4
3	4	0	2
2	2	$8J$	0
2	4	0	0
1	4	0	-2
0	1	$-8J$	-4

Summing over all microstates, we get the partition function

$$Z = \sum_{\text{all microstates}} e^{-\beta E_i} = 2e^{8J\beta} + 2e^{-8J\beta} + 12$$

The expectation value of the energy can then be found from

$$\begin{aligned} \langle E \rangle &= -\frac{\partial \ln(Z)}{\partial \beta} = -\frac{\partial}{\partial \beta} \left( \ln(2e^{8J\beta} + 2e^{-8J\beta} + 12) \right) \\ &= -\frac{1}{2e^{8J\beta} + 2e^{-8J\beta} + 12} \left( 16Je^{8J\beta} - 16Je^{-8J\beta} \right) \\ &= -\frac{8J(e^{8J\beta} - e^{-8J\beta})}{e^{8J\beta} + e^{-8J\beta} + 6} \end{aligned}$$

yielding the heat capacity

$$\begin{aligned} C_V &= \frac{\partial}{\partial T} (\langle E \rangle) = \frac{\partial \beta}{\partial T} \frac{\partial}{\partial \beta} (\langle E \rangle) = -\frac{1}{kT^2} \frac{\partial}{\partial \beta} (\langle E \rangle) \\ &= -\frac{1}{kT^2} \frac{\partial}{\partial \beta} \left( -\frac{8J(e^{8J\beta} - e^{-8J\beta})}{e^{8J\beta} + e^{-8J\beta} + 6} \right) \\ &= \frac{8J}{kT^2} \frac{8J(e^{8J\beta} + e^{-8J\beta})(e^{8J\beta} + e^{-8J\beta} + 6) - (e^{8J\beta} - e^{-8J\beta})8J(e^{8J\beta} - e^{-8J\beta})}{(e^{8J\beta} + e^{-8J\beta} + 6)^2} \\ &= \frac{64J^2}{kT^2} \frac{6(e^{8J\beta} + e^{-8J\beta}) + (e^{8J\beta} + e^{-8J\beta})^2 - (e^{8J\beta} - e^{-8J\beta})^2}{(e^{8J\beta} + e^{-8J\beta} + 6)^2} \\ &= \frac{64J^2}{kT^2} \frac{6(e^{8J\beta} + e^{-8J\beta}) + e^{16J\beta} + 2 + e^{-16J\beta} - (e^{16J\beta} - 2 + e^{-16J\beta})}{(e^{8J\beta} + e^{-8J\beta} + 6)^2} \\ &= \frac{64J^2}{kT^2} \frac{6(e^{8J\beta} + e^{-8J\beta}) + 4}{(e^{8J\beta} + e^{-8J\beta} + 6)^2} \end{aligned}$$

To find the various quantities connected to magnetization, we use the general formula

$$\langle A \rangle = \frac{1}{Z} \sum_{\text{all microstates}} A_i e^{-\beta E_i}$$

Mean magnetic moment(s):

$$\begin{aligned} \langle M \rangle &= \frac{1}{2e^{8J\beta} + 2e^{-8J\beta} + 12} \left( -4e^{8J\beta} + 4 \cdot (-2) + 0 + 0 + 4 \cdot 2 + 4e^{8J\beta} \right) = 0 \\ \langle M^2 \rangle &= \frac{1}{2e^{8J\beta} + 2e^{-8J\beta} + 12} \left( (-4)^2 e^{8J\beta} + 4 \cdot (-2)^2 + 0 + 0 + 4 \cdot 2^2 + 4^2 e^{8J\beta} \right) \\ &= \frac{32(e^{8J\beta} + 1)}{2e^{8J\beta} + 2e^{-8J\beta} + 12} = \frac{16(e^{8J\beta} + 1)}{e^{8J\beta} + e^{-8J\beta} + 6} \\ \langle |M| \rangle &= \frac{1}{2e^{8J\beta} + 2e^{-8J\beta} + 12} \left( |-4|e^{8J\beta} + 4 \cdot |-2| + 0 + 0 + 4 \cdot 2 + 4e^{8J\beta} \right) \end{aligned}$$

$$= \frac{8(e^{8J\beta} + 2)}{2e^{8J\beta} + 2e^{-8J\beta} + 12} = \frac{4(e^{8J\beta} + 2)}{e^{8J\beta} + e^{-8J\beta} + 6}$$

Magnetic susceptibility:

$$\chi = \beta \left( \langle M^2 \rangle - \langle M \rangle^2 \right) = \frac{16\beta(e^{8J\beta} + 1)}{e^{8J\beta} + e^{-8J\beta} + 6}$$

### 3.2 Comparison with numerical results

With  $J = 1$  and  $\beta = 1$ , the above expressions give

$$\begin{aligned} \langle E \rangle &= -7.9839 & C_V &= 0.128329 \\ \langle |M| \rangle &= 3.99464 & \chi &= 15.9732 \end{aligned}$$

Running 100 000 Monte Carlo cycles with the same parameters gives figure 1 on the following page.




Figure 1: Numerical results from 100 000 Monte Carlo cycles.

These results fit fairly well with the numbers calculated above, but we see from the graphs that the values for the heat capacity and magnetic susceptibility have yet to settle down completely. Running with 10000000 Monte Carlo cycles yields

```
9999001 1 -7.98387 63.871 0.027741 15.9731 3.99462 0.128789 15.9723
```

where the numbers are  $i$ ,  $T$ ,  $\langle E \rangle$ ,  $\langle E^2 \rangle$ ,  $\langle M \rangle$ ,  $\langle M^2 \rangle$ ,  $\langle |M| \rangle$ ,  $C_V$  and  $\chi$ . The results fit very well with the analytical results.