

# The intuition

---

- Use magnitude and phase as polar coordinates

# The intuition

---

- Use magnitude and phase as polar coordinates
- Encode both coefficients in a single complex number

# Complex Fourier transform coefficients

---

# Complex Fourier transform coefficients

---

$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left( \int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$d_f = \max_{\varphi \in [0,1)} \left( \int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$c = |c| \cdot e^{i\gamma}$$

# Complex Fourier transform coefficients

---

$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left( \int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$d_f = \max_{\varphi \in [0,1)} \left( \int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$c = |c| \cdot e^{i\gamma}$$

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

# Complex Fourier transform coefficients

---

$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left( \int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$d_f = \max_{\varphi \in [0,1)} \left( \int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$c = |c| \cdot e^{i\gamma}$$

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

# Complex Fourier transform coefficients

---

$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left( \int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$d_f = \max_{\varphi \in [0,1)} \left( \int s(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$


$$c = |c| \cdot e^{i\gamma}$$

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

# Complex Fourier transform coefficients

---

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$






# Complex Fourier transform coefficients

---


$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$



# Complex Fourier transform coefficients

---


$$c_f = \boxed{\frac{d_f}{\sqrt{2}}} \cdot e^{-i2\pi\varphi_f}$$



# Complex Fourier transform coefficients

---


$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$



# Complex Fourier transform coefficients

---


$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$



# Complex Fourier transform coefficients

---


$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$



# Complex Fourier transform coefficients

---

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$



Continuous audio signal

---

$$g(t)$$

## Continuous audio signal

---


$$g(t) \quad g : \mathbb{R} \rightarrow \mathbb{R}$$



# Continuous audio signal

---

$$g(t) \quad g : \mathbb{R} \rightarrow \mathbb{R}$$



## Complex Fourier transform

---

$$\hat{g}(f) = c_f$$

## Complex Fourier transform

---

$$\hat{g}(f) = c_f$$


$$\hat{g} : \mathbb{R} \rightarrow \mathbb{C}$$

# Complex Fourier transform

---

$$\hat{g}(f) = c_f$$

$$\hat{g} : \mathbb{R} \rightarrow \mathbb{C}$$




# Complex Fourier transform

---

$$\hat{g}(f) = c_f$$

$$\hat{g} : \mathbb{R} \rightarrow \mathbb{C}$$




# Complex Fourier transform

---

$$\hat{g}(f) = c_f$$

$$\hat{g} : \mathbb{R} \rightarrow \mathbb{C}$$



# Complex Fourier transform

---

$$d_f = \max_{\varphi \in [0,1)} \left( \int g(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$
$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left( \int g(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

# Complex Fourier transform

---

$$d_f = \max_{\varphi \in [0,1)} \left( \int g(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$
$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left( \int g(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$




# Complex Fourier transform

---

$$d_f = \max_{\varphi \in [0,1)} \left( \int g(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$
$$\varphi_f = \operatorname{argmax}_{\varphi \in [0,1)} \left( \int g(t) \cdot \sin(2\pi \cdot (ft - \varphi)) \cdot dt \right)$$


$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$



# Complex Fourier transform

---

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$



# Complex Fourier transform

---

$$\hat{g}(f) = \int \boxed{g(t)} \cdot e^{-i2\pi ft} dt$$

# Complex Fourier transform

---

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$

# Complex Fourier transform

---

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$

# Complex Fourier transform

---

$$e^{i\gamma} = \cos(\gamma) + i \sin(\gamma)$$

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$

# Complex Fourier transform

---

$$e^{i\gamma} = \cos(\gamma) + i \sin(\gamma)$$

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt = \int g(t) \cdot \cos(-2\pi ft) dt + i \int g(t) \cdot \sin(-2\pi ft) dt$$

# Complex Fourier transform

---

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt = \overset{\text{Real part}}{\int g(t) \cdot \cos(-2\pi ft) dt} + \overset{\text{Imaginary part}}{i \int g(t) \cdot \sin(-2\pi ft) dt}$$



## Magnitude Fourier transform

---

$$|\hat{g}(f)|$$

## Magnitude and phase

---

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

## Magnitude and phase

---

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

$$d_f = \sqrt{2} \cdot |\hat{g}(f)|$$

## Magnitude and phase

---

$$c_f = \boxed{\frac{d_f}{\sqrt{2}}} \cdot e^{-i2\pi\varphi_f}$$

$$d_f = \sqrt{2} \cdot \boxed{|\hat{g}(f)|}$$

## Magnitude and phase

---

$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

$$d_f = \sqrt{2} \cdot |\hat{g}(f)|$$

$$\varphi_f = -\frac{\gamma f}{2\pi}$$

## Magnitude and phase

---


$$c_f = \frac{d_f}{\sqrt{2}} \cdot e^{-i2\pi\varphi_f}$$

$$d_f = \sqrt{2} \cdot |\hat{g}(f)|$$


$$\varphi_f = -\frac{\gamma_f}{2\pi}$$

# Inverse Fourier transform

---



IFT  
←



## Fourier representation

---

$$g(t) = \int c_f \cdot e^{i2\pi f t} df$$



# Fourier representation

---

Pure tone of frequency  $f$


$$g(t) = \int c_f \cdot e^{i2\pi ft} df$$

# Fourier representation

---

Pure tone of frequency  $f$

$$g(t) = \int c_f \cdot e^{i2\pi ft} df$$




# Fourier representation

---

Weight pure tone with magnitude and add phase

$$g(t) = \int c_f \cdot e^{i2\pi ft} df$$




# Fourier representation

---

Add up all (weighted) sinusoids

$$g(t) = \int c_f \cdot e^{i2\pi f t} df$$



## A Fourier roundtrip

---

$$\hat{g}(f) = \int g(t) \cdot e^{-i2\pi ft} dt$$

$$g(t) = \int c_f \cdot e^{i2\pi ft} df$$



