



# Simulation of XXZ Spin Models using Sideband Transitions in Trapped Bosonic Gases

**Anjun Chu (JILA)**

Johannes Will (Leibniz University Hannover)

Jan Arlt (Aarhus University)

Carsten Klempt (Leibniz University Hannover)

Ana Maria Rey (JILA)


DAMOP, June 4<sup>th</sup>, 2020



# Simulation of long-range spin models




- Trapped ions Nature 551, 601 (2017)




K. Hudek & E. Edwards/Univ. of Maryland/IonQ, Inc./JQI

- Rydberg atoms Nature 551, 579 (2017)




- Cavity QED Science 361, 259 (2018)



- Polar molecules & magnetic atoms Science 357, 1002 (2017)

Science 357, 1002 (2017)



# Spin models in trapped atomic gases



## Previous Literature:


- Emulation of long-range spin model via purely contact interaction using carrier transition
- Many-body gap protection in isotropic Heisenberg models

## This Research:

- The role of motional sidebands: adding anisotropy to Heisenberg models
- Applications in quantum simulation and metrology since coherence time is not affected by the sideband protocol


### • Bosonic system

Phys. Rev. Lett. 106, 240801 (2011)




### • Fermionic system

Sci. Adv. 5, eaax1568 (2019)




# Experimental setup

- $^{87}\text{Rb}$  atomic gases  
 $|\downarrow\rangle \equiv |F = 1, m_F = 0\rangle$   
 $|\uparrow\rangle \equiv |F = 2, m_F = 0\rangle$
- Drive motional sidebands via Raman beams
- Gas temperature:  
 $375(25)\text{nK}$
- Trapping frequency:  
 $143\text{Hz}, 21.5\text{Hz}, 171\text{Hz}$
- Atom density:  
 $0.46 \text{ to } 4.8 \times 10^{12}\text{cm}^{-3}$



Collisionless regime: trapping frequency  $\gg$  interaction

# Harmonic trap as mode-space lattice



- Blue sideband transition

$$|\uparrow\uparrow_i\rangle = |\uparrow; n_i^X, n_i^Y, n_i^Z + 1\rangle$$

$$|\downarrow\downarrow_i\rangle = |\downarrow; n_i^X, n_i^Y, n_i^Z\rangle$$

- Similar definitions can apply to carrier and red sideband transitions

$$H_{\text{int}} = \sum_{ij} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j + \sum_{ij} \chi_{ij} S_i^z S_j^z + \sum_i B_i S_i^z$$

# Long-range XXZ models in mode-space lattice

$$H_{\text{int}} = \sum_{ij} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j + \sum_{ij} \chi_{ij} S_i^z S_j^z + \sum_i B_i S_i^z$$

$$J_{ij} = \frac{4\pi\hbar^2 a_{\uparrow\downarrow} V_{ij}^{\text{ex}}}{m} \quad \chi_{ij} = \frac{4\pi\hbar^2 (V_{ij}^{\uparrow\uparrow} a_{\uparrow\uparrow} + V_{ij}^{\downarrow\downarrow} a_{\downarrow\downarrow} - V_{ij}^{\uparrow\downarrow} a_{\uparrow\downarrow} - V_{ij}^{\text{ex}} a_{\uparrow\downarrow})}{m}$$

$$V_{ij}^{\text{ex}} = \int d^3\mathbf{R} \phi_i^{\uparrow}(\mathbf{R}) \phi_i^{\downarrow}(\mathbf{R}) \phi_j^{\uparrow}(\mathbf{R}) \phi_j^{\downarrow}(\mathbf{R}) \quad V_{ij}^{\sigma\sigma'} = \int d^3\mathbf{R} [\phi_i^{\sigma}(\mathbf{R})]^2 [\phi_j^{\sigma'}(\mathbf{R})]^2$$

- Carrier transition  $\longrightarrow \phi_i^{\uparrow}(\mathbf{R}) = \phi_i^{\downarrow}(\mathbf{R}) \longrightarrow$  XXX models
- Sideband transition  $\longrightarrow \phi_i^{\uparrow}(\mathbf{R}) \neq \phi_i^{\downarrow}(\mathbf{R}) \longrightarrow$  XXZ models


# Dynamical phase transition (DPT) in LMG model

$$H_{\text{LMG}} = \chi S^z S^z + \Omega S^x - \tilde{\delta} S^z$$

- Initialization: all spins in  $|\downarrow\rangle$  state
- Order parameter: Long-time average of excitation fraction




$$\overline{N_{\uparrow}}/N = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T N_{\uparrow}(t)/N$$

- Phase diagram: Detuning and interaction vary with a fixed Rabi frequency





# Experimental probe of DPT with inhomogeneity





# Theoretical prediction of spin squeezing

- Ramsey spin squeezing parameter


$$\xi^2 = \min_{\theta} N(\Delta S_{\theta}^{\perp})^2 / |\langle \mathbf{S} \rangle|^2$$

- Discrete truncated Wigner approximation (DTWA)

Phys. Rev. X 5, 011022 (2015)


- 6dB squeezing for N=1000 atoms, limited by the inhomogeneities in Ising couplings
- This sideband protocol can apply to a wide range of systems

(a)




$$H_{\text{int}} = \sum_{ij} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j + \sum_{ij} \chi_{ij} S_i^z S_j^z + \sum_i B_i S_i^z$$

(b)




# Conclusion


- Simulate long-range XXZ model via motional sidebands



- Potential of motional sidebands in spin squeezing generation



- Experimental demonstration via a probe of DPT



Thank you for your attention!

Anjun Chu, Johannes Will, Jan Arlt, Carsten Klempt, and Ana Maria Rey, arXiv:2004.01282