## Generalized linear mixed models

Ben Bolker

McMaster University, Mathematics & Statistics and Biology

30 June 2015

Definitions

- 1me4: Doug Bates, Martin Mächler, Steve Walker
- Data: Josh Banta, Adrian Stier, Sea McKeon, David Julian, Jada-Simone White
- NSERC (Discovery)

Challenges & open questions

SHARCnet

## Outline

- 1 Examples and definitions
- EstimationOverview
  - Overview
  - Methods
- 3 Inference
- 4 Challenges & open questions

## Outline

- 1 Examples and definitions
- 2 Estimation
  - Overview
  - Methods
- 3 Inference
- 4 Challenges & open questions

Definitions

## (G)LMMs: a statistical modeling framework incorporating:

- combinations of categorical and continuous predictors, and interactions

Challenges & open questions

# (Generalized) linear mixed models


## (G)LMMs: a statistical modeling framework incorporating:

- combinations of categorical and continuous predictors, and interactions
- (some) non-Normal responses (e.g. binomial, Poisson, and extensions)

Definitions


## (G)LMMs: a statistical modeling framework incorporating:

- combinations of categorical and continuous predictors, and interactions
- (some) non-Normal responses
   (e.g. binomial, Poisson, and extensions)
- (some) nonlinearity
   (e.g. logistic, exponential, hyperbolic)
- non-independent (grouped) data





References

#### Number of predation events




# Environmental stress: *Glycera* cell survival (D. Julian unpubl.)





# Coral demography (J.-S. White unpubl.)




## Technical definition

$$Y_i \sim \stackrel{ ext{Conditional}}{ ext{Distr}} (g^{-1}(\eta_i), \hspace{0.5cm} \phi)$$
response  $g^{-1}(\eta_i)$  scale parameter function

## Technical definition

## Technical definition



## What are random effects?

#### A method for . . .

- accounting for among-individual, within-block correlation

#### A method for ....

- accounting for among-individual, within-block correlation
- compromising between
   complete pooling (no among-block variance)
   and fixed effects (large among-block variance)
- handling levels selected at random from a larger population
- sharing information among levels (shrinkage estimation)
- estimating variability among levels
- allowing predictions for unmeasured levels

#### A method for ...

- accounting for among-individual, within-block correlation
- compromising between
   complete pooling (no among-block variance)
   and fixed effects (large among-block variance)
- handling levels selected at random from a larger population
- sharing information among levels (shrinkage estimation)
- estimating variability among levels
- allowing predictions for unmeasured levels

#### A method for . . .

- accounting for among-individual, within-block correlation
- compromising between complete pooling (no among-block variance) and fixed effects (large among-block variance)
- handling levels selected at random from a larger population
- sharing information among levels (shrinkage estimation)
- estimating variability among levels
- allowing predictions for unmeasured levels

## What are random effects?

#### A method for . . .

- accounting for among-individual, within-block correlation
- compromising between complete pooling (no among-block variance) and fixed effects (large among-block variance)
- handling levels selected at random from a larger population
- sharing information among levels (shrinkage estimation)
- estimating variability among levels
- allowing predictions for unmeasured levels


#### A method for . . .

- accounting for among-individual, within-block correlation
- compromising between complete pooling (no among-block variance) and fixed effects (large among-block variance)
- handling levels selected at random from a larger population
- sharing information among levels (shrinkage estimation)
- estimating variability among levels
- allowing predictions for unmeasured levels


- levels of random effects must always be sampled at random
- a complete sample cannot be treated as a random effect
- random effects are always a nuisance variable
- nothing can be said about the predictions of a random effect
- you should always use a random effect no matter how few levels you have

- 2 Estimation
  - Overview
  - Methods

- Best fit is a compromise between two components (consistency of data with fixed effects and conditional modes; consistency of random effect with RE distribution)
- Goodness-of-fit integrates over conditional modes



# Shrinkage: Arabidopsis conditional modes



Definitions

## deterministic: various approximate integrals (Breslow, 2004)


- Penalized quasi-likelihood, Laplace, Gauss-Hermite quadrature, . . . (?);
   best methods needed for large variance, small clusters
- flexibility and speed vs. accuracy

. . .

stochastic (Monte Carlo): frequentist and Bayesian (Booth and Hobert, 1999; Ponciano et al., 2009; Sung, 2007)

usually slower but flexible and accurate

# Estimation: Culcita (McKeon et al., 2012)




## Outline


- 1 Examples and definition
- 2 Estimation
  - N/I-41---1-
- 3 Inference
- 4 Challenges & open questions

### Wald tests

- typical results of summary
- exact for ANOVA, regression: approximation for GLM(M)s
- fast
- approximation is sometimes awful (Hauck-Donner effect)



# 2D profiles for Culcita data




## Likelihood ratio tests

- better than Wald, but still have to two problems:
  - "denominator degrees of freedom" (when estimating scale)
  - for GLMMs, distributions are approximate anyway (Bartlett corrections)
  - Kenward-Roger correction? (Stroup, 2014)
- Profile confidence intervals: expensive/fragile

- fit null model to data
- simulate "data" from null model
- fit null and working model, compute likelihood difference
- repeat to estimate null distribution
- should be OK but ??? not well tested (assumes estimated parameters are "sufficiently" good)


# Parametric bootstrap results (Glycera)



# Bayesian inference

- If we have a good sample from the posterior distribution (Markov chains have converged etc. etc.) we get most of the inferences we want for free by summarizing the marginal posteriors
- post hoc Bayesian can work, but mode at zero causes problems

## Culcita confidence intervals



## Outline

- 1 Examples and definition
- 2 Estimation
  - Mathada
  - Methods
- 3 Inference
- 4 Challenges & open questions

# On beyond R

- Julia: MixedModels package
- SAS: PROC MIXED, NLMIXED
- AS-REML
- Stata (GLLAMM, xtmelogit)
- AD Model Builder; Template Model Builder
- HLM, MLWiN

## Challenges

Definitions

- Small clusters: need AGQ/MCMC
- Small numbers of clusters: need finite-size corrections (KR/PB/MCMC)
- Small data sets: issues with singular fits Barr et al. (2013) vs. Bates et al. (2015)
- Big data: speed!
- Model diagnosis
- Confidence intervals accounting for uncertainty in variances

See also: http://rpubs.com/bbolker/glmmchapter, https: //groups.nceas.ucsb.edu/non-linear-modeling/projects


# Spatial and temporal correlations

- Sometimes blocking takes care of non-independence ...
- but sometimes there is temporal or spatial correlation within blocks
- ...also phylogenetic ... (Ives and Zhu, 2006)
- "G-side" vs. "R-side" effects
- tricky to implement for GLMMs, but new possibilities on the horizon (Rousset and Ferdy, 2014; Rue et al., 2009);
  - https://github.com/stevencarlislewalker/lme4ord

- Complex random effects: regularization, model selection, penalized methods (lasso/fence)
- Flexible correlation and variance structures
- Flexible/nonparametric random effects distributions
- hybrid & improved MCMC methods
- Reliable assessment of out-of-sample performance

URL here

- http: //www.math.mcmaster.ca/ bolker/R/misc/foxchapter
- Bolker (2015)



### References

- Banta, J.A., Stevens, M.H.H., and Pigliucci, M., 2010, Oikos, 119(2):359-369, ISSN 1600-0706. doi:10.1111/j.1600-0706.2009.17726.x.
- Barr, D.J., Levy, R., et al., 2013. Journal of Memory and Language, 68(3):255-278. ISSN 0749-596X. doi:10.1016/i.iml.2012.11.001.
- Bates, D., Kliegl, R., et al., 2015, arXiv:1506.04967 [stat]. ArXiv: 1506.04967.
- Bolker, B.M., 2015. In G.A. Fox, S. Negrete-Yankelevich, and V.J. Sosa, editors, Ecological Statistics: Contemporary theory and application, Oxford University Press, ISBN 978-0-19-967255-4.
- Booth, J.G. and Hobert, J.P., 1999, Journal of the Royal Statistical Society, Series B, 61(1):265-285. doi:10.1111/1467-9868.00176.
- Breslow, N.E., 2004. In D.Y. Lin and P.J. Heagerty, editors, Proceedings of the second Seattle symposium in biostatistics: Analysis of correlated data, pages 1-22. Springer. ISBN 0387208623.
- lves, A.R. and Zhu, J., 2006. Ecological Applications, 16(1):20-32.
- McKeon, C.S., Stier, A., et al., 2012. Oecologia, 169(4):1095-1103. ISSN 0029-8549. doi:10.1007/s00442-012-2275-2.
- Ponciano, J.M., Taper, M.L., et al., 2009, Ecology, 90(2):356-362, ISSN 0012-9658.
- Rousset, F. and Ferdy, J.B., 2014. Ecography, page no-no. ISSN 1600-0587. doi:10.1111/ecog.00566.
- Rue, H., Martino, S., and Chopin, N., 2009. Journal of the Royal Statistical Society, Series B, 71(2):319-392.
- Stroup, W.W., 2014. Agronomy Journal, 106:1-17. doi:10.2134/agronj2013.0342.
- Sung, Y.J., 2007. The Annals of Statistics, 35(3):990-1011. ISSN 0090-5364. doi:10.1214/009053606000001389.