

Eco-evolutionary virulence of pathogens: models and speculations

Ben Bolker, McMaster University

Departments of Mathematics & Statistics and Biology

NCBS

3 July 2015

Outline

1 Overview

- The evolution of host-pathogen theory
- Toy models

2 Transient virulence and emerging diseases

- Overview
- Toy model
- Myxomatosis data

3 Transient virulence and seasonality

- Overview
- Toy model
- WNV data

4 More on theory vs. data

- Tradeoff curves
- Conclusions

Acknowledgements

People Arjun Nanda and Dharmini Shah; Christophe Fraser;
Marm Kilpatrick; Anson Wong

Support NSF IRCEB grant 9977063; QSE³ IGERT; NSERC
Discovery grant

Outline

1 Overview

- The evolution of host-pathogen theory
- Toy models

2 Transient virulence and emerging diseases

- Overview
- Toy model
- Myxomatosis data

3 Transient virulence and seasonality

- Overview
- Toy model
- WNV data

4 More on theory vs. data

- Tradeoff curves
- Conclusions

Host-pathogen evolutionary biology

Why is it interesting?

- Intellectual merit
 - Coevolutionary loops
 - Cryptic effects
 - Eco-evolutionary dynamics (Luo and Koelle, 2013)
 - Cool stories
 - Lots of data (sometimes)
- Broader applications
 - Medical
 - Conservation and management
 - Outreach

Host-pathogen evolutionary biology

Why is it interesting?

- Intellectual merit
 - Coevolutionary loops
 - Cryptic effects
 - Eco-evolutionary dynamics (Luo and Koelle, 2013)
 - Cool stories
 - Lots of data (sometimes)
- Broader applications
 - Medical
 - Conservation and management
 - Outreach

Virulence: definitions

- General public: badness
- Plant biologists: infectivity
- Evolutionists: loss of host fitness
- Theoreticians: *rate* of host mortality
(mortality rate vs. case mortality vs. clearance)

Basic tradeoff theory: assumptions

- Homogeneous, non-evolving hosts
- No superinfection/coinfection
- Horizontal, direct transmission
- Tradeoff between *rate* of transmission and length of infectious period
- Infectious period $\propto 1/\text{clearance}$
(= recovery+*disease-induced mortality*+*natural mortality*)


○○○○●○○
○○○○○

○○○
○○○○○
○○○○○○○○○

○○
○○○○
○○○

○○○○○○○
○○○

Tradeoffs, \mathcal{R}_0 , and r




○○○○○○●○
○○○○○○

○○○○
○○○○○○○
○○○○○○○○○○

○○
○○○○○
○○○○

○○○○○○○○
○○○

Tradeoffs, \mathcal{R}_0 , and r




○○○○○○●
○○○○○○

○○○○
○○○○○○○
○○○○○○○○○○

○○
○○○○○
○○○○

○○○○○○○○
○○○

Tradeoffs, \mathcal{R}_0 , and r




Outline

- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 Transient virulence and emerging diseases
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence and seasonality
 - Overview
 - Toy model
 - WNV data
- 4 More on theory vs. data
 - Tradeoff curves
 - Conclusions


Epidemiological model

- SIR model
- Constant population size (birth=death)
- Ignore recovery
- Rescale: $\mu = 1$, $N = 1$ (time units of host lifespan)




Epidemiological model

- SIR model
- Constant population size (birth=death)
- Ignore recovery
- Rescale: $\mu = 1$, $N = 1$ (time units of host lifespan)




Epidemiological model

- SIR model
- Constant population size (birth=death)
- Ignore recovery
- Rescale: $\mu = 1$, $N = 1$ (time units of host lifespan)



Epidemiological model

- SIR model
- Constant population size (birth=death)
- Ignore recovery
- Rescale: $\mu = 1$, $N = 1$ (time units of host lifespan)



The model (2): evolutionary dynamics

Incorporate *trait dynamics*

Standard quantitative genetics model (Abrams, 2001):

Alternatives:

The model (2): evolutionary dynamics

Incorporate *trait dynamics*


Standard quantitative genetics model (Abrams, 2001):

- Fitness depends on mean trait value ($\bar{\alpha}$)
and ecological context (proportion susceptible)
- Constant additive genetic variance V_g
- Trait evolves toward increased fitness:
rate proportional to $\Delta\text{fitness}/\Delta\text{trait}$


Alternatives:

multi-strain, adaptive dynamics, PDEs, agent-based models ...


Evolutionary dynamics, cont.



Evolutionary dynamics, cont.



Power-law tradeoff curves



Outline

- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 Transient virulence and emerging diseases
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence and seasonality
 - Overview
 - Toy model
 - WNV data
- 4 More on theory vs. data
 - Tradeoff curves
 - Conclusions

(Why) are emerging pathogens more virulent?

What might explain initially high, but rapidly decreasing, virulence of emerging pathogens?

- Pathogens with low virulence go unnoticed
- Hosts less resistant to / tolerant of novel parasites
- High transmission → frequent coinfection → selection for virulence
- Disease-induced drop in population density decreases selection for virulence (Lenski and May, 1994)

(Why) are emerging pathogens more virulent?

What might explain initially high, but rapidly decreasing, virulence of emerging pathogens?

- Pathogens with low virulence go unnoticed
- Hosts less resistant to / tolerant of novel parasites
- High transmission → frequent coinfection → selection for virulence
- Disease-induced drop in population density decreases selection for virulence (Lenski and May, 1994)

(Why) are emerging pathogens more virulent?

What might explain initially high, but rapidly decreasing, virulence of emerging pathogens?

- Pathogens with low virulence go unnoticed
- Hosts less resistant to / tolerant of novel parasites
- High transmission → frequent coinfection → selection for virulence
- Disease-induced drop in population density decreases selection for virulence (Lenski and May, 1994)

(Why) are emerging pathogens more virulent?

What might explain initially high, but rapidly decreasing, virulence of emerging pathogens?

- Pathogens with low virulence go unnoticed
- Hosts less resistant to / tolerant of novel parasites
- High transmission → frequent coinfection → selection for virulence
- Disease-induced drop in population density decreases selection for virulence (Lenski and May, 1994)

Transient emerging virulence

When a parasite previously in eco-evolutionary equilibrium emerges in a new host population (at low density) it will show a transient peak in virulence as it spreads

How big is the peak? Does it matter?

Outline

1 Overview

- The evolution of host-pathogen theory
- Toy models

2 Transient virulence and emerging diseases

- Overview
- **Toy model**
- Myxomatosis data

3 Transient virulence and seasonality

- Overview
- Toy model
- WNV data


4 More on theory vs. data

- Tradeoff curves
- Conclusions


Model parameters

Parameter		Alternative	
c	Transmission scale	\mathcal{R}_0^*	Equilibrium \mathcal{R}_0
γ	Transmission curvature	α^*	Equilibrium virulence
$I(0)$	Initial epidemic size	$1/N_0$	Inverse population size
V_g	Genetic variance		


Example



Response variables



Peak height




Estimates for emerging pathogens

Order of magnitude estimates for some emerging high-virulence pathogens:

Pathogen	R_0^*	α^*	Reference
SARS	3	640	Anderson et al. (2004)
West Nile	1.61–3.24	639	Wonham et al. (2004)
HIV	1.43	6.36	Velasco-Hernandez et al. (2002)
myxomatosis	3	5	Dwyer et al. (1990)

Emerging pathogens: where are we?


$CV_g = 0.5$, $I(0) = 10^{-3}$ (middle panel):



Outline

- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 Transient virulence and emerging diseases
 - Overview
 - Toy model
 - **Myxomatosis data**
- 3 Transient virulence and seasonality
 - Overview
 - Toy model
 - WNV data
- 4 More on theory vs. data
 - Tradeoff curves
 - Conclusions

Myxomatosis tradeoff curve



Estimating evolvability (V_g)

- Key parameter: genetic variance in virulence (evolvability)
- Despite case studies of rapid pathogen evolution:
 - myxomatosis (Dwyer et al., 1990)
 - syphilis (Knell, 2004)
 - serial passage experiments (Ebert, 1998)
 - *Plasmodium chabaudi* (Mackinnon and Read, 1999a)

we rarely have enough information to estimate V_g

- Only (?) for myxomatosis do we know the variation in virulence among circulating strains

Estimating evolvability (V_g)

- Key parameter: genetic variance in virulence (evolvability)
- Despite case studies of rapid pathogen evolution:
 - myxomatosis (Dwyer et al., 1990)
 - syphilis (Knell, 2004)
 - serial passage experiments (Ebert, 1998)
 - *Plasmodium chabaudi* (Mackinnon and Read, 1999a)

we rarely have enough information to estimate V_g

- Only (?) for myxomatosis do we know the variation in virulence among circulating strains


○○○○○○○○
○○○○○○

○○○○
○○○○○○
○○○○●○○○○○


○○
○○○○○
○○○○

○○○○○○○○
○○○


Myxomatosis grades vs. time




Myxomatosis variance vs. time



Myxomatosis virulence dynamics: power-law tradeoff



Myxomatosis virulence dynamics: realistic tradeoff




oooooooo
oooooo

oooo
oooooooo
ooooooooo●o


oo
ooooo
oooo

oooooooo
ooo

Myxo virulence: equilibrium start, power-law tradeoff



Myxo virulence: equilibrium start, realistic tradeoff



Outline


- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 Transient virulence and emerging diseases
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence and seasonality
 - Overview
 - Toy model
 - WNV data
- 4 More on theory vs. data
 - Tradeoff curves
 - Conclusions

Outline



- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 Transient virulence and emerging diseases
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence and seasonality
 - Overview
 - **Toy model**
 - WNV data
- 4 More on theory vs. data
 - Tradeoff curves
 - Conclusions

Toy model



- Basic Ross-MacDonald vector-host model
- Simple vector (mosquito) demography
- No host demography
- Two pathogen strains





Case I: $r_1 > r_2$, equal \mathcal{R}_0



Case II: $\mathcal{R}_{0,1} > \mathcal{R}_{0,2}$, equal r




Case III: $\mathcal{R}_{0,1} > \mathcal{R}_{0,2}$, $r_2 > r_1$




Outline

- 1 Overview
 - The evolution of host-pathogen theory
 - Toy models
- 2 Transient virulence and emerging diseases
 - Overview
 - Toy model
 - Myxomatosis data
- 3 Transient virulence and seasonality
 - Overview
 - Toy model
 - WNV data
- 4 More on theory vs. data
 - Tradeoff curves
 - Conclusions

Titer vs infectiousness



Titer curves (American crows)




oooooooo
ooooooo

oooo
oooooooo
oooooooooooo

oo
ooooo
ooo●

oooooooo
ooo

Transmission vs clearance for WNV



Outline

1 Overview

- The evolution of host-pathogen theory
- Toy models

2 Transient virulence and emerging diseases

- Overview
- Toy model
- Myxomatosis data


3 Transient virulence and seasonality

- Overview
- Toy model
- WNV data

4 More on theory vs. data

- Tradeoff curves
- Conclusions

Malaria (Mackinnon and Read, 1999b; Paul et al., 2004)




oooooooo
oooooo

oooo
oooooooo
oooooooooooo


oo
oooo
oooo

ooo●oooo
ooo


Pasteuria ramosa (Jensen et al., 2006)




HIV (Fraser et al., 2007)



HIV dynamics (Shirreff et al., 2011)




Phage dynamics (Berngruber et al., 2013)



What about space?

- Theory: spatial structure should select for decreased virulence
- Experiment: viscosity decreases *infectivity* in *Plodia* (Boots and Meador, 2007)
- Are we ready for space?



Outline

1 Overview

- The evolution of host-pathogen theory
- Toy models

2 Transient virulence and emerging diseases

- Overview
- Toy model
- Myxomatosis data

3 Transient virulence and seasonality

- Overview
- Toy model
- WNV data

4 More on theory vs. data

- Tradeoff curves
- Conclusions

Crome (1997) on theory

When we regard theories as tight, real entities and devote ourselves to their analysis, we can limit our horizons and, worse, attempt to make the world fit them. A lot of ecological discussion is not about nature, but about theories, generalizations, or models supposed to represent nature ...

References

- Abrams, P.A., 2001. *Ecol Lett*, 4:166–175.
- Alizon, S., Hurford, A., et al., 2009. *J. Evol. Biol.*, 22:245–259.
doi:10.1111/j.1420-9101.2008.01658.x.
- Anderson, R.M., Fraser, C., et al., 2004. *Phil Trans R Soc London B*, 359(1447):1091–1105.
- Berngruber, T.W., Froissart, R., et al., 2013. *PLoS Pathog*, 9(3):e1003209.
doi:10.1371/journal.ppat.1003209.
- Boots, M. and Meador, M., 2007. *Science*, 315(5816):1284–1286.
- Crome, F.H.J., 1997. In W.F. Laurance and J. Richard O. Bierregard, editors, *Tropical Forest Remnants: Ecology, Management and Conservation of Fragmented Communities*, chapter 31, pages 485–501. University of Chicago Press, Chicago.
- Day, T. and Proulx, S.R., 2004. *Amer Nat*, 163(4):E40–E63.
- Dwyer, G., Levin, S., and Buttel, L., 1990. *Ecol Monog*, 60:423–447.
- Ebert, D., 1998. *Science*, 282(5393):1432–1435.
- Fenner, F., Day, M.F., and Woodroffe, G.M., 1956. *J Hyg (London)*, 54(2):284–302.
- Frank, S.A., 1996. *Q Rev Biol*, 71(1):37–78.
- Fraser, C., Hollingsworth, T.D., et al., 2007. *PNAS*, 104:17441–17446.
- Jensen, K.H., Little, T., et al., 2006. *PLoS Biology*, 4(7):e197.
- Knell, R.J., 2004. *Proc R Soc London B*, 271:S174–S176.
- Lenski, R.E. and May, R.M., 1994. *J Theor Biol*, 169:253–265.
- Lipsitch, M. and Moxon, E.R., 1997. *Trends Microbiol*, 5(1):31–37.
- Luo, S. and Koelle, K., 2013. *The American Naturalist*, 181(S1):S58–S75. ISSN 0003-0147.
doi:10.1086/669952.
- Mackinnon, M.J. and Read, A.F., 1999a. *Evolution*, 53(3):689–703.
- , 1999b. *Proc R Soc London B*, 266(1420):741–748.
- Paul, R.E.L., Lafond, T., et al., 2004. *BMC Evol Biol*, 4:30.
- Shirreff, G., Pellis, L., et al., 2011. *PLoS Computational Biology*, 7(10). ISSN 1553-734X.
doi:10.1371/journal.pcbi.1002185. WOS:000297262700019.
- Velasco-Hernandez, J.X., Gershgorin, H.B., and Blower, S.M., 2002. *Lancet*, 2:487–493.
- Wonham, M.J., de Camino-Beck, T., and Lewis, M.A., 2004. *Proc R Soc London B*, 271:501–507.