

Results

Repeated Measures ANOVA

Within Subjects Effects

Cases	Sum of Squares	df	Mean Square	F	p
RM Factor 1	4.147 ^a	3 ^a	1.382 ^a	6.896 ^a	.001 ^a
Residuals	5.412	27	0.2005		

Note. Type III Sum of Squares

^a Mauchly's test of sphericity indicates that the assumption of sphericity is violated ($p < .05$).


Between Subjects Effects

Cases	Sum of Squares	df	Mean Square	F	p
Residuals	3.065	9	0.3406		

Note. Type III Sum of Squares

Descriptives

Descriptives plots



Nonparametrics

Friedman Test

Factor	X^2_F	df	p	Kendall's W
RM Factor 1	17.48	3	< .001	0.5828

Conover Test

Conover's Post Hoc Comparisons - RM Factor 1

		T-Stat	df	W_i	W_j	r_{rb}	p	pBonf	pHolm
DeiT	ViT	2.045	27	23.00	15.00	0.7091	.051	.304	.116
	Swin	0.1278	27	23.00	23.50	0.05455	.899	1.000	.899
	Ensemble	3.963	27	23.00	38.50	-1.000	< .001	.003	.002
ViT	Swin	2.173	27	15.00	23.50	-0.4727	.039	.232	.116
	Ensemble	6.009	27	15.00	38.50	-1.000	< .001	< .001	< .001
Swin	Ensemble	3.835	27	23.50	38.50	-0.9333	< .001	.004	.003

Note. Grouped by subject.

Note. Rank-biserial correlation based on individual signed-rank tests.

Descriptive Statistics

Descriptive Statistics

	DeiT	ViT	Swin	Ensemble
Valid	10	10	10	10
Missing	0	0	0	0
Mode	98.74	98.05	98.54	98.94
Median	98.72	98.10	98.57	98.95
Mean (arithmetic)	98.61	98.14	98.45	99.03
Std. Deviation	0.2693	0.5597	0.7000	0.2571
Range	0.9500	2.090	2.570	0.8600
Minimum	98.00	97.05	96.57	98.57
Maximum	98.95	99.14	99.14	99.43

^a More than one mode exists. For nominal and ordinal data, the first mode is reported. For continuous data, the mode with the highest density estimate is reported but multiple modes may exist. We recommend visualizing the data to check for multimodality.