

Mathematical Foundations of AI (Aug'24-Dec'24)

Problem Sheet

1. Prove or disprove the following.

(a) $\lim_{x \rightarrow 0} \frac{\tan 2x}{\sin x} = 2$. (Hint : $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$.)

(b) $\lim_{x \rightarrow 0} \frac{\sin 5x - \sin 3x}{x} = 2$.

(c) $\lim_{x \rightarrow 0} \frac{1 - \sqrt{1 - x^2}}{x^2} = \frac{1}{2}$.

2. Let $f(x)$ be defined as $f(x) = 2 \cos x$ if $x \leq c$ and $f(x) = ax^2 + b$ if $x > c$. a, b, c are constants. Given b and c , find all values of a for which $f(x)$ is continuous at $x = c$.
3. Consider $f(x) = \sin(x^{-1})$ for $x \neq 0$ and $f(0) = a$. Determine (if any) the value of a such that $f(x) \rightarrow A$ (for some A) as $x \rightarrow 0$. Justify your answer.
4. Consider $f(x) = [x^{-1}]$ for $x \neq 0$. $[t]$ denotes the greatest integer less than or equal to t . For example, $[2.5] = 2$ and $[-3.5] = -4$. Can you define $f(0)$ so that f is continuous at 0.
5. Let f and g be functions defined as $f(x) = 1$ if $|x| \leq 2$ and $f(x) = 0$ if $|x| > 2$, and $g(x) = 2 - x^2$ if $|x| \leq 3$ and $g(x) = 2$ if $|x| > 3$. Determine a formula for computing $h(x) = f(g(x))$. For what values of x is h continuous ?
6. Let $f(x)$ be defined by $f(x) = x^2$ if $x \leq c$ and $f(x) = ax^3 + b$ if $x > c$ where a, b, c are constants. Find values of a and b (in terms of c) such that $f'(c)$ exists.
7. Solve Exercise (6.) for $f(x)$ where $f(x) = |x|^{-1}$ if $|x| > c$ and $f(x) = a + bx^2$ if $|x| \leq c$.
8. A reservoir has the shape of a right-circular cone. The altitude is 20 feet, and the radius is 8 feet. Water is poured into the reservoir at a constant rate of 5 cubic feet per minute. How fast is the water level rising when the depth of the water is 5 feet if (a) the vertex of the cone is up ? and (b) the vertex of the cone is down ?

9. Prove that among all rectangles of a given area, square has the smallest perimeter.
10. A farmer has L feet of fencing to enclose a rectangular pasture adjacent to a long stone wall. What dimensions give the maximum area of the pasture ?
11. Find the rectangle of the largest area that can be inscribed in a semi-circle, the lower base being on the diameter.
12. Given n real numbers x_1, \dots, x_n , prove that the sum $\sum_{i=1}^n (x - a_i)^2$ is the smallest when x is the arithmetic mean of a_1, \dots, a_n .
13. Let $0 < a < b$ be given. We want to approximate an unknown $x \in [a, b]$ by another number $t \in [a, b]$ so that the maximum relative error $M(t) = \max_{x \in [a, b]} \frac{|t-x|}{x}$ is as small as possible. Prove that $M(t)$ is achieved at either $x = a$ or $x = b$, for each t . Prove that the smallest value of $M(t)$ is achieved by the harmonic mean of a and b , that is, for t satisfying $\frac{1}{t} = \frac{1}{2} \left(\frac{1}{a} + \frac{1}{b} \right)$.
14. Determine the Taylor series expansion for $f(x)$ around $x = 0$ for each of the following definitions of $f(x)$.
 - (a) $f(x) = a^x$, $a > 0$, for all x .
 - (b) $f(x) = \frac{1}{2-x}$ for $|x| < 2$.
 - (c) $f(x) = \log \sqrt{\frac{1+x}{1-x}}$ for $|x| < 1$.
15. For each of the following scalar functions, determine the set of points (x, y) or (x, y, z) at which (i) f is defined and (ii) f is continuous.
 - (a) $f(x, y, z) = \log_e (x^2 + y^2 + z^2)$ for $(x, y, z) \neq (0, 0, 0)$; $f(0, 0, 0) = 1$.
 - (b) $f(x, y) = \frac{x+y}{x^2+y^2}$ for $(x, y) \neq (0, 0)$ and $f(0, 0) = 0$.
 - (c) $f(x, y) = \frac{x}{\sqrt{x^2+y^2}}$.
16. For $(x, y) \neq (0, 0)$, define $f(x, y) = \frac{x^3-y^3}{x^2+y^2}$. Is it possible to define $f(0, 0)$ so as to make f continuous at $(0, 0)$.

17. Define $v(r, t) = t^n \cdot e^{-r^2/(4t)}$. Find a value of the constant n such that v satisfies the following equation $\frac{\partial v}{\partial t} = \frac{1}{r^2} \cdot \frac{\partial}{\partial r} \left(r^2 \frac{\partial v}{\partial r} \right)$.
18. Find the gradient vector at each point at which it exists for the scalar function given by the following definitions.
- (a) $f(x, y, z) = \log_e(x^2 + 2y^2 - 3z^2)$.
 - (b) $f(x, y, z) = x^{y^z}$.
19. Find the directional derivatives for each of the following scalar functions for the points and directions given :
- (a) $f(x, y, z) = x^2 + 2y^2 + 3z^2$ at $(1, 1, 0)$ in the direction of $(1, -1, 2)$.
 - (b) $f(x, y, z) = \left(\frac{x}{y}\right)^z$ at $(1, 1, 1)$ in the direction of $(2, 1, -1)$.
20. For each of the following definitions of $f(x, y)$, $x = X(t)$ and $y = Y(t)$, define $F(t) = f(X(t), Y(t))$ and compute $F'(t)$ and $F''(t)$.
- (a) $f(x, y) = x^2 + y^2$, $X(t) = t$ and $Y(t) = t^2$.
 - (b) $f(x, y) = e^{xy} \cos(xy^2)$, $X(t) = \cos t$, $Y(t) = \sin t$.