

Lecture Summary: Row Reduction and Determinants

Source: Lec25.pdf

Key Points

- **Elementary Row Operations:**

- **Type 1: Row Interchange.** Swap two rows, denoted as $R_i \leftrightarrow R_j$.
- **Type 2: Scalar Multiplication.** Multiply a row by a non-zero scalar t , denoted as tR_i .
- **Type 3: Row Addition.** Add a multiple of one row to another, denoted as $R_i \leftarrow R_i + tR_j$.

- **Row Reduction:**

- A systematic method to transform a matrix into **Row Echelon Form (REF)** or **Reduced Row Echelon Form (RREF)**.
- Steps:
 1. Identify the leftmost non-zero column.
 2. Use row operations to create a leading 1 in the top position of this column.
 3. Use row operations to create zeros below this leading 1.
 4. Repeat for the submatrix below the current row.

- **Reduced Row Echelon Form:**

- In addition to REF:
 1. Each leading 1 in a row has zeros above and below it.
 2. Leading 1s move from left to right in subsequent rows.

- **Applications of Row Reduction:**

- Solve systems of linear equations $Ax = b$.
- Compute the determinant of square matrices.

- **Determinants via Row Reduction:**

- Row reduce a square matrix to REF (upper triangular form).
- If the diagonal entries are all zeros, $\det(A) = 0$.
- If non-zero, compute $\det(A)$ as:

$$\det(A) = \prod (\text{Diagonal Entries in REF}) \cdot \text{Product of scaling factors from Type 2 operations.}$$

- Type 1 row operations change the sign of the determinant, while Type 3 row operations do not affect it.

Simplified Explanation

Example: Row Reduction to REF Given:

$$A = \begin{bmatrix} 2 & 4 & 1 \\ 3 & 8 & 7 \\ 5 & 6 & 9 \end{bmatrix}.$$

Steps:

1. Divide R_1 by 2:

$$\begin{bmatrix} 1 & 2 & 0.5 \\ 3 & 8 & 7 \\ 5 & 6 & 9 \end{bmatrix}.$$

2. Subtract $3R_1$ from R_2 and $5R_1$ from R_3 :

$$\begin{bmatrix} 1 & 2 & 0.5 \\ 0 & 2 & 5.5 \\ 0 & -4 & 6.5 \end{bmatrix}.$$

3. Divide R_2 by 2 and eliminate the entry below $R_2(2,2)$ using $R_3 + 2R_2$:

$$\begin{bmatrix} 1 & 2 & 0.5 \\ 0 & 1 & 2.75 \\ 0 & 0 & 8 \end{bmatrix}.$$

4. Divide R_3 by 8:

$$\begin{bmatrix} 1 & 2 & 0.5 \\ 0 & 1 & 2.75 \\ 0 & 0 & 1 \end{bmatrix}.$$

Determinant Calculation

- Diagonal entries: $1, 1, 1$.
- Scaling factors from Type 2 operations: $\frac{1}{2} \cdot 2 \cdot 8 = 8$.
- Final determinant:

$$\det(A) = 1 \cdot 1 \cdot 1 \cdot 8 = 8.$$

Conclusion

In this lecture, we:

- Defined row reduction and the elementary row operations.
- Showed how to transform matrices into REF and RREF systematically.
- Illustrated the computation of determinants using row reduction as an efficient alternative to the definition.

Row reduction is a versatile tool for solving linear systems and analyzing matrix properties. It simplifies computations, especially for large matrices.