

Lecture Summary: Tangent Hyperplanes and Linear Approximations

Source: Week 11 - Lec 03.pdf

Key Points

- **Tangent Hyperplanes:**

- A tangent hyperplane generalizes the concept of a tangent plane (in \mathbb{R}^3) to higher dimensions.
- For a scalar-valued function $f : D \subset \mathbb{R}^n \rightarrow \mathbb{R}$, the tangent hyperplane at $\vec{a} \in \mathbb{R}^n$ is the affine flat that contains all tangent lines at \vec{a} .
- **Equation of the Tangent Hyperplane:**

$$z = f(\vec{a}) + \sum_{i=1}^n \frac{\partial f}{\partial x_i}(\vec{a})(x_i - a_i),$$

or equivalently,

$$z = f(\vec{a}) + \nabla f(\vec{a}) \cdot (\vec{x} - \vec{a}),$$

where ∇f is the gradient of f .

- **Interpretation in \mathbb{R}^3 :**

- For $f(x, y)$, the tangent plane at (a, b) is given by:

$$z = f(a, b) + \frac{\partial f}{\partial x}(a, b)(x - a) + \frac{\partial f}{\partial y}(a, b)(y - b).$$

- Geometrically, all tangent lines at (a, b) lie on this plane.

- **Linear Approximation:**

- The tangent hyperplane provides the best linear approximation to f near \vec{a} .
- Define $L_f(\vec{x})$ as:

$$L_f(\vec{x}) = f(\vec{a}) + \nabla f(\vec{a}) \cdot (\vec{x} - \vec{a}).$$

- This is the linear approximation of f around \vec{a} .

- **Examples:**

- **Example 1:** $f(x, y) = x + y$ at $(1, 1)$:

$$\nabla f = (1, 1), \quad z = 2 + 1(x - 1) + 1(y - 1).$$

Simplify to:

$$z = x + y.$$

- **Example 2:** $f(x, y) = xy$ at $(1, 1)$:

$$\nabla f = (y, x), \quad z = 1 + 1(x - 1) + 1(y - 1).$$

Simplify to:

$$z = x + y - 1.$$

- **Example 3:** $f(x, y, z) = x^2 + y^2 + z^2$ at $(2, 3, -1)$:

$$\nabla f = (2x, 2y, 2z), \quad \nabla f(2, 3, -1) = (4, 6, -2).$$

Tangent hyperplane:

$$z = 14 + 4(x - 2) + 6(y - 3) - 2(z + 1).$$

Simplify to:

$$z = 14 + 4x + 6y - 2z - 32 \implies z = 4x + 6y - 2z - 18.$$

- **Conditions for Existence:**

- The gradient ∇f must exist and be continuous in a neighborhood of \vec{a} .
- If ∇f is discontinuous or undefined, the tangent hyperplane may not exist.

Simplified Explanation

Tangent Hyperplanes: The tangent hyperplane is a generalization of the tangent plane for higher dimensions. It contains all tangent lines at a point and approximates the function locally.

Equation: For $f(x, y)$ at (a, b) :

$$z = f(a, b) + \frac{\partial f}{\partial x}(a, b)(x - a) + \frac{\partial f}{\partial y}(a, b)(y - b).$$

Example: For $f(x, y) = x + y$ at $(1, 1)$:

$$z = x + y.$$

Conclusion

In this lecture, we:

- Defined tangent hyperplanes and their role in multivariable calculus.
- Derived their equations and discussed their geometric significance.
- Demonstrated their use in linear approximations.

Tangent hyperplanes are essential for approximating and understanding the behavior of multivariable functions near a point.