

Assignment-15

Ankur Aditya - EE20RESCH11010

Abstract—This document contains the problem related to Linear Transformations (Hoffman:- Page-111,Q-1a)

Download the latex-file from

<https://github.com/ankuraditya13/EE5609-Assignment15>

1 PROBLEM

Let n be a positive integer and \mathbf{F} a field. let \mathbf{W} be the set of all vectors $(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) \in \mathbf{F}^n$ such that $\mathbf{x}_1 + \mathbf{x}_2 + \mathbf{x}_3 + \dots + \mathbf{x}_n = 0$. Prove that \mathbf{W}^0 consists of all linear functional f of the form

$$f(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = c \sum_{j=1}^n \mathbf{x}_j \quad (1.0.1)$$

2 DEFINITIONS

2.1 Definition 1

If the vector space \mathbf{V} is finite dimensional (say dimension $=n$), the dimension of null-space \mathbf{N}_f by rank nullity theorem is given by,

$$|\mathbf{N}_f| = |\mathbf{V}| - 1 = n - 1 \quad (2.1.1)$$

2.2 Definition 2

If \mathbf{V} is a vector space over the field \mathbf{F} and \mathbf{S} is a subset of \mathbf{V} , the annihilator of \mathbf{S} is the set \mathbf{S}^0 of linear functional f on \mathbf{V} such that $f(\alpha) = 0$ for every α in \mathbf{S} .

3 SOLUTION

Let h be the functional,

$$h(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \mathbf{x}_1 + \mathbf{x}_2 + \dots + \mathbf{x}_n \quad (3.0.1)$$

$$\text{Let, } \mathbf{X} = \begin{pmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \\ \vdots \\ \mathbf{x}_n \end{pmatrix} \quad (3.0.2)$$

$$\Rightarrow h(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = 0 \quad (3.0.3)$$

Then \mathbf{W} is in null-space of h . Hence by definition-1, the dimension of \mathbf{W} is,

$$|\mathbf{W}| = n - 1 \quad (3.0.4)$$

Now let,

$$a_j = \epsilon_1 - \epsilon_{i+1}, \text{ for } i = (1, \dots, n-1) \quad (3.0.5)$$

Hence clearly $\{a_1, a_2, \dots, a_{n-1}\}$ are linearly independent. Hence from (3.0.4) and above statement we can conclude that $\{a_1, a_2, \dots, a_{n-1}\}$ are all in \mathbf{W} so they must form basis for \mathbf{W} . Now, it is given that f is linear functional hence,

$$f(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \sum_{j=1}^n c_j \mathbf{x}_j \quad (3.0.6)$$

$$\Rightarrow f(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \mathbf{C}^T \mathbf{X} \quad (3.0.7)$$

Where,

$$\mathbf{C} = \begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix} \quad (3.0.8)$$

Now $f \in \mathbf{W}^0$ from definition-2 is given as,

$$f(a_1) = f(a_2) = \dots = f(a_n) = 0 \text{ for every } a_j \in \mathbf{W} \quad (3.0.9)$$

$$\Rightarrow c_1 - c_i = 0 \forall i = 2 \dots n \quad (3.0.10)$$

$$\Rightarrow c_i = c_1 = c \forall i \quad (3.0.11)$$

$$\Rightarrow \mathbf{C} = c \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \quad (3.0.12)$$

Hence,

$$f(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = \mathbf{C}^T \mathbf{X} \quad (3.0.13)$$

$$\Rightarrow f(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = c \sum_{j=1}^n \mathbf{x}_j \quad (3.0.14)$$

Hence, Proved