

Assignment-16

Ankur Aditya - EE20RESCH11010

Abstract—This document contains the problem related to Eigenvalue and Eigenvectors (UGC-June-2017 Maths Q-78)

Download the latex-file from

<https://github.com/ankuraditya13/EE5609–Assignment16>

1 PROBLEM

Consider the matrix

$$A(x) = \begin{pmatrix} 1+x^2 & 7 & 11 \\ 3x & 2x & 4 \\ 8x & 17 & 13 \end{pmatrix}; x \in \mathbf{R}. \quad (1.0.1)$$

Then,

- a) $A(x)$ has eigenvalue 0 for some $x \in \mathbf{R}$.
- b) 0 is not an eigenvalue of $A(x)$ for any $x \in \mathbf{R}$.
- c) $A(x)$ has eigenvalue 0 $\forall x \in \mathbf{R}$.
- d) $A(x)$ is invertible $\forall x \in \mathbf{R}$.

2 SOLUTION

Let $\lambda = 0$ be an eigenvalue. Hence,

$$|A - \lambda I| = 0 \quad (2.0.1)$$

$$\Rightarrow |A| = 0 \quad (2.0.2)$$

$$\Rightarrow |A| = \begin{vmatrix} 1+x^2 & 7 & 11 \\ 3x & 2x & 4 \\ 8x & 17 & 13 \end{vmatrix} = 0 \quad (2.0.3)$$

Performing row reduction we get,

$$\begin{vmatrix} 1+x^2 & 7 & 11 \\ 0 & \frac{2x^3-19x}{1+x^2} & \frac{4x^2-33x+4}{1+x^2} \\ 0 & 0 & \frac{26x^3-244x^2+538x-68}{2x^3-19x} \end{vmatrix} = 0 \quad (2.0.4)$$

$$\Rightarrow 26x^3 - 244x^2 + 538x - 68 = 0 \quad (2.0.5)$$

$$\Rightarrow x_1 = 6.01, x_2 = 3.23, x_3 = 0.13 \quad (2.0.6)$$

OPTIONS	Explanation
Option (b)	At the Values of x given by (2.0.6), eigen value $\lambda = 0$. Hence option (b) can't be correct.
Option (c)	If one of the eigenvalue is 0 for $A(x)$ then, $ A(x) = 0 \forall x \in \mathbf{R}$. But from (2.0.6) we have concluded that $ A = 0$ only for, $x_1 = 6.01, x_2 = 3.23, x_3 = 0.13$. Hence, Option (c) is incorrect.
Option (d)	Now for the values of x given by (2.0.6), $ A = 0$. Hence it is not invertible $\forall x \in \mathbf{R}$ Hence Option (d) is incorrect.
Option (a)	Now clearly from above arguments $A(x)$ has eigenvalue 0 for some $x \in \mathbf{R}$ Hence Option (a) is Correct.