

## PICTURES IN DIGESTIVE PATHOLOGY

### Atypical left paraduodenal hernia

M. Socas Macías, J. M. Álamo Martín, J. M. Suárez Grau, G. Suárez Artacho, A. Tejada, J. Martín Cartes, J. M. Hissnard Cadet, M. Bustos, J. D. Tutosaus Gómez and S. Morales Méndez

*Service of General Surgery. University Hospitals "Virgen del Rocío" and "Hospital General". Sevilla, Spain*

#### CASE REPORT

A 47-year-old man came to the hospital because acute abdominal pain and vomits. The patient had suffered from episodes of intestinal obstruction since adolescence, and has been studied by a gastroenterologist who found no abnormalities. These episodes have been solved spontaneously in nearly all occasions. However, he had come to the emergency room for the same reason twice last year. The exploration showed abdominal distension and tympanism in the epigastrium. There were neither laparotomic scars nor hernia defects in the abdominal wall. Abdominal X-rays showed air-fluid levels at the small bowel, and marked distension in the proximal and medium jejunum. Gastrointestinal X-ray series with gastrographin showed no abnormal findings. After 48-72 hours the patient persisted with the same complaints, and we decided to operate. During the operation we found a congenital malformation: an intestinal hernia of jejunum inside a peritoneal sac formed by a peritoneal flap of the left mesentery (Fig. 1), as well as an abnormal implantation of the mesentery above the rectum. We performed a resection of the peritoneal sac (Fig. 2) and the adhesences inside it. The postoperative period was normal.

#### DISCUSSION

Paraduodenal hernias are rare congenital malformations, but they are the most frequent cause of internal hernias (50% of them). They are caused by a failure in the intestinal rotation, and by an asynchronism in the future development of adhesion of the mesocolon. In right paraduodenal hernias, the small bowel is placed totally or partially behind the ascending mesocolon; in left paraduodenal hernias or mesocolic hernias, which are more frequent than the right ones, the small bowel is placed behind the descending mesocolon (1). We present a rare case of a paraduodenal sac formed by a peritoneal




Fig. 1.- A picture of the peritoneal sac forming the left paraduodenal hernia.  
*Imagen del saco peritoneal que formaba la hernia paraduodenal izquierda.*




Fig. 2.- The mesentery after the resection of the peritoneal sac, and its intestinal contents.  
*Raíz del mesenterio tras extirpación del saco peritoneal, y contenido herniario.*

membrane extending from the left side of the mesentery to the descending colon involving a large part of the jejunum, which was not behind the mesocolon as it usually is. Although this is a congenital problem, it is usually diagnosed around the age of 38 years (2). Symptoms vary from recurrent atypical abdominal pain to total intestinal obstruction; however, this condition may be asymptomatic (3). The diagnosis is difficult – CT scans and intestinal follow-throughs are the most sensitive tests, but the condition is usually diagnosed during urgent laparotomy (4). When we operate on a patient with a small bowel obstruction, and the patient has not been operated on before and has no external hernias, we must bear this possibility in mind. Early intervention allows a good postoperative course, as complications due to hernia incarceration are avoided (5).

## REFERENCES

1. Forgacs B, Istvan G, Sugar I, Ondrejka P. Uncommon case of incomplete left paraduodenal hernia. *Magy Seb* 2003; 56 (6): 243-6.
2. Morán JM, Salas J, Sanjuán S, Amaya JL, Rincón P, Serrano A, et al. Paramesocolic hernias: consequences of delayed diagnosis. Report of three new cases. *J Pediatr Surg* 2004; 39 (1): 112-6.
3. Ramachandran P, Sridharan S. Strangulated left paraduodenal hernia in an infant. *Pediatr Surg Int* 2003; 19 (1-2):120-1. Epub 2003 Mar 29.
4. Dritsas ER, Ruiz OR, Kennedy GM, Blackford J, Hasl D. Paraduodenal hernia: a report of two cases. *Am Surg* 2001; 67 (8): 733-6.
5. Huang YC, Chen HL, Hsu WM, Chen SJ, Lai MW, Chang MH. Left paraduodenal hernia presenting as intestinal obstruction: report of one case. *Acta Paediatr Taiwan* 2001; 42 (3): 172-4.

## Hernia paraduodenal izquierda atípica

M. Socas Macías, J. M. Álamo Martín, J. M. Suárez Grau, G. Suárez Artacho, A. Tejada, J. Martín Cartes, J. M. Hissnard Cadet, M. Bustos, J. D. Tutosaus Gómez y S. Morales Méndez

*Servicio de Cirugía General. Hospitales Universitarios Virgen del Rocío. Hospital General. Sevilla*

## CASO CLÍNICO

Varón de 47 años, que acude a urgencias por cuadro de dolor abdominal tipo cólico, acompañado de náuseas y vómitos. Refiere cuadros suboclusivos desde la adolescencia, que han sido estudiados por digestivo sin evidenciar anomalías y resueltos en la mayoría de los casos de forma espontánea, pero que le obligan a acudir a urgencias en dos ocasiones en el último año. A la exploración presenta abdomen distendido y timpanizado en hemiabdomen superior. No presenta cicatrices de laparotomías previas ni defectos herniarios. Radiografías de abdomen, con distensión de asas y niveles hidroaéreos dependientes de yeyuno proximal y medio. Estudio gastroduodenal practicado con gastrografía, sin hallazgos significativos. Tras 48-72 horas de observación no se aprecia mejoría del cuadro, decidiéndose intervención quirúrgica; en la cual se pone de manifiesto una malformación congénita consistente en una hernia interna de asas de yeyuno medio, dentro de un saco peritoneal dependiente de la hoja peritoneal de la cara izquierda del mesenterio (Fig. 1), así como una implantación de la raíz del mesenterio sobre el recto, quedando este en posición retroperitoneal. Se procede a realizar resección del saco peritoneal (Fig. 2) y adhesiolisis de adherencias existentes en su interior. Postoperatorio sin incidencias.

## DISCUSIÓN

Las hernias paraduodenales son hernias congénitas infrecuentes, pero son la causa más frecuente de hernia interna, constituyendo el 50% de las mismas. Es una malformación congénita debida al fallo del proceso de rotación intestinal, y

de asincronismo con el desarrollo de la coalescencia posterior del mesocolon. En la hernia paraduodenal derecha, el intestino delgado se encuentra localizado total o parcialmente por detrás del meso del colon ascendente en la hernia paraduodenal izquierda (HPDI) o mesocólica, más frecuente que la derecha, el intestino delgado se encuentra por detrás del mesocolon descendente (1). Presentamos el caso infrecuente, de un saco paraduodenal formado por una membrana peritoneal localizada entre la cara izquierda del mesenterio y el colon descendente, que englobaba gran parte del yeyuno, no encontrándose este por detrás del mesocolon como suele ocurrir en las HPDI. A pesar de ser congénitas se suelen diagnosticar en torno a los 38 años (2). La sintomatología varía desde dolor abdominal atípico y recurrente, a la obstrucción intestinal completa, pudiendo ser asintomática (3). El diagnóstico es difícil, el TAC y el tránsito intestinal, son los más precisos en el diagnóstico; siendo diagnosticadas en la mayoría de los casos durante una laparotomía exploradora de urgencias (4). Cuando se interviene quirúrgicamente a un paciente con un síndrome de oclusión intestinal de delgado, en el que no existen antecedentes intervenciones previas o hernias externas, hay que pensar en esta posibilidad. Una intervención quirúrgica temprana permite un postoperatorio sin incidencias ya que previene las complicaciones de la estrangulación herniaria (5).