

# **Unit V : Model Evaluation and Selection**

Metrics for Evaluating Classifier  
Performance Model Selection Using  
Statistical Tests of Significance  
Comparing Classifiers Based on  
Cost–Benefit and ROC Curves,  
Confusion Matrix, F-Measure,  
Precision, Recall

# Model Evaluation and Selection

- Evaluation metrics: How can we measure accuracy? Other metrics to consider?
- Use **validation test set** of class-labeled tuples instead of training set when assessing accuracy
- Methods for estimating a classifier's accuracy:
  - Holdout method, random subsampling
  - Cross-validation
  - Bootstrap
- Comparing classifiers:
  - Confidence intervals
  - Cost-benefit analysis and ROC Curves

# Classifier Evaluation Metrics: Confusion Matrix

## Confusion Matrix:

Actual class\Predicted class	$C_1$	$\neg C_1$
$C_1$	True Positives (TP)	False Negatives (FN)
$\neg C_1$	False Positives (FP)	True Negatives (TN)

## Example of Confusion Matrix:

Actual class\Predicted class	buy_computer = yes	buy_computer = no	Total
buy_computer = yes	6954	46	7000
buy_computer = no	412	2588	3000
Total	7366	2634	10000

- Given  $m$  classes, an entry,  $\mathbf{CM}_{i,j}$  in a **confusion matrix** indicates # of tuples in class  $i$  that were labeled by the classifier as class  $j$
- May have extra rows/columns to provide totals

# Classifier Evaluation Metrics: Accuracy, Error Rate, Sensitivity and Specificity

A\P	C	¬C	
C	TP	FN	P
¬C	FP	TN	N
	P'	N'	All

- **Classifier Accuracy**, or recognition rate: percentage of test set tuples that are correctly classified

$$\text{Accuracy} = (TP + TN)/All$$

- **Error rate**:  $1 - \text{accuracy}$ , or  
$$\text{Error rate} = (FP + FN)/All$$

- **Class Imbalance Problem:**

- One class may be *rare*, e.g. fraud, or HIV-positive
- Significant *majority of the negative class* and minority of the positive class
- **Sensitivity**: True Positive recognition rate
  - **Sensitivity** =  $TP/P$
- **Specificity**: True Negative recognition rate
  - **Specificity** =  $TN/N$

# Classifier Evaluation Metrics:

## Precision and Recall, and F-measures

- **Precision:** exactness – what % of tuples that the classifier labeled as positive are actually positive

$$\text{precision} = \frac{TP}{TP + FP}$$

- **Recall:** completeness – what % of positive tuples did the classifier label as positive?

$$\text{recall} = \frac{TP}{TP + FN}$$

- Perfect score is 1.0
- Inverse relationship between precision & recall
- **F measure ( $F_1$  or F-score):** harmonic mean of precision and recall,

- $F_\beta$ : weighted measure of precision and recall  
– assigns  $\beta$  times as much weight to recall as to precision

$$F_\beta = \frac{(1 + \beta^2) \times \text{precision} \times \text{recall}}{\beta^2 \times \text{precision} + \text{recall}}$$

# Classifier Evaluation Metrics: Example

Actual Class\Predicted class	cancer = yes	cancer = no	Total	Recognition(%)
cancer = yes	<b>90</b>	<b>210</b>	300	30.00 ( <i>sensitivity</i> )
cancer = no	<b>140</b>	<b>9560</b>	9700	98.56 ( <i>specificity</i> )
Total	230	9770	10000	96.40 ( <i>accuracy</i> )

–  $Precision = 90/230 = 39.13\%$

$Recall = 90/300 = 30.00\%$

# Evaluating Classifier Accuracy: Holdout & Cross-Validation Methods

- **Holdout method**
  - Given data is randomly partitioned into two independent sets
    - Training set (e.g., 2/3) for model construction
    - Test set (e.g., 1/3) for accuracy estimation
  - Random sampling: a variation of holdout
    - Repeat holdout  $k$  times, accuracy = avg. of the accuracies obtained
- **Cross-validation** ( $k$ -fold, where  $k = 10$  is most popular)
  - Randomly partition the data into  $k$  *mutually exclusive* subsets, each approximately equal size
  - At  $i$ -th iteration, use  $D_i$  as test set and others as training set
  - Leave-one-out:  $k$  folds where  $k = \#$  of tuples, for small sized data
  - **\*Stratified cross-validation\***: folds are stratified so that class dist. in each fold is approx. the same as that in the initial data

# Evaluating Classifier Accuracy: Bootstrap

- **Bootstrap**
  - Works well with small data sets
  - Samples the given training tuples uniformly *with replacement*
    - i.e., each time a tuple is selected, it is equally likely to be selected again and re-added to the training set
- Several bootstrap methods, and a common one is **.632 bootstrap**
  - A data set with  $d$  tuples is sampled  $d$  times, with replacement, resulting in a training set of  $d$  samples. The data tuples that did not make it into the training set end up forming the test set. About 63.2% of the original data end up in the bootstrap, and the remaining 36.8% form the test set (since  $(1 - 1/d)^d \approx e^{-1} = 0.368$ )
  - Repeat the sampling procedure  $k$  times, overall accuracy of the model:

$$Acc(M) = \frac{1}{k} \sum_{i=1}^k (0.632 \times Acc(M_i)_{test\_set} + 0.368 \times Acc(M_i)_{train\_set})$$



# Estimating Confidence Intervals: Classifier Models $M_1$ vs. $M_2$

- Suppose we have 2 classifiers,  $M_1$  and  $M_2$ , which one is better?
- Use 10-fold cross-validation to obtain  $\overline{err}(M_1)$  and  $\overline{err}(M_2)$
- These mean error rates are just *estimates* of error on the true population of *future* data cases
- What if the difference between the 2 error rates is just attributed to *chance*?
  - Use a **test of statistical significance**
  - Obtain **confidence limits** for our error estimates

# Estimating Confidence Intervals: Null Hypothesis

- Perform 10-fold cross-validation
- Assume samples follow a **t distribution** with  $k-1$  **degrees of freedom** (here,  $k=10$ )
- Use **t-test** (or **Student's t-test**)
- **Null Hypothesis:**  $M_1$  &  $M_2$  are the same
- If we can **reject** null hypothesis, then
  - we conclude that the difference between  $M_1$  &  $M_2$  is **statistically significant**
  - Chose model with lower error rate

# Estimating Confidence Intervals: t-test

- If only 1 test set available: **pairwise comparison**

- For  $i^{\text{th}}$  round of 10-fold cross-validation,  $k$  samples are used to obtain  $err(M_1)_i$  and  $err(M_2)_i$

- Average over 10 rounds to get  $\overline{err}(M_1)$  and  $\overline{err}(M_2)$

- **t-test** compute  $t = \frac{\overline{err}(M_1) - \overline{err}(M_2)}{\sqrt{var(M_1 - M_2)/k}}$  **1 degree of freedom** where

$$var(M_1 - M_2) = \frac{1}{k} \sum_{i=1}^k \left[ err(M_1)_i - err(M_2)_i - (\overline{err}(M_1) - \overline{err}(M_2)) \right]^2$$

where


$$var(M_1 - M_2) = \sqrt{\frac{var(M_1)}{k_1} + \frac{var(M_2)}{k_2}}$$

- If two test sets are available: **red**  
 where  $k_1$  &  $k_2$  are # of cross-validation samples used for  $M_1$  &  $M_2$ , resp.  
**t-test**

# Estimating Confidence Intervals:

## Table for t-distribution

TABLE B: t-DISTRIBUTION CRITICAL VALUES



- Symmetric
- **Significance level**, e.g.,  $\text{sig} = 0.05$  or 5% means  $M_1$  &  $M_2$  are *significantly different* for 95% of population
- **Confidence limit**,  $z = \text{sig}/2$

df	Tail probability p											
	.25	.20	.15	.10	.05	.025	.02	.01	.005	.0025	.001	.0005
1	1.000	1.376	1.963	3.078	6.314	12.71	15.89	31.82	63.66	127.3	318.3	636.6
2	.816	1.061	1.386	1.886	2.920	4.303	4.849	6.965	9.925	14.09	22.33	31.60
3	.765	.978	1.250	1.638	2.353	3.182	3.482	4.541	5.841	7.453	10.21	12.92
4	.741	.941	1.190	1.533	2.132	2.776	2.999	3.747	4.604	5.598	7.173	8.610
5	.727	.920	1.156	1.476	2.015	2.571	2.757	3.365	4.032	4.773	5.893	6.869
6	.718	.906	1.134	1.440	1.943	2.447	2.612	3.143	3.707	4.317	5.208	5.959
7	.711	.896	1.119	1.415	1.895	2.365	2.517	2.998	3.499	4.029	4.785	5.408
8	.706	.889	1.108	1.397	1.860	2.306	2.449	2.896	3.355	3.833	4.501	5.041
9	.703	.883	1.100	1.383	1.833	2.262	2.398	2.821	3.250	3.690	4.297	4.781
10	.700	.879	1.093	1.372	1.812	2.228	2.359	2.764	3.169	3.581	4.144	4.587
11	.697	.876	1.088	1.363	1.796	2.201	2.328	2.718	3.106	3.497	4.025	4.437
12	.695	.873	1.083	1.356	1.782	2.179	2.303	2.681	3.055	3.428	3.930	4.318
13	.694	.870	1.079	1.350	1.771	2.160	2.282	2.650	3.012	3.372	3.852	4.221
14	.692	.868	1.076	1.345	1.761	2.145	2.264	2.624	2.977	3.326	3.787	4.140
15	.691	.866	1.074	1.341	1.753	2.131	2.249	2.602	2.947	3.286	3.733	4.073
16	.690	.865	1.071	1.337	1.746	2.120	2.235	2.583	2.921	3.252	3.686	4.015
17	.689	.863	1.069	1.333	1.740	2.110	2.224	2.567	2.898	3.222	3.646	3.965
18	.688	.862	1.067	1.330	1.734	2.101	2.214	2.552	2.878	3.197	3.611	3.922
19	.688	.861	1.066	1.328	1.729	2.093	2.205	2.539	2.861	3.174	3.579	3.883
20	.687	.860	1.064	1.325	1.725	2.086	2.197	2.528	2.845	3.153	3.552	3.850
21	.686	.859	1.063	1.323	1.721	2.080	2.189	2.518	2.831	3.135	3.527	3.819
22	.686	.858	1.061	1.321	1.717	2.074	2.183	2.508	2.819	3.119	3.505	3.792
23	.685	.858	1.060	1.319	1.714	2.069	2.177	2.500	2.807	3.104	3.485	3.768
24	.685	.857	1.059	1.318	1.711	2.064	2.172	2.492	2.797	3.091	3.467	3.745
25	.684	.856	1.058	1.316	1.708	2.060	2.167	2.485	2.787	3.078	3.450	3.725
26	.684	.856	1.058	1.315	1.706	2.056	2.162	2.479	2.779	3.067	3.435	3.707
27	.684	.855	1.057	1.314	1.703	2.052	2.158	2.473	2.771	3.057	3.421	3.690
28	.683	.855	1.056	1.313	1.701	2.048	2.154	2.467	2.763	3.047	3.408	3.674
29	.683	.854	1.055	1.311	1.699	2.045	2.150	2.462	2.756	3.038	3.396	3.659
30	.683	.854	1.055	1.310	1.697	2.042	2.147	2.457	2.750	3.030	3.385	3.646
40	.681	.851	1.050	1.303	1.684	2.021	2.123	2.423	2.704	2.971	3.307	3.551
50	.679	.849	1.047	1.299	1.676	2.009	2.109	2.403	2.678	2.937	3.261	3.496
60	.679	.848	1.045	1.296	1.671	2.000	2.099	2.390	2.660	2.915	3.232	3.460
80	.678	.846	1.043	1.292	1.664	1.990	2.088	2.374	2.639	2.887	3.195	3.416
100	.677	.845	1.042	1.290	1.660	1.984	2.081	2.364	2.626	2.871	3.174	3.390
1000	.675	.842	1.037	1.282	1.646	1.962	2.056	2.330	2.581	2.813	3.098	3.300
∞	.674	.841	1.036	1.282	1.645	1.960	2.054	2.326	2.576	2.807	3.091	3.291
	50%	60%	70%	80%	90%	95%	96%	98%	99%	99.5%	99.8%	99.9%
	Confidence level C											


# Estimating Confidence Intervals:

## Statistical Significance

- Are  $M_1$  &  $M_2$  **significantly different**?
  - Compute  $t$ . Select *significance level* (e.g.  $sig = 5\%$ )
  - Consult table for t-distribution: Find  $t$  *value* corresponding to  $k-1$  *degrees of freedom* (here, 9)
  - t-distribution is symmetric: typically upper % points of distribution shown → look up value for **confidence limit**  $z=sig/2$  (here, 0.025)
  - **If  $t > z$  or  $t < -z$** , then  $t$  value lies in rejection region:
    - **Reject null hypothesis** that mean error rates of  $M_1$  &  $M_2$  are same
    - Conclude: statistically significant difference between  $M_1$  &  $M_2$
  - **Otherwise**, conclude that any difference is **chance**

# Model Selection: ROC Curves

- **ROC** (Receiver Operating Characteristics) curves: for visual comparison of classification models
- Originated from signal detection theory
- Shows the trade-off between the true positive rate and the false positive rate
- The area under the ROC curve is a measure of the accuracy of the model
- Rank the test tuples in decreasing order: the one that is most likely to belong to the positive class appears at the top of the list
- The closer to the diagonal line (i.e., the closer the area is to 0.5), the less accurate is the model



- Vertical axis represents the true positive rate
- Horizontal axis rep. the false positive rate
- The plot also shows a diagonal line
- A model with perfect accuracy will have an area of 1.0

# Issues Affecting Model Selection

- **Accuracy**
  - classifier accuracy: predicting class label
- **Speed**
  - time to construct the model (training time)
  - time to use the model (classification/prediction time)
- **Robustness**: handling noise and missing values
- **Scalability**: efficiency in disk-resident databases
- **Interpretability**
  - understanding and insight provided by the model
- Other measures, e.g., goodness of rules, such as decision tree size or compactness of classification rules