

SURGICAL PATHOLOGY REPORT [SYNTHETIC]

ACCESSION #: UC-2025-68229

DATE OF PROCEDURE: 04/24/2025

DATE OF REPORT: 04/27/2025

REQUESTING PHYSICIAN: Dr. Zachary Perez, Gastroenterology

PATHOLOGIST: Dr. Jacqueline Stephens, Anatomic Pathology

CLINICAL HISTORY:

73 year old female with 2 week history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed moderate inflammation with patchy erosions from rectum to splenic flexure. Clinical suspicion for ulcerative colitis.

SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 5 tan-pink tissue fragments measuring 4 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 2 tan-pink tissue fragments measuring 3 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 6 tan-pink tissue fragments measuring 4 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 5 tan-pink tissue fragments measuring 8 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 5 tan-pink tissue fragments measuring 3 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 2 tan-pink tissue fragments measuring 7 mm in aggregate.

All specimens are entirely submitted in 3 cassette(s).

MICROSCOPIC DESCRIPTION:

A. Rectal mucosa shows moderate active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Occasional apoptotic bodies are present in crypts.

B. Sigmoid colonic mucosa shows severe active chronic inflammation with severe cryptitis, crypt architectural distortion, and Paneth cell metaplasia. The inflammatory process is limited to the mucosa without evidence of granulomas. No evidence of dysplasia is identified.

C. Descending colonic mucosa shows moderate to severe active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Surface epithelium shows reactive changes.

D. Transverse colonic mucosa shows mild to moderate active chronic inflammation with crypt branching, crypt atrophy, and focal crypt abscesses. Reactive epithelial changes are seen adjacent to areas of active inflammation.

E. Ascending colonic mucosa shows mild active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. Lamina propria shows increased plasma cells and lymphocytes.

F. Terminal ileal mucosa shows mild non-specific inflammation without architectural distortion. No evidence of chronic inflammatory bowel disease identified in this section.

DIAGNOSIS:

A. Rectum, biopsy:

- moderate active chronic colitis with crypt architectural distortion and goblet cell depletion
- moderate consistent with ulcerative colitis
- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

B. Sigmoid colon, biopsy:

- severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- severe consistent with ulcerative colitis
- No dysplasia identified

C-E. Descending, transverse, and ascending colon, biopsies:

- moderate to severe active chronic colitis with crypt architectural distortion
- Features consistent with ulcerative colitis
- No dysplasia identified

F. Terminal ileum, biopsy:

- Mild non-specific inflammation
- No evidence of inflammatory bowel disease

COMMENT:

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The presence of diffuse crypt architectural distortion, basal plasmacytosis, and continuous inflammatory pattern strongly supports the diagnosis of ulcerative colitis.

SPECIAL STUDIES:

Immunohistochemical stain for p53 shows no evidence of dysplasia-associated molecular alterations.