

# **SURGICAL PATHOLOGY REPORT [SYNTHETIC]**

**ACCESSION #:** UC-2025-68328

**DATE OF PROCEDURE:** 04/14/2025

**DATE OF REPORT:** 04/16/2025

**REQUESTING PHYSICIAN:** Dr. Joseph Bradley, Gastroenterology

**PATHOLOGIST:** Dr. Jennifer Myers, Anatomic Pathology

## **CLINICAL HISTORY:**

55 year old female with longstanding history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed severe ulceration and spontaneous bleeding from rectum to descending colon. Clinical suspicion for ulcerative colitis.

## **SPECIMEN RECEIVED:**

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

## **GROSS DESCRIPTION:**

- A. Received in formalin labeled "rectum" are 6 tan-pink tissue fragments measuring 8 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 5 tan-pink tissue fragments measuring 7 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 3 tan-pink tissue fragments measuring 4 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 2 tan-pink tissue fragments measuring 5 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 5 tan-pink tissue fragments measuring 7 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 5 tan-pink tissue fragments measuring 4 mm in aggregate.

All specimens are entirely submitted in 4 cassette(s).

## **MICROSCOPIC DESCRIPTION:**

A. Rectal mucosa shows moderate active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Basal plasmacytosis is prominent.

B. Sigmoid colonic mucosa shows moderate active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Reactive epithelial changes are seen adjacent to areas of active inflammation.

C. Descending colonic mucosa shows moderate active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Occasional apoptotic bodies are present in crypts.

D. Transverse colonic mucosa shows mild active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. No evidence of dysplasia is identified.

E. Ascending colonic mucosa shows mild to moderate active chronic inflammation with crypt architectural distortion and crypt abscesses. Basal plasmacytosis is prominent.

F. Terminal ileal mucosa shows normal small intestinal mucosa with preserved villous architecture. No evidence of chronic inflammatory bowel disease identified in this section.

## **DIAGNOSIS:**

### **A. Rectum, biopsy:**

- moderate active chronic colitis with crypt architectural distortion and goblet cell depletion
- moderate consistent with ulcerative colitis
- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

### ***B. Sigmoid colon, biopsy:***

- moderate active chronic colitis with crypt architectural distortion and goblet cell depletion
- moderate consistent with ulcerative colitis
- No dysplasia identified

### ***C-E. Descending, transverse, and ascending colon, biopsies:***

- moderate active chronic colitis with crypt architectural distortion
- Features consistent with ulcerative colitis
- No dysplasia identified

### ***F. Terminal ileum, biopsy:***

- Mild non-specific inflammation
- No evidence of inflammatory bowel disease

## **COMMENT:**

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The histologic findings show classic features of ulcerative colitis with diffuse crypt architectural distortion and diffuse mucosal inflammation.

## **SPECIAL STUDIES:**

Immunohistochemical stain for p53 shows no evidence of dysplasia-associated molecular alterations.