

SURGICAL PATHOLOGY REPORT [SYNTHETIC]

ACCESSION #: UC-2025-46352

DATE OF PROCEDURE: 05/04/2025

DATE OF REPORT: 05/08/2025

REQUESTING PHYSICIAN: Dr. Heather Owens, Gastroenterology

PATHOLOGIST: Dr. Brandy Gonzalez, Anatomic Pathology

CLINICAL HISTORY:

65 year old male with 2 year history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed moderate inflammation with patchy erosions from rectum to splenic flexure. Clinical suspicion for ulcerative colitis.

SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 4 tan-pink tissue fragments measuring 7 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 4 tan-pink tissue fragments measuring 3 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 4 tan-pink tissue fragments measuring 4 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 6 tan-pink tissue fragments measuring 6 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 5 tan-pink tissue fragments measuring 6 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 4 tan-pink tissue fragments measuring 5 mm in aggregate.

All specimens are entirely submitted in 3 cassette(s).

MICROSCOPIC DESCRIPTION:

A. Rectal mucosa shows severe active chronic inflammation with severe cryptitis, crypt architectural distortion, and mucosal ulceration. The inflammatory process is limited to the mucosa without evidence of granulomas. Reactive epithelial changes are seen adjacent to areas of active inflammation.

B. Sigmoid colonic mucosa shows moderate to severe active chronic inflammation with marked crypt architectural distortion, numerous crypt abscesses, and complete goblet cell depletion. The inflammatory process is limited to the mucosa without evidence of granulomas. Mucosal edema and congestion are present.

C. Descending colonic mucosa shows moderate to severe active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. The inflammatory process is limited to the mucosa without evidence of granulomas. Occasional apoptotic bodies are present in crypts.

D. Transverse colonic mucosa shows mild to moderate active chronic inflammation with crypt architectural distortion and crypt abscesses. No evidence of dysplasia is identified.

E. Ascending colonic mucosa shows mild to moderate active chronic inflammation with crypt architectural distortion and crypt abscesses. Reactive epithelial changes are seen adjacent to areas of active inflammation.

F. Terminal ileal mucosa shows essentially normal ileal mucosa with intact villous architecture and no active inflammation. No evidence of chronic inflammatory bowel disease identified in this section.

DIAGNOSIS:

A. Rectum, biopsy:

- severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- severe consistent with ulcerative colitis
- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

B. Sigmoid colon, biopsy:

- moderate to severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- moderate to severe consistent with ulcerative colitis
- No dysplasia identified

C-E. Descending, transverse, and ascending colon, biopsies:

- moderate to severe active chronic colitis with crypt architectural distortion
- Features consistent with ulcerative colitis
- No dysplasia identified

F. Terminal ileum, biopsy:

- Mild non-specific inflammation
- No evidence of inflammatory bowel disease

COMMENT:

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The lack of granulomas, ileal involvement, and transmural inflammation favors ulcerative colitis over Crohn's disease.

SPECIAL STUDIES:

Periodic acid-Schiff (PAS) stain is negative for fungal organisms.