

SURGICAL PATHOLOGY REPORT [SYNTHETIC]

ACCESSION #: UC-2025-17240

DATE OF PROCEDURE: 04/11/2025

DATE OF REPORT: 04/15/2025

REQUESTING PHYSICIAN: Dr. Tyler Scott, Gastroenterology

PATHOLOGIST: Dr. Donald Knapp, Anatomic Pathology

CLINICAL HISTORY:

20 year old female with longstanding history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed moderate inflammation with patchy erosions from rectum to splenic flexure. Clinical suspicion for ulcerative colitis.

SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 4 tan-pink tissue fragments measuring 8 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 3 tan-pink tissue fragments measuring 3 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 4 tan-pink tissue fragments measuring 6 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 2 tan-pink tissue fragments measuring 2 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 2 tan-pink tissue fragments measuring 5 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 6 tan-pink tissue fragments measuring 8 mm in aggregate.

All specimens are entirely submitted in 3 cassette(s).

MICROSCOPIC DESCRIPTION:

A. Rectal mucosa shows moderate to severe active chronic inflammation with crypt architectural distortion, lamina propria plasma cells, and basal plasmacytosis. The inflammatory process is limited to the mucosa without evidence of granulomas. No evidence of dysplasia is identified.

B. Sigmoid colonic mucosa shows mild to moderate active chronic inflammation with severe cryptitis, crypt architectural distortion, and Paneth cell metaplasia. The inflammatory process is limited to the mucosa without evidence of granulomas. No evidence of dysplasia is identified.

C. Descending colonic mucosa shows moderate active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Lamina propria shows increased plasma cells and lymphocytes. Areas of crypt dropout and lamina propria fibrosis are present, suggesting chronicity and possible treatment effect.

D. Transverse colonic mucosa shows mild to moderate active chronic inflammation with marked crypt architectural distortion, numerous crypt abscesses, and complete goblet cell depletion. Lamina propria shows increased plasma cells and lymphocytes.

E. Ascending colonic mucosa shows mild to moderate active chronic inflammation with crypt architectural distortion and crypt abscesses. Lamina propria shows increased plasma cells and lymphocytes.

F. Terminal ileal mucosa shows minimal increase in lamina propria lymphoplasmacytic infiltrates, likely reactive. No evidence of chronic inflammatory bowel disease identified in this section. Terminal ileal mucosa shows mild active inflammation with neutrophilic cryptitis, likely representing backwash ileitis.

DIAGNOSIS:

A. Rectum, biopsy:

- moderate to severe active chronic colitis with crypt architectural distortion and goblet cell depletion

- moderate to severe consistent with ulcerative colitis
- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

B. Sigmoid colon, biopsy:

- mild to moderate active chronic colitis with crypt architectural distortion and goblet cell depletion
- mild to moderate consistent with ulcerative colitis
- No dysplasia identified

C-E. Descending, transverse, and ascending colon, biopsies:

- moderate active chronic colitis with crypt architectural distortion
- Mild active ileitis, consistent with backwash ileitis in the setting of ulcerative colitis
- Features consistent with ulcerative colitis
- No dysplasia identified

F. Terminal ileum, biopsy:

- Mild non-specific inflammation
- Changes consistent with chronicity and treatment effect
- No evidence of inflammatory bowel disease

COMMENT:

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The pattern of inflammation is consistent with ulcerative colitis as evidenced by the continuous mucosal involvement with greatest intensity distally. The mild ileal inflammation in the context of pancolitis is consistent with backwash ileitis, which can be seen in ulcerative colitis and does not necessarily indicate Crohn's disease. Histologic features suggesting chronicity and treatment effect are present. Correlation with treatment history is recommended.

SPECIAL STUDIES:

Immunohistochemical stain for p53 shows no evidence of dysplasia-associated molecular alterations.

This is a synthetic educational pathology report created for AI training purposes. It does not represent a real patient case.