

# SURGICAL PATHOLOGY REPORT [SYNTHETIC]

**ACCESSION #:** UC-2025-38211

**DATE OF PROCEDURE:** 05/09/2025

**DATE OF REPORT:** 05/12/2025

**REQUESTING PHYSICIAN:** Dr. Amber Jackson, Gastroenterology

**PATHOLOGIST:** Dr. Omar Murphy, Anatomic Pathology

## CLINICAL HISTORY:

38 year old male with 10 year history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed diffuse erythema, loss of vascular pattern, and contact bleeding from rectum to hepatic flexure. Clinical suspicion for ulcerative colitis.

## SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

## GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 5 tan-pink tissue fragments measuring 4 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 2 tan-pink tissue fragments measuring 3 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 2 tan-pink tissue fragments measuring 7 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 6 tan-pink tissue fragments measuring 3 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 4 tan-pink tissue fragments measuring 7 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 6 tan-pink tissue fragments measuring 6 mm in aggregate.

All specimens are entirely submitted in 5 cassette(s).

## **MICROSCOPIC DESCRIPTION:**

A. Rectal mucosa shows moderate to severe active chronic inflammation with crypt architectural distortion, lamina propria plasma cells, and basal plasmacytosis. The inflammatory process is limited to the mucosa without evidence of granulomas. No evidence of dysplasia is identified.

B. Sigmoid colonic mucosa shows moderate to severe active chronic inflammation with crypt branching, crypt atrophy, and focal crypt abscesses. The inflammatory process is limited to the mucosa without evidence of granulomas. Marked decrease in goblet cell population.

C. Descending colonic mucosa shows moderate to severe active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Occasional Paneth cell metaplasia is noted.

D. Transverse colonic mucosa shows mild to moderate active chronic inflammation with crypt branching, crypt atrophy, and focal crypt abscesses. Reactive epithelial changes are seen adjacent to areas of active inflammation. Areas of crypt dropout and lamina propria fibrosis are present, suggesting chronicity and possible treatment effect.

E. Ascending colonic mucosa shows mild active chronic inflammation with crypt architectural distortion and crypt abscesses. Mucosal edema and congestion are present.

F. Terminal ileal mucosa shows normal small intestinal mucosa with preserved villous architecture. No evidence of chronic inflammatory bowel disease identified in this section. The inflammatory pattern shows overlapping features of both ulcerative colitis and Crohn's disease.

## **DIAGNOSIS:**

### ***A. Rectum, biopsy:***

- moderate to severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- moderate to severe consistent with ulcerative colitis

- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

### ***B. Sigmoid colon, biopsy:***

- moderate to severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- Features of chronicity consistent with inflammatory bowel disease, with overlapping features of both UC and CD
- moderate to severe consistent with ulcerative colitis
- No dysplasia identified

### ***C-E. Descending, transverse, and ascending colon, biopsies:***

- moderate to severe active chronic colitis with crypt architectural distortion
- Features consistent with ulcerative colitis
- No dysplasia identified

### ***F. Terminal ileum, biopsy:***

- Mild non-specific inflammation
- Changes consistent with chronicity and treatment effect
- No evidence of inflammatory bowel disease

## **COMMENT:**

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The pattern of inflammation is consistent with ulcerative colitis as evidenced by the continuous mucosal involvement with greatest intensity distally. The histologic findings show overlapping features of both ulcerative colitis and Crohn's disease. This pattern may represent an 'indeterminate colitis' and correlation with clinical, endoscopic, and serologic markers is strongly recommended for further classification. Histologic features suggesting chronicity and treatment effect are present. Correlation with treatment history is recommended.

## **SPECIAL STUDIES:**

Acid-fast bacilli (AFB) stain is negative for mycobacterial organisms.

\_This is a synthetic educational pathology report created for AI training purposes. It does not represent a real patient case.\_