

SURGICAL PATHOLOGY REPORT [SYNTHETIC]

ACCESSION #: UC-2025-87689

DATE OF PROCEDURE: 05/03/2025

DATE OF REPORT: 05/05/2025

REQUESTING PHYSICIAN: Dr. Megan Brown, Gastroenterology

PATHOLOGIST: Dr. Anthony Smith, Anatomic Pathology

CLINICAL HISTORY:

57 year old female with 2 week history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed continuous erythema and friability from rectum to splenic flexure. Clinical suspicion for ulcerative colitis.

SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 2 tan-pink tissue fragments measuring 6 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 4 tan-pink tissue fragments measuring 4 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 6 tan-pink tissue fragments measuring 5 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 3 tan-pink tissue fragments measuring 2 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 6 tan-pink tissue fragments measuring 4 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 3 tan-pink tissue fragments measuring 8 mm in aggregate.

All specimens are entirely submitted in 4 cassette(s).

MICROSCOPIC DESCRIPTION:

A. Rectal mucosa shows fulminant active chronic inflammation with crypt architectural distortion, lamina propria plasma cells, and basal plasmacytosis. The inflammatory process is limited to the mucosa without evidence of granulomas. Marked decrease in goblet cell population.

B. Sigmoid colonic mucosa shows mild to moderate active chronic inflammation with severe cryptitis, crypt architectural distortion, and Paneth cell metaplasia. The inflammatory process is limited to the mucosa without evidence of granulomas. Lamina propria shows increased plasma cells and lymphocytes.

C. Descending colonic mucosa shows moderate active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Lamina propria shows increased plasma cells and lymphocytes.

D. Transverse colonic mucosa shows moderate active chronic inflammation with crypt architectural distortion and crypt abscesses. Lamina propria shows increased plasma cells and lymphocytes.

E. Ascending colonic mucosa shows mild to moderate active chronic inflammation with crypt architectural distortion and crypt abscesses. Occasional apoptotic bodies are present in crypts.

F. Terminal ileal mucosa shows essentially normal ileal mucosa with intact villous architecture and no active inflammation. No evidence of chronic inflammatory bowel disease identified in this section.

DIAGNOSIS:

A. Rectum, biopsy:

- fulminant active chronic colitis with crypt architectural distortion and goblet cell depletion
- fulminant consistent with ulcerative colitis
- No dysplasia identified

- No evidence of cytomegalovirus (CMV) infection

B. Sigmoid colon, biopsy:

- mild to moderate active chronic colitis with crypt architectural distortion and goblet cell depletion
- mild to moderate consistent with ulcerative colitis
- No dysplasia identified

C-E. Descending, transverse, and ascending colon, biopsies:

- moderate active chronic colitis with crypt architectural distortion
- Features consistent with ulcerative colitis
- No dysplasia identified

F. Terminal ileum, biopsy:

- Mild non-specific inflammation
- No evidence of inflammatory bowel disease

COMMENT:

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. Clinical correlation and follow-up biopsies are recommended to monitor disease activity and treatment response.

SPECIAL STUDIES:

Acid-fast bacilli (AFB) stain is negative for mycobacterial organisms.

This is a synthetic educational pathology report created for AI training purposes. It does not represent a real patient case.