

SURGICAL PATHOLOGY REPORT [SYNTHETIC]

ACCESSION #: UC-2025-89151

DATE OF PROCEDURE: 04/24/2025

DATE OF REPORT: 04/27/2025

REQUESTING PHYSICIAN: Dr. Cynthia Gonzalez, Gastroenterology

PATHOLOGIST: Dr. Matthew Middleton, Anatomic Pathology

CLINICAL HISTORY:

46 year old male with 1 year history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed moderate erythema and loss of vascular pattern from rectum to descending colon. Clinical suspicion for ulcerative colitis.

SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 5 tan-pink tissue fragments measuring 5 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 4 tan-pink tissue fragments measuring 2 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 2 tan-pink tissue fragments measuring 2 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 2 tan-pink tissue fragments measuring 6 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 6 tan-pink tissue fragments measuring 8 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 4 tan-pink tissue fragments measuring 7 mm in aggregate.

All specimens are entirely submitted in 2 cassette(s).

MICROSCOPIC DESCRIPTION:

A. Rectal mucosa shows fulminant active chronic inflammation with severe cryptitis, crypt architectural distortion, and mucosal ulceration. The inflammatory process is limited to the mucosa without evidence of granulomas. Occasional apoptotic bodies are present in crypts.

B. Sigmoid colonic mucosa shows moderate active chronic inflammation with marked crypt architectural distortion, numerous crypt abscesses, and complete goblet cell depletion. The inflammatory process is limited to the mucosa without evidence of granulomas. No evidence of dysplasia is identified.

C. Descending colonic mucosa shows moderate to severe active chronic inflammation with crypt branching, crypt atrophy, and focal crypt abscesses. The inflammatory process is limited to the mucosa without evidence of granulomas. No evidence of dysplasia is identified.

D. Transverse colonic mucosa shows moderate active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. Reactive epithelial changes are seen adjacent to areas of active inflammation.

E. Ascending colonic mucosa shows mild active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. Reactive epithelial changes are seen adjacent to areas of active inflammation.

F. Terminal ileal mucosa shows normal small intestinal mucosa with preserved villous architecture. No evidence of chronic inflammatory bowel disease identified in this section.

DIAGNOSIS:

A. Rectum, biopsy:

- fulminant active chronic colitis with crypt architectural distortion and goblet cell depletion
- fulminant consistent with ulcerative colitis
- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

B. Sigmoid colon, biopsy:

- moderate active chronic colitis with crypt architectural distortion and goblet cell depletion
- moderate consistent with ulcerative colitis
- No dysplasia identified

C-E. Descending, transverse, and ascending colon, biopsies:

- moderate to severe active chronic colitis with crypt architectural distortion
- Features consistent with ulcerative colitis
- No dysplasia identified

F. Terminal ileum, biopsy:

- Mild non-specific inflammation
- No evidence of inflammatory bowel disease

COMMENT:

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The histologic findings show classic features of ulcerative colitis with diffuse crypt architectural distortion and diffuse mucosal inflammation.

SPECIAL STUDIES:

Acid-fast bacilli (AFB) stain is negative for mycobacterial organisms.