

# SURGICAL PATHOLOGY REPORT [SYNTHETIC]

**ACCESSION #:** UC-2025-22578

**DATE OF PROCEDURE:** 04/23/2025

**DATE OF REPORT:** 04/25/2025

**REQUESTING PHYSICIAN:** Dr. David Castaneda, Gastroenterology

**PATHOLOGIST:** Dr. Thomas Hart, Anatomic Pathology

## CLINICAL HISTORY:

52 year old male with recent onset history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed severe friability, superficial ulcerations, and pseudopolyps throughout the colon. Clinical suspicion for ulcerative colitis.

## SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

## GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 4 tan-pink tissue fragments measuring 3 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 3 tan-pink tissue fragments measuring 8 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 5 tan-pink tissue fragments measuring 7 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 4 tan-pink tissue fragments measuring 4 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 4 tan-pink tissue fragments measuring 2 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 2 tan-pink tissue fragments measuring 8 mm in aggregate.

All specimens are entirely submitted in 3 cassette(s).

## **MICROSCOPIC DESCRIPTION:**

A. Rectal mucosa shows fulminant active chronic inflammation with crypt branching, crypt atrophy, and focal crypt abscesses. The inflammatory process is limited to the mucosa without evidence of granulomas. Reactive epithelial changes are seen adjacent to areas of active inflammation.

B. Sigmoid colonic mucosa shows mild to moderate active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. No evidence of dysplasia is identified.

C. Descending colonic mucosa shows mild to moderate active chronic inflammation with severe cryptitis, crypt architectural distortion, and Paneth cell metaplasia. The inflammatory process is limited to the mucosa without evidence of granulomas. Basal plasmacytosis is prominent.

D. Transverse colonic mucosa shows moderate active chronic inflammation with marked crypt architectural distortion, numerous crypt abscesses, and complete goblet cell depletion. Lamina propria shows increased plasma cells and lymphocytes.

E. Ascending colonic mucosa shows mild active chronic inflammation with crypt architectural distortion and crypt abscesses. Mucosal edema and congestion are present.

F. Terminal ileal mucosa shows essentially normal ileal mucosa with intact villous architecture and no active inflammation. No evidence of chronic inflammatory bowel disease identified in this section.

## **DIAGNOSIS:**

### **A. Rectum, biopsy:**

- fulminant active chronic colitis with crypt architectural distortion and goblet cell depletion
- fulminant consistent with ulcerative colitis
- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

***B. Sigmoid colon, biopsy:***

- mild to moderate active chronic colitis with crypt architectural distortion and goblet cell depletion
- mild to moderate consistent with ulcerative colitis
- No dysplasia identified

***C-E. Descending, transverse, and ascending colon, biopsies:***

- mild to moderate active chronic colitis with crypt architectural distortion
- Features consistent with ulcerative colitis
- No dysplasia identified

***F. Terminal ileum, biopsy:***

- Mild non-specific inflammation
- No evidence of inflammatory bowel disease

**COMMENT:**

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The overall histologic features are characteristic of ulcerative colitis in the active phase.

**SPECIAL STUDIES:**

CD3 and CD20 immunostains show a normal distribution of T and B lymphocytes without evidence of lymphoma.