

# SURGICAL PATHOLOGY REPORT [SYNTHETIC]

**ACCESSION #:** UC-2025-62834

**DATE OF PROCEDURE:** 04/18/2025

**DATE OF REPORT:** 04/21/2025

**REQUESTING PHYSICIAN:** Dr. Jacob Harris, Gastroenterology

**PATHOLOGIST:** Dr. Kyle Campbell, Anatomic Pathology

## CLINICAL HISTORY:

48 year old female with 5 year history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed circumferential ulceration and pseudopolyps from rectum to mid-transverse colon. Clinical suspicion for ulcerative colitis.

## SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

## GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 5 tan-pink tissue fragments measuring 7 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 5 tan-pink tissue fragments measuring 3 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 5 tan-pink tissue fragments measuring 3 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 4 tan-pink tissue fragments measuring 2 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 6 tan-pink tissue fragments measuring 8 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 5 tan-pink tissue fragments measuring 8 mm in aggregate.

All specimens are entirely submitted in 1 cassette(s).

## **MICROSCOPIC DESCRIPTION:**

A. Rectal mucosa shows severe active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Basal plasmacytosis is prominent.

B. Sigmoid colonic mucosa shows severe active chronic inflammation with crypt branching, crypt atrophy, and focal crypt abscesses. The inflammatory process is limited to the mucosa without evidence of granulomas. Occasional Paneth cell metaplasia is noted.

C. Descending colonic mucosa shows mild to moderate active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Occasional Paneth cell metaplasia is noted.

D. Transverse colonic mucosa shows mild active chronic inflammation with crypt architectural distortion and crypt abscesses. Mucosal edema and congestion are present.

E. Ascending colonic mucosa shows mild to moderate active chronic inflammation with crypt architectural distortion and crypt abscesses. Basal plasmacytosis is prominent.

F. Terminal ileal mucosa shows normal small intestinal mucosa with preserved villous architecture. No evidence of chronic inflammatory bowel disease identified in this section.

## **DIAGNOSIS:**

### ***A. Rectum, biopsy:***

- severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- severe consistent with ulcerative colitis
- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

### ***B. Sigmoid colon, biopsy:***

- severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- severe consistent with ulcerative colitis
- No dysplasia identified

***C-E. Descending, transverse, and ascending colon, biopsies:***

- mild to moderate active chronic colitis with crypt architectural distortion
- Features consistent with ulcerative colitis
- No dysplasia identified

***F. Terminal ileum, biopsy:***

- Mild non-specific inflammation
- No evidence of inflammatory bowel disease

**COMMENT:**

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The lack of granulomas, ileal involvement, and transmural inflammation favors ulcerative colitis over Crohn's disease.

**SPECIAL STUDIES:**

Acid-fast bacilli (AFB) stain is negative for mycobacterial organisms.

\_This is a synthetic educational pathology report created for AI training purposes. It does not represent a real patient case.\_