

# **SURGICAL PATHOLOGY REPORT [SYNTHETIC]**

**ACCESSION #:** UC-2025-44652

**DATE OF PROCEDURE:** 04/19/2025

**DATE OF REPORT:** 04/21/2025

**REQUESTING PHYSICIAN:** Dr. Leroy Rubio DVM, Gastroenterology

**PATHOLOGIST:** Dr. Kathleen Carroll, Anatomic Pathology

## **CLINICAL HISTORY:**

73 year old male with 5 year history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed circumferential ulceration and pseudopolyps from rectum to mid-transverse colon. Clinical suspicion for ulcerative colitis.

## **SPECIMEN RECEIVED:**

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

## **GROSS DESCRIPTION:**

- A. Received in formalin labeled "rectum" are 4 tan-pink tissue fragments measuring 2 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 6 tan-pink tissue fragments measuring 7 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 2 tan-pink tissue fragments measuring 2 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 3 tan-pink tissue fragments measuring 3 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 2 tan-pink tissue fragments measuring 8 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 2 tan-pink tissue fragments measuring 2 mm in aggregate.

All specimens are entirely submitted in 4 cassette(s).

## **MICROSCOPIC DESCRIPTION:**

A. Rectal mucosa shows moderate to severe active chronic inflammation with crypt architectural distortion, lamina propria plasma cells, and basal plasmacytosis. The inflammatory process is limited to the mucosa without evidence of granulomas. Basal plasmacytosis is prominent.

B. Sigmoid colonic mucosa shows severe active chronic inflammation with marked crypt architectural distortion, numerous crypt abscesses, and complete goblet cell depletion. The inflammatory process is limited to the mucosa without evidence of granulomas. Surface epithelium shows reactive changes.

C. Descending colonic mucosa shows mild to moderate active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Occasional apoptotic bodies are present in crypts.

D. Transverse colonic mucosa shows moderate active chronic inflammation with crypt architectural distortion and crypt abscesses. No evidence of dysplasia is identified.

E. Ascending colonic mucosa shows mild to moderate active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. Occasional apoptotic bodies are present in crypts.

F. Terminal ileal mucosa shows mild non-specific inflammation without architectural distortion. No evidence of chronic inflammatory bowel disease identified in this section.

## **DIAGNOSIS:**

### ***A. Rectum, biopsy:***

- moderate to severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- moderate to severe consistent with ulcerative colitis
- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

***B. Sigmoid colon, biopsy:***

- severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- severe consistent with ulcerative colitis
- No dysplasia identified

***C-E. Descending, transverse, and ascending colon, biopsies:***

- mild to moderate active chronic colitis with crypt architectural distortion
- Features consistent with ulcerative colitis
- No dysplasia identified

***F. Terminal ileum, biopsy:***

- Mild non-specific inflammation
- No evidence of inflammatory bowel disease

**COMMENT:**

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The histologic findings show classic features of ulcerative colitis with diffuse crypt architectural distortion and diffuse mucosal inflammation.

**SPECIAL STUDIES:**

Periodic acid-Schiff (PAS) stain is negative for fungal organisms.