

SURGICAL PATHOLOGY REPORT [SYNTHETIC]

ACCESSION #: UC-2025-23273

DATE OF PROCEDURE: 05/02/2025

DATE OF REPORT: 05/04/2025

REQUESTING PHYSICIAN: Dr. Rhonda Carter, Gastroenterology

PATHOLOGIST: Dr. Amy Adams MD, Anatomic Pathology

CLINICAL HISTORY:

69 year old male with 3 week history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed pancolitis with diffuse ulceration and spontaneous bleeding. Clinical suspicion for ulcerative colitis.

SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 6 tan-pink tissue fragments measuring 4 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 2 tan-pink tissue fragments measuring 6 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 4 tan-pink tissue fragments measuring 4 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 6 tan-pink tissue fragments measuring 4 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 2 tan-pink tissue fragments measuring 3 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 5 tan-pink tissue fragments measuring 2 mm in aggregate.

All specimens are entirely submitted in 4 cassette(s).

MICROSCOPIC DESCRIPTION:

A. Rectal mucosa shows moderate active chronic inflammation with severe cryptitis, crypt architectural distortion, and mucosal ulceration. The inflammatory process is limited to the mucosa without evidence of granulomas. Surface epithelium shows reactive changes.

B. Sigmoid colonic mucosa shows mild to moderate active chronic inflammation with crypt architectural distortion, lamina propria plasma cells, and basal plasmacytosis. The inflammatory process is limited to the mucosa without evidence of granulomas. Marked decrease in goblet cell population. Areas of crypt dropout and lamina propria fibrosis are present, suggesting chronicity and possible treatment effect.

C. Descending colonic mucosa shows mild to moderate active chronic inflammation with severe cryptitis, crypt architectural distortion, and Paneth cell metaplasia. The inflammatory process is limited to the mucosa without evidence of granulomas. Basal plasmacytosis is prominent.

D. Transverse colonic mucosa shows mild active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. No evidence of dysplasia is identified.

E. Ascending colonic mucosa shows mild to moderate active chronic inflammation with crypt architectural distortion and crypt abscesses. No evidence of dysplasia is identified.

F. Terminal ileal mucosa shows no significant pathologic abnormality. No evidence of chronic inflammatory bowel disease identified in this section.

DIAGNOSIS:

A. Rectum, biopsy:

- moderate active chronic colitis with crypt architectural distortion and goblet cell depletion
- moderate consistent with ulcerative colitis
- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

B. Sigmoid colon, biopsy:

- mild to moderate active chronic colitis with crypt architectural distortion and goblet cell depletion
- mild to moderate consistent with ulcerative colitis
- No dysplasia identified

C-E. Descending, transverse, and ascending colon, biopsies:

- mild to moderate active chronic colitis with crypt architectural distortion
- Changes consistent with chronicity and treatment effect
- Features consistent with ulcerative colitis
- No dysplasia identified

F. Terminal ileum, biopsy:

- Mild non-specific inflammation
- No evidence of inflammatory bowel disease

COMMENT:

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The pattern of inflammation is consistent with ulcerative colitis as evidenced by the continuous mucosal involvement with greatest intensity distally. Histologic features suggesting chronicity and treatment effect are present. Correlation with treatment history is recommended.

SPECIAL STUDIES:

Cytomegalovirus (CMV) immunohistochemistry is negative for viral inclusions.

This is a synthetic educational pathology report created for AI training purposes. It does not represent a real patient case.