

SURGICAL PATHOLOGY REPORT [SYNTHETIC]

ACCESSION #: UC-2025-50777

DATE OF PROCEDURE: 05/06/2025

DATE OF REPORT: 05/10/2025

REQUESTING PHYSICIAN: Dr. Sarah Black, Gastroenterology

PATHOLOGIST: Dr. Karen Simmons, Anatomic Pathology

CLINICAL HISTORY:

64 year old male with longstanding history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed diffuse erythema, loss of vascular pattern, and contact bleeding from rectum to hepatic flexure. Clinical suspicion for ulcerative colitis.

SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 4 tan-pink tissue fragments measuring 6 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 4 tan-pink tissue fragments measuring 7 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 2 tan-pink tissue fragments measuring 5 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 4 tan-pink tissue fragments measuring 2 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 4 tan-pink tissue fragments measuring 7 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 2 tan-pink tissue fragments measuring 7 mm in aggregate.

All specimens are entirely submitted in 3 cassette(s).

MICROSCOPIC DESCRIPTION:

A. Rectal mucosa shows severe active chronic inflammation with severe cryptitis, crypt architectural distortion, and Paneth cell metaplasia. The inflammatory process is limited to the mucosa without evidence of granulomas. Occasional Paneth cell metaplasia is noted.

B. Sigmoid colonic mucosa shows moderate active chronic inflammation with marked crypt architectural distortion, numerous crypt abscesses, and complete goblet cell depletion. The inflammatory process is limited to the mucosa without evidence of granulomas. Occasional apoptotic bodies are present in crypts.

C. Descending colonic mucosa shows moderate active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. The inflammatory process is limited to the mucosa without evidence of granulomas. Marked decrease in goblet cell population.

D. Transverse colonic mucosa shows mild active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. Occasional apoptotic bodies are present in crypts.

E. Ascending colonic mucosa shows mild active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. Mucosal edema and congestion are present.

F. Terminal ileal mucosa shows mild reactive lymphoid hyperplasia without evidence of chronic inflammatory bowel disease. No evidence of chronic inflammatory bowel disease identified in this section.

DIAGNOSIS:

A. Rectum, biopsy:

- severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- severe consistent with ulcerative colitis
- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

B. Sigmoid colon, biopsy:

- moderate active chronic colitis with crypt architectural distortion and goblet cell depletion
- moderate consistent with ulcerative colitis
- No dysplasia identified

C-E. Descending, transverse, and ascending colon, biopsies:

- moderate active chronic colitis with crypt architectural distortion
- Features consistent with ulcerative colitis
- No dysplasia identified

F. Terminal ileum, biopsy:

- Mild non-specific inflammation
- No evidence of inflammatory bowel disease

COMMENT:

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The presence of diffuse crypt architectural distortion, basal plasmacytosis, and continuous inflammatory pattern strongly supports the diagnosis of ulcerative colitis.

SPECIAL STUDIES:

Cytomegalovirus (CMV) immunohistochemistry is negative for viral inclusions.