

SURGICAL PATHOLOGY REPORT [SYNTHETIC]

ACCESSION #: UC-2025-19873

DATE OF PROCEDURE: 04/24/2025

DATE OF REPORT: 04/26/2025

REQUESTING PHYSICIAN: Dr. Nathan Williams, Gastroenterology

PATHOLOGIST: Dr. Cassandra Anderson, Anatomic Pathology

CLINICAL HISTORY:

55 year old female with 1 month history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed continuous erythema and friability from rectum to splenic flexure. Clinical suspicion for ulcerative colitis.

SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 5 tan-pink tissue fragments measuring 2 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 3 tan-pink tissue fragments measuring 8 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 6 tan-pink tissue fragments measuring 8 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 5 tan-pink tissue fragments measuring 5 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 6 tan-pink tissue fragments measuring 8 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 3 tan-pink tissue fragments measuring 7 mm in aggregate.

All specimens are entirely submitted in 4 cassette(s).

MICROSCOPIC DESCRIPTION:

A. Rectal mucosa shows severe active chronic inflammation with severe cryptitis, crypt architectural distortion, and mucosal ulceration. The inflammatory process is limited to the mucosa without evidence of granulomas. Mucosal edema and congestion are present.

B. Sigmoid colonic mucosa shows severe active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Reactive epithelial changes are seen adjacent to areas of active inflammation. Rare cells with intranuclear and cytoplasmic inclusions suspicious for cytomegalovirus (CMV) infection are identified.

C. Descending colonic mucosa shows mild to moderate active chronic inflammation with crypt branching, crypt atrophy, and focal crypt abscesses. The inflammatory process is limited to the mucosa without evidence of granulomas. Basal plasmacytosis is prominent. Terminal ileal mucosa shows mild active inflammation with neutrophilic cryptitis, likely representing backwash ileitis.

D. Transverse colonic mucosa shows mild active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. Lamina propria shows increased plasma cells and lymphocytes.

E. Ascending colonic mucosa shows mild to moderate active chronic inflammation with crypt architectural distortion and crypt abscesses. No evidence of dysplasia is identified.

F. Terminal ileal mucosa shows mild reactive lymphoid hyperplasia without evidence of chronic inflammatory bowel disease. No evidence of chronic inflammatory bowel disease identified in this section.

DIAGNOSIS:

A. Rectum, biopsy:

- severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- severe consistent with ulcerative colitis
- No dysplasia identified

- No evidence of cytomegalovirus (CMV) infection

B. Sigmoid colon, biopsy:

- severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- severe consistent with ulcerative colitis
- No dysplasia identified

C-E. Descending, transverse, and ascending colon, biopsies:

- mild to moderate active chronic colitis with crypt architectural distortion
- Mild active ileitis, consistent with backwash ileitis in the setting of ulcerative colitis
- Viral cytopathic changes suspicious for cytomegalovirus (CMV) infection
- Features consistent with ulcerative colitis
- No dysplasia identified

F. Terminal ileum, biopsy:

- Mild non-specific inflammation
- No evidence of inflammatory bowel disease

COMMENT:

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The lack of granulomas, ileal involvement, and transmural inflammation favors ulcerative colitis over Crohn's disease. Immunohistochemical staining for CMV is positive, confirming the presence of CMV infection. This may contribute to the severity of colitis and should be considered in treatment planning. The mild ileal inflammation in the context of pancolitis is consistent with backwash ileitis, which can be seen in ulcerative colitis and does not necessarily indicate Crohn's disease.

SPECIAL STUDIES:

No special stains were performed. Cytomegalovirus (CMV) immunohistochemistry reveals scattered positive cells confirming viral infection.

This is a synthetic educational pathology report created for AI training purposes. It does not represent a real patient case.