

SURGICAL PATHOLOGY REPORT [SYNTHETIC]

ACCESSION #: UC-2025-24626

DATE OF PROCEDURE: 04/23/2025

DATE OF REPORT: 04/27/2025

REQUESTING PHYSICIAN: Dr. Thomas Griffin, Gastroenterology

PATHOLOGIST: Dr. Terri Cameron, Anatomic Pathology

CLINICAL HISTORY:

35 year old male with 1 year history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed severe friability, superficial ulcerations, and pseudopolyps throughout the colon. Clinical suspicion for ulcerative colitis.

SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 6 tan-pink tissue fragments measuring 8 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 5 tan-pink tissue fragments measuring 3 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 2 tan-pink tissue fragments measuring 2 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 3 tan-pink tissue fragments measuring 7 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 4 tan-pink tissue fragments measuring 7 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 4 tan-pink tissue fragments measuring 8 mm in aggregate.

All specimens are entirely submitted in 2 cassette(s).

MICROSCOPIC DESCRIPTION:

A. Rectal mucosa shows moderate active chronic inflammation with crypt branching, crypt atrophy, and focal crypt abscesses. The inflammatory process is limited to the mucosa without evidence of granulomas. Lamina propria shows increased plasma cells and lymphocytes.

B. Sigmoid colonic mucosa shows moderate active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. No evidence of dysplasia is identified.

C. Descending colonic mucosa shows mild to moderate active chronic inflammation with marked crypt architectural distortion, numerous crypt abscesses, and complete goblet cell depletion. The inflammatory process is limited to the mucosa without evidence of granulomas. No evidence of dysplasia is identified.

D. Transverse colonic mucosa shows mild to moderate active chronic inflammation with marked crypt architectural distortion, numerous crypt abscesses, and complete goblet cell depletion. Marked decrease in goblet cell population.

E. Ascending colonic mucosa shows mild to moderate active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. Mucosal edema and congestion are present.

F. Terminal ileal mucosa shows normal small intestinal mucosa with preserved villous architecture. No evidence of chronic inflammatory bowel disease identified in this section. Areas of crypt dropout and lamina propria fibrosis are present, suggesting chronicity and possible treatment effect.

DIAGNOSIS:

A. Rectum, biopsy:

- moderate active chronic colitis with crypt architectural distortion and goblet cell depletion
- moderate consistent with ulcerative colitis
- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

B. Sigmoid colon, biopsy:

- moderate active chronic colitis with crypt architectural distortion and goblet cell depletion
- Changes consistent with chronicity and treatment effect
- moderate consistent with ulcerative colitis
- No dysplasia identified

C-E. Descending, transverse, and ascending colon, biopsies:

- mild to moderate active chronic colitis with crypt architectural distortion
- Features consistent with ulcerative colitis
- No dysplasia identified

F. Terminal ileum, biopsy:

- Mild non-specific inflammation
- No evidence of inflammatory bowel disease

COMMENT:

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The lack of granulomas, ileal involvement, and transmural inflammation favors ulcerative colitis over Crohn's disease. Histologic features suggesting chronicity and treatment effect are present. Correlation with treatment history is recommended.

SPECIAL STUDIES:

Acid-fast bacilli (AFB) stain is negative for mycobacterial organisms.

This is a synthetic educational pathology report created for AI training purposes. It does not represent a real patient case.