

# SURGICAL PATHOLOGY REPORT [SYNTHETIC]

**ACCESSION #:** UC-2025-71009

**DATE OF PROCEDURE:** 04/12/2025

**DATE OF REPORT:** 04/16/2025

**REQUESTING PHYSICIAN:** Dr. Charles Ortiz, Gastroenterology

**PATHOLOGIST:** Dr. Keith Brown, Anatomic Pathology

## CLINICAL HISTORY:

47 year old male with 2 week history of bloody diarrhea, abdominal pain, and urgency. Colonoscopy showed pancolitis with diffuse ulceration and spontaneous bleeding. Clinical suspicion for ulcerative colitis.

## SPECIMEN RECEIVED:

- A. Rectum, biopsy
- B. Sigmoid colon, biopsy
- C. Descending colon, biopsy
- D. Transverse colon, biopsy
- E. Ascending colon, biopsy
- F. Terminal ileum, biopsy

## GROSS DESCRIPTION:

- A. Received in formalin labeled "rectum" are 4 tan-pink tissue fragments measuring 2 mm in aggregate.
- B. Received in formalin labeled "sigmoid colon" are 2 tan-pink tissue fragments measuring 4 mm in aggregate.
- C. Received in formalin labeled "descending colon" are 6 tan-pink tissue fragments measuring 3 mm in aggregate.
- D. Received in formalin labeled "transverse colon" are 3 tan-pink tissue fragments measuring 8 mm in aggregate.
- E. Received in formalin labeled "ascending colon" are 2 tan-pink tissue fragments measuring 7 mm in aggregate.

F. Received in formalin labeled "terminal ileum" are 2 tan-pink tissue fragments measuring 5 mm in aggregate.

All specimens are entirely submitted in 1 cassette(s).

## **MICROSCOPIC DESCRIPTION:**

A. Rectal mucosa shows fulminant active chronic inflammation with severe cryptitis, crypt architectural distortion, and Paneth cell metaplasia. The inflammatory process is limited to the mucosa without evidence of granulomas. Marked decrease in goblet cell population.

B. Sigmoid colonic mucosa shows severe active chronic inflammation with crypt architectural distortion, lamina propria plasma cells, and basal plasmacytosis. The inflammatory process is limited to the mucosa without evidence of granulomas. Mucosal edema and congestion are present. In addition to the chronic inflammatory changes, there are numerous neutrophils and pseudomembranes suspicious for superimposed *Clostridioides difficile* infection.

C. Descending colonic mucosa shows mild to moderate active chronic inflammation with diffuse neutrophilic cryptitis, crypt abscesses, and epithelial injury. The inflammatory process is limited to the mucosa without evidence of granulomas. Lamina propria shows increased plasma cells and lymphocytes.

D. Transverse colonic mucosa shows mild active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. Reactive epithelial changes are seen adjacent to areas of active inflammation.

E. Ascending colonic mucosa shows mild active chronic inflammation with diffuse crypt architectural distortion, crypt abscesses, and goblet cell depletion. Basal plasmacytosis is prominent.

F. Terminal ileal mucosa shows mild reactive lymphoid hyperplasia without evidence of chronic inflammatory bowel disease. No evidence of chronic inflammatory bowel disease identified in this section. The inflammatory pattern shows overlapping features of both ulcerative colitis and Crohn's disease.

## **DIAGNOSIS:**

### ***A. Rectum, biopsy:***

- fulminant active chronic colitis with crypt architectural distortion and goblet cell depletion
- Features of chronicity consistent with inflammatory bowel disease, with overlapping features of both UC and CD

- fulminant consistent with ulcerative colitis
- No dysplasia identified
- No evidence of cytomegalovirus (CMV) infection

#### ***B. Sigmoid colon, biopsy:***

- severe active chronic colitis with crypt architectural distortion and goblet cell depletion
- severe consistent with ulcerative colitis
- No dysplasia identified

#### ***C-E. Descending, transverse, and ascending colon, biopsies:***

- mild to moderate active chronic colitis with crypt architectural distortion
- Features consistent with ulcerative colitis
- No dysplasia identified

#### ***F. Terminal ileum, biopsy:***

- Mild non-specific inflammation
- Features suggestive of superimposed *Clostridioides difficile* infection
- No evidence of inflammatory bowel disease

### **COMMENT:**

The histologic findings show a pattern of continuous chronic active colitis with greatest severity in the distal colon and rectum, with relative sparing of the proximal colon. The absence of granulomas, transmural inflammation, and terminal ileal involvement are features favoring ulcerative colitis over Crohn's disease. Correlation with clinical, endoscopic, and radiologic findings is recommended for definitive classification. The lack of granulomas, ileal involvement, and transmural inflammation favors ulcerative colitis over Crohn's disease. The histologic features suggestive of superimposed *Clostridioides difficile* infection should be correlated with clinical presentation and stool testing. The histologic findings show overlapping features of both ulcerative colitis and Crohn's disease. This pattern may represent an 'indeterminate colitis' and correlation with clinical, endoscopic, and serologic markers is strongly recommended for further classification.

### **SPECIAL STUDIES:**

Grocott's methenamine silver (GMS) stain is negative for fungal organisms. Gram stain highlights numerous gram-positive bacilli morphologically consistent with *Clostridioides difficile*.

\_This is a synthetic educational pathology report created for AI training purposes. It does not represent a real patient case.\_