

1 Transcripts with high distal heritability mediate genetic effects on
2 complex traits

3

4 **Abstract**

5 The transcriptome is increasingly viewed as a bridge between genetic risk factors for complex disease and
6 their associated pathophysiology. Powerful insights into disease mechanism can be made by linking genetic
7 variants affecting gene expression (expression quantitative trait loci - eQTLs) to phenotypes.

8 **Introduction**

9 In the quest to understand the genetic architecture of complex traits, gene expression is an important bridge
10 between genotype and phenotype. By identifying mediating transcripts, we get one step closer to a molecular
11 understanding of how genetic variants influence traits. Moreover, there is evidence from genome-wide
12 association studies (GWAS) that regulation of gene expression accounts for the bulk of the genetic effect
13 on complex traits, as most trait-associated variants lie in gene regulatory regions [1, 2, 3, 4, 5, 6, 7]. It is
14 widely assumed that these variants influence local transcription, and methods such as transcriptome-wide
15 association studies (TWAS) [8, 9, 10, 11], summary data-based Mendelian randomization (SMR) [10], and
16 others have capitalized on this idea to identify genes associated with multiple disease traits [12, 13, 14, 15]

17 Despite the great promise of these methods, however, they have not been as widely successful as it seemed
18 they could have been, and the vast majority of complex trait heritability remains unexplained. Although
19 trait-associated variants tend to lie in non-coding, regulatory regions, they often do not have detectable effects
20 on gene expression [16] and tend not to co-localize with expression quantitative trait loci (eQTLs) [17, 18].

21 One possible explanation for these observations is that gene expression is not being measured in the appropriate
22 cell types and thus true eQTLs influencing traits cannot be detected [16]. An alternative explanation that
23 has been discussed in recent years is that effects of these variants are mediated not through local regulation
24 of gene expression, but through distal regulation [18, 19, 20, 15].

25 However, assessing the role of wide-spread distal gene regulation on complex traits requires large, dedicated data

26 sets that include high-dimensional, clinically relevant phenotyping, dense genotyping in a highly recombined
27 population, and transcriptome-wide measurements of gene expression in multiple tissues. Measuring gene
28 expression in multiple tissues is critical to adequately assess the extent to which local gene regulation varies
29 across multiple tissues and whether such variability might account for previous failed attempts to identify
30 trait-relevant local eQTL. Such data sets are extremely difficult to obtain in human populations, particularly
31 in the large numbers of subjects required for statistical testing. Thus, to investigate further the role of local
32 and distal gene regulation on complex traits, we have generated an appropriate data set in a large population
33 of diversity outbred (DO) mice [21] in a population model of diet-induced obesity and metabolic disease [12].

34 The DO mice were derived from eight inbred founder mouse strains, five classical lab strains, and three
35 strains more recently derived from wild mice [21]. They represent three subspecies of mouse *Mus musculus*
36 *domesticus*, *Mus musculus musculus*, and *Mus musculus castaneus*, and capture 90% of the known variation
37 in laboratory mice [cite]. They are maintained with a breeding scheme that ensures equal contributions from
38 each founder across the genome thus rendering almost the whole genome visible to genetic inquiry [21]. We
39 measured clinically relevant metabolic traits, including body weight, plasma levels of insulin and glucose,
40 and plasma lipids in 500 DO mice. We further measured transcriptome-wide gene expression in four tissues
41 related to metabolic disease: adipose tissue, pancreatic islets, liver, and skeletal muscle.

42 To assess the role of gene regulation in mediating variation in metabolic traits in this population, we propose
43 high-dimensional mediation (HDM). In univariate approaches, such as TWAS, SMR, and other Mendelian
44 randomization approaches, each transcript is tested independently for mediation of a local variant on a
45 trait. This process requires huge numbers of statistical tests, which is computationally expensive, requires
46 strict corrections for multiple testing, and assumes independence of genetic variants and transcripts. Such
47 methods are therefore limited to detecting only the largest statistical effects and are biased toward local gene
48 regulation. In contrast, with high-dimensional mediation we assessed broad relationships among the genome,
49 transcriptome, and phenotype as a whole and identified a highly heritable composite trait that was perfectly
50 mediated by a composite transcript. We show that composite transcripts were tissue-specific and highly
51 interpretable in terms of biological processes as well as cell type composition. Heritability analysis of the
52 transcripts showed that the strongest transcriptional mediators of metabolic disease had low local heritability
53 and high distal heritability. Finally, we show that the composite transcripts identified in the DO population
54 predicted obesity in an independent population of Collaborative Cross recombinant inbred (CC-RIX) mice
55 and in human subjects. In contrast, local eQTL were unable to predict obesity in the CC-RIX mice. Together
56 our results suggest that both the tissue used for gene expression analysis as well as distal gene regulation are
57 critically important in identifying transcriptional mediators of the genome on complex traits.

58 **Results**

59 **Genetic variation contributed to wide phenotypic variation**

60 Although the environment was consistent across all animals, the genetic diversity present in this population
61 resulted in widely varying distributions across physiological measurements (Fig. 1). For example, body
62 weights of adult individuals varied from less than the average adult B6 body weight to several times the body
63 weight of a B6 adult in both sexes (Fig. 1A). Fasting blood glucose (FBG) also varied considerably (Fig. 1B)
64 although few of the animals had FBG levels that would indicate pre-diabetes (animals,), or diabetes (7
65 animals, 1.4) according to previously developed cutoffs (pre-diabetes: $\text{FBG} \geq 250 \text{ mg/dL}$, diabetes: $\text{FBG} \geq$
66 300, mg/dL) [22]. Males had higher FBG than females on average (Fig. 1C) as has been observed before
67 suggesting either that males were more susceptible to metabolic disease on the high-fat diet, or that males
68 and females may require different thresholds for pre-diabetes and diabetes.

69 Body weight was strongly positively correlated with food consumption (Fig. 1D $R^2 = 0.51, p = 1.5 \times 10^{-75}$)
70 and fasting blood glucose (FBG) (Fig. 1E, $R^2 = 0.21, p = 1.4 \times 10^{-26}$) suggesting a link between behavioral
71 factors and metabolic disease. However, the heritability of this trait and others (Fig. 1F) indicates that
72 background genetics contribute substantially to correlates of metabolic disease in this population.

73 The landscape of trait correlations (Fig. 1G) shows that most of the metabolic trait pairs were relatively
74 weakly correlated indicating complex relationships among the measured traits. This low level of redundancy
75 suggests a broad sampling of multiple heritable aspects of metabolic disease including overall body weight,
76 glucose homeostasis, pancreatic composition and liver function.

77 **Distal Heritability Correlated with Phenotype Relevance**

78 We performed eQTL analysis using R/qltl2 [23] (Methods) and identified both local and distal eQTL for
79 transcripts in each of the four tissues (Supp. Fig 9). Significant local eQTLs far outnumbered distal eQTLs
80 (Supp. Fig. 9F) and tended to be shared across tissues (Supp. Fig. 9G) whereas the few significant distal
81 eQTL we identified tended to be tissue-specific (Supp. Fig. 9H)

82 We calculated the heritability of each transcript in terms of local and distal genetic factors (Methods). Overall,
83 local and distal genetic factors contributed approximately equally to transcript abundance. In all tissues,
84 both local and distal factors explained between 8 and 18% of the variance in the median transcript (Fig 2A).

85 Local heritability of transcripts was negatively correlated with their trait relevance, defined as the maximum
86 correlation of a transcript across all traits (Fig. 2B). This suggests that the more local genotype influenced
87 transcript abundance, the less effect variation in transcript abundance had on the measured traits. Conversely,




Figure 1: Clinical overview. **A.** Distributions of final body weight in the diversity outbred mice. Sex is indicated by color. The average B6 male and female adult weights at 24 weeks of age are indicated by blue and green bars on the x-axis. **B.** The distribution of final fasting glucose across the population split by sex. Normal, pre-diabetic, and diabetic fasting glucose levels for mice are shown by colored bars along the x-axis. **C.** Males had higher fasting blood glucose on average than females. **D.** The relationship between food consumption and body weight for both sexes. **E.** Relationship between body weight and fasting glucose for both sexes. **F.** Heritability estimates for each physiological trait. Bars show standard error of the estimate. **G.** Correlation structure between pairs of physiological traits.

distal heritability of transcripts was positively correlated with trait relevance (Fig. 2C). That is, transcripts that were more highly correlated with the measured traits tended to be distally, rather than locally, heritable. That trait-correlated transcripts have low local heritability is consistent with previous observations that low-heritability transcripts explain more expression-mediated disease heritability than high-heritability transcripts [19]. However, the positive relationship between trait correlation and distal heritability suggests that there are alternative mechanisms through which genetic regulation of transcripts may influence traits.




Figure 2: Transcript heritability and trait relevance. **A.** Distributions of distal and local heritability of transcripts across the four tissues. Overall local and distal factors contribute equally to transcript heritability. The relationship between **(B.)** local and **(C.)** distal heritability and trait relevance across all four tissues. Here trait relevance is defined as the maximum correlation between the transcript and all traits. Local heritability was negatively correlated with trait relevance, and distal heritability is positively correlated with trait relevance. Pearson (r) and p values for each correlation are shown in the upper-right of each panel.

- 94 **High-Dimensional Mediation identified a high-heritability composite trait that was perfectly
95 mediated by a composite transcript**
- 96 We used high-dimensional mediation to identify the major axis of variation in the transcriptome that mediated
97 the effects of the genome on metabolic traits (Fig. 3). We kernelized the genome, phenome, and transcriptome
98 matrices and used generalized canonical correlation analysis (RGCCA) [24] to identify a composite transcript
99 (T_C) that perfectly mediated the effect of the composite genome (G_C) on the composite phenome (P_C).
100 Fig. 3A shows the partial correlations (ρ) between the pairs of these composite vectors. The partial correlation
101 between G_C and T_S was 0.42, and the partial correlation between T_S and P_S was 0.78. However, when the




Figure 3: High-dimensional mediation. **A.** Workflow indicating major steps of high-dimensional mediation. The genotype, transcriptome, and phenotype matrices were kernelized to yield single matrices representing the relationships between all individuals for each data modality (G_K = genome kernel, T_K = transcriptome kernel; P_K = phenotype kernel). High-dimensional mediation was applied to these matrices to maximize the direct path $G \rightarrow T \rightarrow P$, the mediating pathway (arrows), while simultaneously minimizing the direct $G \rightarrow P$ pathway (dotted line). The composite vectors that resulted from high-dimensional mediation were G_c , T_c , and P_c . The partial correlations ρ between these vectors indicated perfect mediation. Transcript and trait loadings were calculated as described in the methods. **B.** The null distribution of the path coefficient derived from 10,000 permutations compared to the observed path coefficient (red line). **C.** The null distribution of the G_c-T_c correlation vs. the T_c-P_c correlation compared with the observed value (red dot).

- 102 transcriptome was taken into account, the partial correlation between G_S and P_S was effectively 0 (0.039).
- 103 The estimated heritability of the composite phenotype was heritability of 0.71 ± 0.084 , which was higher than
- 104 any of the individual traits (Fig. 1F). Thus, we have identified a maximally heritable metabolic trait that is
- 105 perfectly mediated by a heritable component of the transcriptome.
- 106 Standard CCA is prone to over-fitting because in any two large matrices it can be trivial to identify
- 107 highly correlated composite vectors. To assess whether RGCCA was similarly prone to over-fitting in

108 a high-dimensional space, we performed permutation testing. We permuted the individual labels on the
109 transcriptome kernel matrix 1000 times and recalculated the path coefficient, which is the partial correlation of
110 G_C and T_C multiplied by the partial correlation of T_C and P_C . This represents the path from G_C to P_C that is
111 mediated through T_C . The null distribution of the path coefficient is shown in Fig. 3B, and the observed path
112 coefficient from the original data is indicated by the red line. The observed path coefficient was well outside
113 the null distribution generated by permutations. Fig. 3C illustrates this observation in more detail. Although
114 we identified high correlations between G_C and T_C , and modest correlations between T_C and P_C in the null
115 data (Fig 3C), these two values could not be maximized simultaneously. The red dot shows that in the real
116 data both the G_C - T_C correlation and the T_C - P_C correlation could be maximized simultaneously suggesting
117 that the path from genotype to phenotype through transcriptome is highly non-trivial and identifiable in
118 this case. These results suggest that these composite vectors represent genetically determined variation in
119 phenotype that is mediated through genetically determined variation in transcription.

120 **Body weight and insulin resistance were highly represented in the expression-mediated composite trait**

122 The loadings of each measured trait onto P_C indicate how much each contributed to the composite phenotype.
123 Final body weight contributed the most (Fig. 4), followed by homeostatic insulin resistance (HOMA_IR) and
124 fasting plasma insulin levels (Insulin_Fasting). We can thus interpret P_C as an index of metabolic disease (Fig.
125 4B). Individuals with high values of P_C have a higher metabolic index and greater metabolic disease, including
126 higher body weight and higher insulin resistance. We refer to P_C as the metabolic index going forward. Traits
127 contributing the least to the metabolic index were measures of cholesterol and pancreas composition. Thus,
128 when we interpret the transcriptomic signature identified by HDM, we are explaining primarily transcriptional
129 mediation of body weight and insulin resistance, as opposed to cholesterol measurements.

130 **High-loading transcripts have low local heritability, high distal heritability, and were linked
131 mechanistically to obesity**

132 We interpreted large loadings onto transcripts as indicating strong mediation of the effect of genetics on
133 metabolic index. Large positive loadings indicate that higher expression was associated with a higher
134 metabolic index (i.e. higher risk of obesity and metabolic disease on the high-fat diet) (Fig. 4C). Conversely,
135 large negative loadings indicate that high expression of these transcripts was associated with a lower metabolic
136 index (i.e. lower risk of obesity and metabolic disease on the high-fat diet) (Fig. 4C). We used gene set
137 enrichment analysis (GSEA) [25, 26] to look for biological processes and pathways that were enriched at the
138 top and bottom of this list (Methods).




Figure 4: Interpretation of loadings. **A.** Loadings across traits. Body weight and insulin resistance contributed the most to the composite trait. **B.** Phenotype scores across individuals. Individuals with large positive phenotype scores had higher body weight and insulin resistance than average. Individuals with large negative phenotype scores had lower body weight and insulin resistance than average. **C.** Distribution of transcript loadings in adipose tissue. For transcripts with large positive loadings, higher expression was associated with higher phenotype scores. For transcripts with large negative loadings, higher expression was associated with lower phenotype scores. **D.** Distribution of absolute value of transcript loadings across tissues. Transcripts in adipose tissue had the largest loadings indicating that transcripts in adipose tissue were the best mediators of the genetic effects on body weight and insulin resistance.

- 139 In adipose tissue, both GO processes and KEGG pathway enrichments pointed to an axis of inflammation
 140 and metabolism (Supp. Fig. 10 and 11). GP terms and KEGG pathways associated with inflammation,
 141 particularly macrophage infiltration, were positively associated with metabolic index, indicating that increased
 142 expression in inflammatory pathways was associated with a higher metabolic index. It is well established
 143 that adipose tissue in obese individuals is highly inflamed [cite] and infiltrated by macrophages [cite], and the
 144 results here suggest that this may be a heritable component of metabolic disease.
- 145 The strongest negative enrichments in adipose tissue were related to mitochondrial activity in general, and
 146 thermogenesis in particular (Supp. Fig. 10 and 11). It has been shown mouse strains with greater thermogenic
 147 potential are also less susceptible to obesity on a high-fat diet [cite].
- 148 Transcripts associated with the citric acid (TCA) cycle as well as the catabolism of branched-chain amino


149 acids (BCAA), valine, leuceine, and isoleucine were strongly enriched with negative loadings in adipose
150 tissue (Supp. Fig. XXX). Expression of genes in both pathways (for which there is some overlap) has been
151 previously associated with insulin sensitivity [12, 27, 28], suggesting that heritable variation in regulation of
152 these pathways may influence risk of insulin resistance.

153 Looking at the 10 strongest positive and negative loaded transcripts from each tissue, it is apparent that
154 transcripts in the adipose tissue had the largest loadings, both positive and negative, of all tissues (Fig.
155 5A bar plot) This suggesting that much of the effect of genetics on body weight and insulin resistance
156 is mediated through gene expression in adipose tissue. The strongest loadings in liver and pancreas were
157 comparable, and those in skeletal muscle were the weakest (Fig. 5A), suggesting that less of the genetic
158 effects were mediated through transcription in skeletal muscle. Heritability analysis showed that transcripts
159 with the largest loadings tended to have relatively high distal heritability compared with local heritability
160 (Fig. 5A heat map and box plot). This pattern contrasts with transcripts nominated by TWAS (Fig. 5B),
161 which tended to have lower loadings, higher local heritability and lower distal heritability. Transcripts with
162 the highest local heritability in each tissue (Fig. 5C) had the lowest loadings.

163 We performed a literature search for the genes in each of these groups along with the terms “diabetes”,
164 “obesity”, and the name of the expressing tissue to determine whether any of these genes had previous
165 associations with metabolic disease in the literature (Methods). Multiple genes in each group had been
166 previously associated with obesity and diabetes (Fig. 5 bolded gene names). Genes with high loadings were
167 most highly enriched for previous literature support. They were 2.25 more likely than TWAS hits and 3.6
168 times more likely than genes with high local heritability to be previously associated with obesity or diabetes.

169 **Tissue-specific transcriptional programs were associated with metabolic traits**

170 Clustering of transcripts with top loadings in each tissue showed tissue-specific functional modules associated
171 with obesity and insulin resistance (Fig. 6A) (Methods). The clustering highlights the importance of immune
172 activation particularly in adipose tissue. Except for the “mitosis” cluster, which had large positive loadings in
173 three of the four tissues, all clusters were strongly loaded in only one or two tissues. For example, the lipid
174 metabolism cluster was loaded most heavily in liver. The positive loadings suggest that high expression of
175 these genes particularly in the liver was associated with increased metabolic disease. This cluster included
176 the gene *Pparg*, whose primary role is in the adipose tissue where it is considered a master regulator of
177 adipogenesis [29]. Agonists of *Pparg*, such as Thiazolidinediones, which are FDA-approved to treat type II
178 diabetes, reduce inflammation and adipose hypertrophy [29]. Consistent with this role, the loading for *Pparg*
179 in adipose tissue was negative, suggesting that higher expression was associated with leaner mice (Fig. 6B).



183 mice with a *Ppara* knockout, causes upregulation of genes involved in adipogenesis [32]. In the livers of both
184 mice and humans high *Pparg* expression is associated with hepatocytes that accumulate large lipid droplets
185 and have gene expression profiles similar to adipocytes [33, 34].

186 The local and distal heritability of *Pparg* is low in adipose tissue suggesting its expression in this tissue is
187 highly constrained in the population (Fig. 6B). However, the distal heritability of *Pparg* in liver is relatively
188 high suggesting it is complexly regulated and has sufficient variation in this population to drive variation
189 in phenotype. Both local and distal heritability of *Pparg* in the islet are fairly high, but the loading is
190 low, suggesting that variability of expression in the islet does not drive phenotypic variation. These results
191 highlight the importance of tissue context when investigating the role of heritable transcript variability in
192 driving phenotype.

193 Gene lists for all clusters are available in Supplemental File XXX.

194 **Gene expression, but not local eQTLs, predicted body weight in an independent population**

195 To test whether the transcript loadings identified in the DO could be translated to another population, we
196 tested whether they could predict metabolic a phenotype in an independent population of CC-RIX mice, which
197 were F1 mice derived from multiple pairings of Collaborative Cross (CC) [cite] strains (Fig. 7) (Methods).
198 We tested two questions. First, we asked whether the loadings identified in the DO mice were relevant to the
199 relationship between the transcriptome and the genome in the CC-RIX. We predicted body weight in each
200 CC-RIX individual using measured gene expression in each tissue and the transcript loadings identified in the
201 DO (Methods). The predicted body weight and actual body weight were highly correlated in all tissues (Fig.
202 7B left column). The best prediction was achieved for adipose tissue, which supports the observation in the
203 DO that adipose expression was the strongest mediator of the genetic effect on metabolic index. This result
204 also confirms the validity and translatability of the transcript loadings and their relationship to metabolic
205 disease.

206 The second question related to the source of the relevant variation in gene expression. If local regulation was
207 the predominant factor influencing gene expression, we should be able to predict phenotype in the CC-RIX
208 using transcripts imputed from local genotype (Fig. 7A). The DO and the CC-RIX were derived from the
209 same eight founder strains and so carry the same alleles throughout the genome. We imputed gene expression
210 in the CC-RIX using local genotype and were able to estimate variation in gene transcription robustly (Supp.
211 Fig. 12). However, these imputed values failed to predict body weight in the CC-RIX when weighted with
212 the loadings from HDM. (Fig. 7B right column). This result suggests that local regulation of gene expression
213 is not the primary factor driving heritability of complex traits.




Figure 6: Tissue-specific transcriptional programs were associated with obesity and insulin resistance. **A** Heat map showing the loadings of all transcripts with loadings greater than 2.5 standard deviations from the mean in any tissue. The heat map was clustered using k medoid clustering. Functional enrichments of each cluster are indicated along the left margin. **B** Loadings for *Pparg* in different tissues. **C** Local and distal of *Pparg* expression in different tissues.

214 **Distally heritable transcriptomic signatures reflected variation in composition of adipose tissue
215 and islets**

216 Interpretation of global genetic influences on gene expression and phenotype is potentially more challenging
217 than interpretation and translation of local genetic influences, as genetic effects cannot be localized to
218 individual gene variants or transcripts. However, there are global patterns across the loadings that can
219 inform mechanism. For example, heritable variation in cell type composition can be derived from transcript
220 loadings. We noted earlier that immune activation in the adipose tissues was an important driver of obesity
221 in the DO population. To determine whether this is reflected as an increase in macrophages in adipose
222 tissue, we compared loadings of cell-type specific genes in adipose tissue (Methods). The mean loading
223 of macrophage-specific genes was substantially greater than 0 (Fig. 8A), indicating that obese mice were




Figure 7: Transcription, but not local genotype, predicts phenotype in the CC-RIX. **A.** Workflow showing procedure for translating HDM results to an independent population of mice. **B.** Relationships between the predicted metabolic index and measured body weight. The left column shows the predictions using measured transcripts. The right column shows the prediction using transcript levels imputed from local genotype. Gray boxes indicate measured quantities, and blue boxes indicate calculated quantities. The dots in each panel represent individual CC-RIX strains. The gray lines show the standard deviation on body weight for the strain.

- genetically predisposed to have high levels of macrophage infiltration in adipose tissue in response to the high-fat, high-sugar diet.
- We also compared loadings of cell-type specific transcripts in islet (Methods). The mean loadings for alpha-cell specific transcripts were significantly greater than 0, while the mean loadings for delta- and endothelial-cell specific genes were significantly less than 0 (Fig. 8B). These results suggest that obese mice had inherited higher proportions of alpha cells, and lower proportions of endothelial and delta cells in their pancreatic islets.
- The loadings for pancreatic beta cell-type specific loadings was not significantly different from zero. This is not necessarily reflective of the function of the beta cells in the obese mice, but rather suggests that any variation in the number of beta cells in these mice was unrelated to obesity and insulin resistance.




Figure 8: HDM results translate to humans. **A.** Distribution of loadings for cell-type-specific transcripts in adipose tissue. **B.** Distribution of loadings for cell-type-specific transcripts in pancreatic islets (green). **C.** Null distributions for the mean loading of randomly selected transcripts in each cell type compared with the observed mean loading of each group of transcripts (red asterisk). **D.** Predictions of metabolic phenotypes in four adipose transcription data sets downloaded from GEO. In each study the obese/diabetic patients were predicted to have greater metabolic disease than the lean/non-diabetic patients based on the HDM results from DO mice.

233 Heritable transcriptomic signatures translated to human disease

234 Ultimately, the heritable transcriptomic signatures that we identified in DO mice will be useful if they inform
 235 pathogenicity and treatment of human disease. To investigate the potential for translation of the gene
 236 signatures identified in DO mice, we compared them to transcriptional profiles in obese and non-obese human
 237 subjects (Methods). We limited our analysis to adipose tissue because the adipose tissue signature had the
 238 strongest relationship to obesity and insulin resistance in the DO.

239 We calculated a predicted obesity score for each individual in the human studies based on their adipose
 240 tissue gene expression (Methods) and compared the predicted scores for obese and non-obese groups as well
 241 as diabetic and non-diabetic groups. In all cases, the predicted obesity scores were higher on average for
 242 individuals in the obese and diabetic groups compared with the lean and non-diabetic groups (Fig. 8D). This
 243 indicates that the heritable signature of obesity identified in DO mice is relevant to obesity and diabetes in
 244 human subjects.

245 **Targeting gene signatures**

246 Another global view of the transcript loading landscape is in ranking potential drug candidates for the
247 treatment of metabolic disease. Although high-loading transcripts may be good candidates for understanding
248 specific biology related to obesity, the transcriptome overall is highly interconnected and redundant, and
249 focusing on individual transcripts for treatment may be less effective than using a broader transcriptomic
250 signatures. The ConnectivityMap (CMAP) database [cite] developed by the Broad Institute allows us to
251 query thousands of compounds that reverse or enhance the extreme ends of transcriptomic signatures in
252 multiple different cell types. By identifying drugs that reverse pathogenic transcriptomic signatures, we can
253 potentially identify compounds that have favorable effects on gene expression.

254 To test this hypothesis we queried the CMAP database through the CLUE online query tool [cite] (Methods).
255 We identified top anti-correlated hits both across all cell types, as well as in adipocytes and pancreatic tumor
256 cells (Supplemental Figure XXX and XXX).

257 Looking broadly across cell types, the notable top hits from the adipose tissue loadings included mTOR
258 inhibitors and glucocorticoid agonists (Supplemental Figure XXX). It is thought that metformin, which is
259 commonly used to improve glycemic control, acts, at least in part, by inhibiting mTOR signaling [35, 36].
260 However, long-term use of other mTOR inhibitors, such as rapamycin, are known to cause insulin resistance and
261 β -cell toxicity [36, 37, 38]. Glucocorticoids are used to reduce inflammation, which was a prominent signature
262 in the adipose tissues, but these drugs also promote hyperglycemia and diabetes [39, 40]. Acute treatment
263 with glucocorticoids has further been shown to reduce thermogenesis in rodent adipocytes [41, 42, 43], but
264 increase thermogenesis in human adipocytes [44, 45]. Thus, the pathways identified by CMAP across all cell
265 types were highly related to the transcript loading profiles, but the relationship was not a simple reversal.

266 The top hit in adipocytes was a PARP inhibitor (Supplemental Figure XXXB). PARPs play a role in lipid
267 metabolism and are involved in the development of obesity and diabetes [46]. PARP1 inhibition increases
268 mitochondrial biogenesis [47]. Inhibition of PARP1 activity can further prevent necrosis in favor of the less
269 inflammatory apoptosis [48], thereby potentially reducing inflammation in stressed adipocytes. Other notable
270 hits in the top 20 were BTK inhibitors, which have been observed to suppress inflammation and improve
271 insulin resistance [49] as well as to reduce insulin antibodies in type I diabetes [50]. Similarly, IKK has been
272 shown to be associated with insulin resistance [51], and inhibitors have been shown to improve glucose control
273 in type II diabetes [52].

274 Among the top hits for the query with transcript loadings from pancreatic islets (Fig. XXX), was suppression
275 of T cell receptor signaling, which is known to be involved in Type 1 diabetes [53], as well as TNFR1, which

276 has been associated with mortality in diabetes patients [54]. Suppression of NOD1/2 signaling was also
277 among the top hits. NOD1 and 2 sense ER stress [55, 56], which is associated with β -cell death in type 1 and
278 type 2 diabetes [57]. This cell death process is dependent on NOD1/2 signaling [55], although the specifics
279 have not yet been worked out.

280 Among the top hits in pancreatic tumor cells were known diabetes drugs, including sulfonylureas, PPAR
281 receptor agonists, and insulin sensitizers. Rosiglitazone is a PPAR- γ agonist and was one of the most
282 prescribed drugs for type 2 diabetes before its use was reduced due to cardiac side-effects [58]. Sulfonylureas
283 are another commonly prescribed drug class for type 2 diabetes, but also have notable side effects including
284 hypoglycemia and accelerated β -cell death [59].

285 Discussion

286 It is thought that the bulk of the effect of genomic variation on complex traits is mediated through regulation
287 of gene expression. It has widely been assumed that this regulation is largely in *cis*, but attempts to use
288 local gene regulation to explain phenotypic variation have yet to explain much trait heritability. In recent
289 years, the discussion has turned to distal gene regulation. Although, distal gene regulation is more complex
290 to identify, evidence suggests that it is an important component of trait heritability.

291 Yao *et al.* [19] observed that genes with low local heritability explained more expression-mediated disease
292 heritability than genes with high local heritability. This observation is consistent with principles of robustness
293 in complex systems. If a transcript were both important to a trait and subject to strong local regulation,
294 a population would be susceptible to extremes in phenotype that might frequently cross the threshold to
295 disease. Indeed, strong disruption of highly trait-relevant genes is the cause of Mendelian disease.

296 Rather, observations suggest that genes that are near GWAS hits and have obvious functional relevance to a
297 trait tend to have highly complex regulatory landscapes under strong selection pressures [18]. In contrast,
298 genes with strong local regulation tend to be depleted of functional annotations and are under looser selection
299 constraints [18]. These observations and others led Liu *et al.* [60] to suggest that most heritability of complex
300 traits is driven by weak trans-eQTLs. They proposed a framework of understanding heritability of complex
301 traits in which massive polygenicity is distributed across common variants in both functional “core genes”, as
302 well as more peripheral genes that may not seem obviously related to the trait.

303 Here, we used a large, comprehensive, and purpose-built data set to investigate the roles of local and distal
304 gene regulation in mediating complex traits related to metabolic disease in mice. We propose a systems-level
305 method called high-dimension mediation (HDM). With this method we tested the hypothesis of the omnigenic

306 model directly, while allowing for the contributions of both local and distal gene regulation. The omnigenic
307 model posits that once the expression of the core genes (i.e. trait-mediating genes) is accounted for, there
308 should be no residual correlation between the genome and the phenotype. This hypothesis lends itself well to
309 systems approaches, such as HDM, that can account for arbitrarily complex gene regulation, as well as the
310 interconnectedness and redundancy of the transcriptome without explicitly modeling them.

311 Using HDM, we identified a highly heritable composite trait (71% heritable) that was perfectly mediated
312 by a composite transcript that included expression from four tissues known to be involved in metabolic
313 disease. Gene expression in adipose tissue was the strongest mediator of genetic effects on metabolic disease.
314 Further analysis of the loadings onto transcripts in each tissue revealed that the mediating signatures
315 were tissue-specific transcriptional programs many of which were previously known to be involved in the
316 pathogenesis of metabolic disease. We showed here that regulation of these programs is heritable and explains
317 a large proportion of disease risk.

318 We did not see evidence of distinct classes of core genes and peripheral genes. Rather, loadings on transcripts
319 were smoothly distributed across the whole transcriptome with genes more functionally related to metabolic
320 disease receiving larger loadings. The strength of mediation (transcript loading) was negatively correlated
321 with local heritability and positively correlated with distal heritability, suggesting that distal gene regulation
322 is the dominant mode through which gene expression mediates the effect of genotype on phenotype. This
323 pattern also suggests that distal regulation may be quite strong for many mediating genes. For example,
324 Nucb2, was a strongly mediating gene in islet and was also strongly distally regulated (66% distal heritability).
325 This gene is expressed in pancreatic β cells and is involved in insulin and glucagon release [61, 62, 63].
326 Although it was strongly distally regulated in islets, the regulation was distributed across the genome, with
327 no clear distal eQTL (Supplemental Figure XXX). Thus, although distal regulation of some genes may be
328 strong, this regulation is likely to be highly complex and not easily localized.

329 The high complexity of gene regulation combined with a systems-level analysis yields vague, continuous results
330 that do not implicate individual transcripts or genetic loci in disease pathogenesis. Historically, geneticists
331 have been interested in pinpointing individual loci whose mechanistic roles can be dissected through further
332 experiments. The continuous results, can also be quite informative, however, and paired with increasing
333 amounts of high-dimensional data in public data bases can be useful for generating hypotheses and potential
334 drug treatments. We showed that ranked lists of genes can be analyzed for enriched meaningful biological
335 functions and pathways using GSEA. These lists can be paired with information about cell-type specific genes
336 to generate hypotheses about cell composition in individual tissues. Finally, we used the CMAP database to
337 show that the transcriptomic signatures we identified in mice could be translated into human drug targets, as

338 currently used diabetes drugs were among the top hits for reversing the disease-associated signatures.

339 Discussion about biological relevance of high-loading transcripts in adipose tissue. TCA cycle and BCAA
340 catabolism: Selective PPAR γ modulation by insulin-sensitizing thiazolidinedione drugs has further been
341 shown to influence both inflammation and BCAA metabolism in obese rats suggesting a relationship between
342 these pathways and insulin resistance [64]. BCAA levels are also related to insulin resistance in human
343 subjects and are elevated in insulin-resistant obese individuals relative to weight-matched non-insulin resistant
344 individuals [65]. In the DO mice studied here, inheriting increased expression of genes involved in BCAA
345 catabolism was associated with reduced body weight and insulin resistance.

346 • distal heritability correlates with phenotype relevance

347 • others who use local eQTL to associate genotype with traits often say “we nominated this gene” even
348 though other nearby genes have higher eQTL LOD scores (27019110, 31465442) Our method supports
349 the idea that the transcripts with the strongest local regulation are less likely to be functionally related
350 to the trait

351 **Data Availability**

352 Here we tell people where to find the data

353 **Acknowledgements**

354 Here we thank people

355 **Supplemental Figures**




Figure 9: Overview of eQTL analysis in DO mice. **A.** RNA seq samples from the four different tissues clustered by tissue. **B.-E.** eQTL maps are shown for each tissue. The *x*-axis shows the position of the mapped eQTL, and the *y*-axis shows the physical position of the gene encoding each mapped transcript. Each dot represents an eQTL with a minimum LOD score of 8. The dots on the diagonal are locally regulated eQTL for which the mapped eQTL is at the within 4Mb of the encoding gene. Dots off the diagonal are distally regulated eQTL for which the mapped eQTL is distant from the gene encoding the transcript. **F.** Comparison of the total number of local and distal eQTL with a minimum LOD score of 8 in each tissue. All tissues have comparable numbers of eQTL. Local eQTL are much more numerous than distal eQTL. **G.** Counts of transcripts with local eQTL shared across multiple tissues. The majority of local eQTL were shared across all four tissues. **H.** Counts of transcripts with distal eQTL shared across multiple tissues. The majority of distal eQTL were tissue-specific and not shared across multiple tissues. For both G and H, eQTL for a given transcript were considered shared in two tissues if they were within 4Mb of each other. Colored bars indicate the counts for individual tissues for easy of visualization.

KEGG pathway enrichments by GSEA




Figure 10: Bar plots showing normalized enrichment scores (NES) for KEGG pathways as determined by fast gene score enrichment analysis (fgsea). Only the top 10 positive and top 10 negative scores are shown. Colors indicate tissue. The name beside each bar shows the name of each enriched KEGG pathway.

References

- 357 [1] M. T. Maurano, R. Humbert, E. Rynes, R. E. Thurman, E. Haugen, H. Wang, A. P. Reynolds,
 358 R. Sandstrom, H. Qu, J. Brody, A. Shafer, F. Neri, K. Lee, T. Kutyavin, S. Stehling-Sun, A. K.
 359 Johnson, T. K. Canfield, E. Giste, M. Diegel, D. Bates, R. S. Hansen, S. Neph, P. J. Sabo, S. Heimfeld,
 360 A. Raubitschek, S. Ziegler, C. Cotsapas, N. Sotoodehnia, I. Glass, S. R. Sunyaev, R. Kaul, and J. A.
 361 Stamatoyannopoulos. Systematic localization of common disease-associated variation in regulatory DNA.
 362 *Science*, 337(6099):1190–1195, Sep 2012.
- 363 [2] K. K. Farh, A. Marson, J. Zhu, M. Kleinewietfeld, W. J. Housley, S. Beik, N. Shores, H. Whitton, R. J.
 364 Ryan, A. A. Shishkin, M. Hatan, M. J. Carrasco-Alfonso, D. Mayer, C. J. Luckey, N. A. Patsopoulos,
 365 P. L. De Jager, V. K. Kuchroo, C. B. Epstein, M. J. Daly, D. A. Hafler, and B. E. Bernstein. Genetic

Top GO term enrichments by GSEA




Figure 11: Bar plots showing normalized enrichment scores (NES) for GO terms as determined by fast gene score enrichment analysis (fgsea). Only the top 10 positive and top 10 negative scores are shown. Colors indicate tissue. The name beside each bar shows the name of each enriched GO term. The letters in parentheses indicate whether the term is from the biological process ontology (BP), the molecular function ontology (MF), or the cellular compartment ontology (CC).

- 366 and epigenetic fine mapping of causal autoimmune disease variants. *Nature*, 518(7539):337–343, Feb
367 2015.
- 368 [3] E. Pennisi. The Biology of Genomes. Disease risk links to gene regulation. *Science*, 332(6033):1031, May
369 2011.
- 370 [4] L. A. Hindorff, P. Sethupathy, H. A. Junkins, E. M. Ramos, J. P. Mehta, F. S. Collins, and T. A. Manolio.
371 Potential etiologic and functional implications of genome-wide association loci for human diseases and
372 traits. *Proc Natl Acad Sci*, 106(23):9362–9367, Jun 2009.
- 373 [5] J. K. Pickrell. Joint analysis of functional genomic data and genome-wide association studies of 18
374 human traits. *Am J Hum Genet*, 94(4):559–573, Apr 2014.




Figure 12: Validation of transcript imputation in the CC-RIX. **A.** Distributions of correlations between imputed and measured transcripts in the CC-RIX. The mean of each distribution is shown by the red line. All distributions were skewed toward positive correlations and had positive means near a Pearson correlation (r) of 0.5. **B.** The relationship between the correlation between measured and imputed expression in the CC-RIX (x-axis) and eQTL LOD score. As expected, imputations are more accurate for transcripts with strong local eQTL. **C.** Variance explained by local genotype in the DO and CC-RIX.

- 375 [6] D. Welter, J. MacArthur, J. Morales, T. Burdett, P. Hall, H. Junkins, A. Klemm, P. Flicek, T. Manolio,
376 L. Hindorff, and H. Parkinson. The NHGRI GWAS Catalog, a curated resource of SNP-trait associations.
377 *Nucleic Acids Res*, 42(Database issue):D1001–1006, Jan 2014.
- 378 [7] Y. I. Li, B. van de Geijn, A. Raj, D. A. Knowles, A. A. Petti, D. Golan, Y. Gilad, and J. K. Pritchard.
379 RNA splicing is a primary link between genetic variation and disease. *Science*, 352(6285):600–604, Apr
380 2016.
- 381 [8] D. Zhou, Y. Jiang, X. Zhong, N. J. Cox, C. Liu, and E. R. Gamazon. A unified framework for joint-tissue
382 transcriptome-wide association and Mendelian randomization analysis. *Nat Genet*, 52(11):1239–1246,
383 Nov 2020.
- 384 [9] E. R. Gamazon, H. E. Wheeler, K. P. Shah, S. V. Mozaffari, K. Aquino-Michaels, R. J. Carroll, A. E.
385 Eyler, J. C. Denny, D. L. Nicolae, N. J. Cox, and H. K. Im. A gene-based association method for

- 386 mapping traits using reference transcriptome data. *Nat Genet*, 47(9):1091–1098, Sep 2015.
- 387 [10] Z. Zhu, F. Zhang, H. Hu, A. Bakshi, M. R. Robinson, J. E. Powell, G. W. Montgomery, M. E. Goddard,
388 N. R. Wray, P. M. Visscher, and J. Yang. Integration of summary data from GWAS and eQTL studies
389 predicts complex trait gene targets. *Nat Genet*, 48(5):481–487, May 2016.
- 390 [11] A. Gusev, A. Ko, H. Shi, G. Bhatia, W. Chung, B. W. Penninx, R. Jansen, E. J. de Geus, D. I. Boomsma,
391 F. A. Wright, P. F. Sullivan, E. Nikkola, M. Alvarez, M. Civelek, A. J. Lusis, T. ki, E. Raitoharju,
392 M. nen, I. ä, O. T. Raitakari, J. Kuusisto, M. Laakso, A. L. Price, P. Pajukanta, and B. Pasaniuc.
393 Integrative approaches for large-scale transcriptome-wide association studies. *Nat Genet*, 48(3):245–252,
394 Mar 2016.
- 395 [12] M. P. Keller, D. M. Gatti, K. L. Schueler, M. E. Rabaglia, D. S. Stapleton, P. Simecek, M. Vincent,
396 S. Allen, A. T. Broman, R. Bacher, C. Kendziora, K. W. Broman, B. S. Yandell, G. A. Churchill, and
397 A. D. Attie. Genetic Drivers of Pancreatic Islet Function. *Genetics*, 209(1):335–356, May 2018.
- 398 [13] W. L. Crouse, G. R. Keele, M. S. Gastonguay, G. A. Churchill, and W. Valdar. A Bayesian model
399 selection approach to mediation analysis. *PLoS Genet*, 18(5):e1010184, May 2022.
- 400 [14] J. M. Chick, S. C. Munger, P. Simecek, E. L. Huttlin, K. Choi, D. M. Gatti, N. Raghupathy, K. L. Svenson,
401 G. A. Churchill, and S. P. Gygi. Defining the consequences of genetic variation on a proteome-wide scale.
402 *Nature*, 534(7608):500–505, Jun 2016.
- 403 [15] H. E. Wheeler, S. Ploch, A. N. Barbeira, R. Bonazzola, A. Andaleon, A. Fotuhi Siahpirani, A. Saha,
404 A. Battle, S. Roy, and H. K. Im. Imputed gene associations identify replicable trans-acting genes enriched
405 in transcription pathways and complex traits. *Genet Epidemiol*, 43(6):596–608, Sep 2019.
- 406 [16] B. D. Umans, A. Battle, and Y. Gilad. Where Are the Disease-Associated eQTLs? *Trends Genet*,
407 37(2):109–124, Feb 2021.
- 408 [17] N. J. Connally, S. Nazeen, D. Lee, H. Shi, J. Stamatoyannopoulos, S. Chun, C. Cotsapas, C. A. Cassa,
409 and S. R. Sunyaev. The missing link between genetic association and regulatory function. *Elife*, 11, Dec
410 2022.
- 411 [18] H. Mostafavi, J. P. Spence, S. Naqvi, and J. K. Pritchard. Systematic differences in discovery of genetic
412 effects on gene expression and complex traits. *Nat Genet*, 55(11):1866–1875, Nov 2023.
- 413 [19] D. W. Yao, L. J. O’Connor, A. L. Price, and A. Gusev. Quantifying genetic effects on disease mediated
414 by assayed gene expression levels. *Nat Genet*, 52(6):626–633, Jun 2020.

- 415 [20] X. Liu, J. A. Mefford, A. Dahl, Y. He, M. Subramaniam, A. Battle, A. L. Price, and N. Zaitlen. GBAT:
416 a gene-based association test for robust detection of trans-gene regulation. *Genome Biol*, 21(1):211, Aug
417 2020.
- 418 [21] G. A. Churchill, D. M. Gatti, S. C. Munger, and K. L. Svenson. The Diversity Outbred mouse population.
419 *Mamm Genome*, 23(9-10):713–718, Oct 2012.
- 420 [22] S. M. Clee and A. D. Attie. The genetic landscape of type 2 diabetes in mice. *Endocr Rev*, 28(1):48–83,
421 Feb 2007.
- 422 [23] K. W. Broman, D. M. Gatti, P. Simecek, N. A. Furlotte, P. Prins, Š. Sen, B. S. Yandell, and G. A.
423 Churchill. R/qtL2: Software for Mapping Quantitative Trait Loci with High-Dimensional Data and
424 Multiparent Populations. *Genetics*, 211(2):495–502, Feb 2019.
- 425 [24] Fabien Girka, Etienne Camenen, Caroline Peltier, Arnaud Gloaguen, Vincent Guillemot, Laurent Le
426 Brusquet, and Arthur Tenenhaus. *RGCCA: Regularized and Sparse Generalized Canonical Correlation*
427 *Analysis for Multiblock Data*, 2023. R package version 3.0.3.
- 428 [25] Gennady Korotkevich, Vladimir Sukhov, and Alexey Sergushichev. Fast gene set enrichment analysis.
429 *bioRxiv*, 2019.
- 430 [26] A. Subramanian, P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert, M. A. Gillette, A. Paulovich,
431 S. L. Pomeroy, T. R. Golub, E. S. Lander, and J. P. Mesirov. Gene set enrichment analysis: a
432 knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*,
433 102(43):15545–15550, Oct 2005.
- 434 [27] C. B. Newgard. Interplay between lipids and branched-chain amino acids in development of insulin
435 resistance. *Cell Metab*, 15(5):606–614, May 2012.
- 436 [28] D. D. Sears, G. Hsiao, A. Hsiao, J. G. Yu, C. H. Courtney, J. M. Ofrecio, J. Chapman, and S. Subramaniam.
437 Mechanisms of human insulin resistance and thiazolidinedione-mediated insulin sensitization. *Proc Natl*
438 *Acad Sci U S A*, 106(44):18745–18750, Nov 2009.
- 439 [29] R. Stienstra, C. Duval, M. ller, and S. Kersten. PPARs, Obesity, and Inflammation. *PPAR Res*,
440 2007:95974, 2007.
- 441 [30] O. Gavrilova, M. Haluzik, K. Matsusue, J. J. Cutson, L. Johnson, K. R. Dietz, C. J. Nicol, C. Vinson,
442 F. J. Gonzalez, and M. L. Reitman. Liver peroxisome proliferator-activated receptor gamma contributes
443 to hepatic steatosis, triglyceride clearance, and regulation of body fat mass. *J Biol Chem*, 278(36):34268–
444 34276, Sep 2003.

- 445 [31] K. Matsusue, M. Haluzik, G. Lambert, S. H. Yim, O. Gavrilova, J. M. Ward, B. Brewer, M. L. Reitman,
446 and F. J. Gonzalez. Liver-specific disruption of PPARgamma in leptin-deficient mice improves fatty
447 liver but aggravates diabetic phenotypes. *J Clin Invest*, 111(5):737–747, Mar 2003.
- 448 [32] D. Patsouris, J. K. Reddy, M. Iller, and S. Kersten. Peroxisome proliferator-activated receptor alpha
449 mediates the effects of high-fat diet on hepatic gene expression. *Endocrinology*, 147(3):1508–1516, Mar
450 2006.
- 451 [33] S. E. Schadinger, N. L. Bucher, B. M. Schreiber, and S. R. Farmer. PPARgamma2 regulates lipogenesis
452 and lipid accumulation in steatotic hepatocytes. *Am J Physiol Endocrinol Metab*, 288(6):E1195–1205,
453 Jun 2005.
- 454 [34] W. Motomura, M. Inoue, T. Ohtake, N. Takahashi, M. Nagamine, S. Tanno, Y. Kohgo, and T. Okumura.
455 Up-regulation of ADRP in fatty liver in human and liver steatosis in mice fed with high fat diet. *Biochem
456 Biophys Res Commun*, 340(4):1111–1118, Feb 2006.
- 457 [35] S. Amin, A. Lux, and F. O’Callaghan. The journey of metformin from glycaemic control to mTOR
458 inhibition and the suppression of tumour growth. *Br J Clin Pharmacol*, 85(1):37–46, Jan 2019.
- 459 [36] A. Kezic, L. Popovic, and K. Lalic. mTOR Inhibitor Therapy and Metabolic Consequences: Where Do
460 We Stand? *Oxid Med Cell Longev*, 2018:2640342, 2018.
- 461 [37] A. D. Barlow, M. L. Nicholson, and T. P. Herbert. -cells and a review of the underlying molecular
462 mechanisms. *Diabetes*, 62(8):2674–2682, Aug 2013.
- 463 [38] Y. Gu, J. Lindner, A. Kumar, W. Yuan, and M. A. Magnuson. Rictor/mTORC2 is essential for
464 maintaining a balance between beta-cell proliferation and cell size. *Diabetes*, 60(3):827–837, Mar 2011.
- 465 [39] E. B. Geer, J. Islam, and C. Buettner. Mechanisms of glucocorticoid-induced insulin resistance: focus on
466 adipose tissue function and lipid metabolism. *Endocrinol Metab Clin North Am*, 43(1):75–102, Mar 2014.
- 467 [40] J. X. Li and C. L. Cummins. Fresh insights into glucocorticoid-induced diabetes mellitus and new
468 therapeutic directions. *Nat Rev Endocrinol*, 18(9):540–557, Sep 2022.
- 469 [41] R. A. Lee, C. A. Harris, and J. C. Wang. Glucocorticoid Receptor and Adipocyte Biology. *Nucl Receptor
470 Res*, 5, 2018.
- 471 [42] S. Viengchareun, P. Penfornis, M. C. Zennaro, and M. s. Mineralocorticoid and glucocorticoid receptors
472 inhibit UCP expression and function in brown adipocytes. *Am J Physiol Endocrinol Metab*, 280(4):E640–
473 649, Apr 2001.

- 474 [43] J. Liu, X. Kong, L. Wang, H. Qi, W. Di, X. Zhang, L. Wu, X. Chen, J. Yu, J. Zha, S. Lv, A. Zhang,
475 P. Cheng, M. Hu, Y. Li, J. Bi, Y. Li, F. Hu, Y. Zhong, Y. Xu, and G. Ding. -HSD1 in regulating brown
476 adipocyte function. *J Mol Endocrinol*, 50(1):103–113, Feb 2013.
- 477 [44] L. E. Ramage, M. Akyol, A. M. Fletcher, J. Forsythe, M. Nixon, R. N. Carter, E. J. van Beek,
478 N. M. Morton, B. R. Walker, and R. H. Stimson. Glucocorticoids Acutely Increase Brown Adipose
479 Tissue Activity in Humans, Revealing Species-Specific Differences in UCP-1 Regulation. *Cell Metab*,
480 24(1):130–141, Jul 2016.
- 481 [45] J. L. Barclay, H. Agada, C. Jang, M. Ward, N. Wetzig, and K. K. Ho. Effects of glucocorticoids on
482 human brown adipocytes. *J Endocrinol*, 224(2):139–147, Feb 2015.
- 483 [46] M. ó, R. Gupte, W. L. Kraus, P. Pacher, and P. Bai. PARPs in lipid metabolism and related diseases.
484 *Prog Lipid Res*, 84:101117, Nov 2021.
- 485 [47] P. Bai, C. ó, H. Oudart, A. nszki, Y. Cen, C. Thomas, H. Yamamoto, A. Huber, B. Kiss, R. H.
486 Houtkooper, K. Schoonjans, V. Schreiber, A. A. Sauve, J. Menissier-de Murcia, and J. Auwerx. PARP-1
487 inhibition increases mitochondrial metabolism through SIRT1 activation. *Cell Metab*, 13(4):461–468,
488 Apr 2011.
- 489 [48] A. Chiarugi and M. A. Moskowitz. Cell biology. PARP-1—a perpetrator of apoptotic cell death? *Science*,
490 297(5579):200–201, Jul 2002.
- 491 [49] M. Althubiti, R. Almaimani, S. Y. Eid, M. Elzubaier, B. Refaat, S. Idris, T. A. Alqurashi, and M. Z.
492 El-Readi. BTK targeting suppresses inflammatory genes and ameliorates insulin resistance. *Eur Cytokine
493 Netw*, 31(4):168–179, Dec 2020.
- 494 [50] C. Skrabs, W. F. Pickl, T. Perkmann, U. ger, and A. Gessl. Rapid decline in insulin antibodies and
495 glutamic acid decarboxylase autoantibodies with ibrutinib therapy of chronic lymphocytic leukaemia. *J
496 Clin Pharm Ther*, 43(1):145–149, Feb 2018.
- 497 [51] M. C. Arkan, A. L. Hevener, F. R. Greten, S. Maeda, Z. W. Li, J. M. Long, A. Wynshaw-Boris, G. Poli,
498 J. Olefsky, and M. Karin. IKK-beta links inflammation to obesity-induced insulin resistance. *Nat Med*,
499 11(2):191–198, Feb 2005.
- 500 [52] E. A. Oral, S. M. Reilly, A. V. Gomez, R. Meral, L. Butz, N. Ajluni, T. L. Chenevert, E. Korytnaya,
501 A. H. Neidert, R. Hench, D. Rus, J. F. Horowitz, B. Poirier, P. Zhao, K. Lehmann, M. Jain, R. Yu,
502 C. Liddle, M. Ahmadian, M. Downes, R. M. Evans, and A. R. Saltiel. and TBK1 Improves Glucose
503 Control in a Subset of Patients with Type 2 Diabetes. *Cell Metab*, 26(1):157–170, Jul 2017.

- 504 [53] M. Clark, C. J. Kroger, Q. Ke, and R. M. Tisch. The Role of T Cell Receptor Signaling in the
505 Development of Type 1 Diabetes. *Front Immunol*, 11:615371, 2020.
- 506 [54] P. ndell, A. C. Carlsson, A. Larsson, O. Melander, T. Wessman, J. v, and T. Ruge. TNFR1 is associated
507 with short-term mortality in patients with diabetes and acute dyspnea seeking care at the emergency
508 department. *Acta Diabetol*, 57(10):1145–1150, Oct 2020.
- 509 [55] A. M. Keestra-Gounder, M. X. Byndloss, N. Seyffert, B. M. Young, A. vez Arroyo, A. Y. Tsai, S. A.
510 Cevallos, M. G. Winter, O. H. Pham, C. R. Tiffany, M. F. de Jong, T. Kerrinnes, R. Ravindran, P. A.
511 Luciw, S. J. McSorley, A. J. umler, and R. M. Tsolis. NOD1 and NOD2 signalling links ER stress with
512 inflammation. *Nature*, 532(7599):394–397, Apr 2016.
- 513 [56] A. M. Keestra-Gounder and R. M. Tsolis. NOD1 and NOD2: Beyond Peptidoglycan Sensing. *Trends
514 Immunol*, 38(10):758–767, Oct 2017.
- 515 [57] J. Montane, L. Cadavez, and A. Novials. Stress and the inflammatory process: a major cause of
516 pancreatic cell death in type 2 diabetes. *Diabetes Metab Syndr Obes*, 7:25–34, 2014.
- 517 [58] B. B. Kahn and T. E. McGraw. , and type 2 diabetes. *N Engl J Med*, 363(27):2667–2669, Dec 2010.
- 518 [59] S. Del Prato and N. Pulizzi. The place of sulfonylureas in the therapy for type 2 diabetes mellitus.
519 *Metabolism*, 55(5 Suppl 1):S20–27, May 2006.
- 520 [60] X. Liu, Y. I. Li, and J. K. Pritchard. Trans Effects on Gene Expression Can Drive Omnipotent Inheritance.
521 *Cell*, 177(4):1022–1034, May 2019.
- 522 [61] M. Riva, M. D. Nitert, U. Voss, R. Sathanoori, A. Lindqvist, C. Ling, and N. Wierup. Nesfatin-1
523 stimulates glucagon and insulin secretion and beta cell NUCB2 is reduced in human type 2 diabetic
524 subjects. *Cell Tissue Res*, 346(3):393–405, Dec 2011.
- 525 [62] M. Nakata and T. Yada. Role of NUCB2/nesfatin-1 in glucose control: diverse functions in islets,
526 adipocytes and brain. *Curr Pharm Des*, 19(39):6960–6965, 2013.
- 527 [63] H. Shimizu and A. Osaki. Nesfatin/Nucleobindin-2 (NUCB2) and Glucose Homeostasis. *Curr Hypertens
528 Rev*, pages Nesfatin/Nucleobindin-2 (NUCB2) and Glucose Homeostasis., Jul 2014.
- 529 [64] G. Hsiao, J. Chapman, J. M. Ofrecio, J. Wilkes, J. L. Resnik, D. Thapar, S. Subramaniam, and D. D.
530 Sears. modulation of insulin sensitivity and metabolic pathways in obese rats. *Am J Physiol Endocrinol
531 Metab*, 300(1):E164–174, Jan 2011.
- 532 [65] D. E. Lackey, C. J. Lynch, K. C. Olson, R. Mostaedi, M. Ali, W. H. Smith, F. Karpe, S. Humphreys,

533 D. H. Bedinger, T. N. Dunn, A. P. Thomas, P. J. Oort, D. A. Kieffer, R. Amin, A. Bettaieb, F. G.
534 Haj, P. Permana, T. G. Anthony, and S. H. Adams. Regulation of adipose branched-chain amino acid
535 catabolism enzyme expression and cross-adipose amino acid flux in human obesity. *Am J Physiol*
536 *Endocrinol Metab*, 304(11):E1175–1187, Jun 2013.