

PROJECT NARRATIVE

A. Background and Rationale

The **goal** of this proposal is to map genetic modifiers of tuberous sclerosis complex (TSC) outcomes in a genetically diverse mouse population (Fig. 1). TSC is a rare autosomal dominant disease caused by loss-of-function (LoF) mutations in either of the genes *TSC1* or *TSC2* (*TSC1/2*). These mutations cause aberrant up-regulation of mammalian target of rapamycin (mTOR) signaling, resulting in high rates of epilepsy, autism spectrum disorder (ASD), and tumor formation in multiple organ systems[?]. However, presentation of TSC in the patient population is highly heterogeneous ranging from subclinical to severe^{??????}. There are at least three distinct contributors to this phenotypic complexity: 1) a patient’s specific TSC gene mutation; 2) the random occurrence of somatic second-hit *TSC1/2* mutations, and 3) genetic modifiers, i.e., alleles inherited separately from the *TSC1/2* genes that alter the severity of TSC-associated phenotypes. The **central hypothesis** of this project is that genetic modifiers are a major source of patient heterogeneity. Despite high heterogeneity within affected families sharing the same mutation^{??}, TSC outcomes are more similar in identical twins and within families than across families[?], supporting the existence of genetic modifiers in the human population.




Figure 1: Project overview. Breeding schematics show generation of A) DO-*Tsc1* and B) DO-*Tsc2* mice by crossing a DO female to an inbred mouse carrying a *Tsc1* or *Tsc2* allele. DO-*Tsc1* and DO-*Tsc2* mice will be used in both aims of this proposal. Both mouse subpopulations will have brain, kidneys, lung, liver, and heart harvested and will be run through all phenotyping pipelines. RNA sequencing will be performed in brain for the DO-*Tsc1* subpopulation and kidney for the DO-*Tsc2* subpopulation. We will use AI technology to quantify C) seizures and B) histological phenotypes in all mice. We will map TSC-related phenotypic outcomes as E) individual QTL and F) as genetic interaction networks. G) Gene expression will be used as a mediator of genetic effects on phenotypes to identify molecular pathways influencing tumor and neurological outcomes.

The **rationale** for this study is that by mapping genetic modifiers, we will identify novel genes and pathways that can be therapeutically targeted to mitigate poor outcomes. Although current therapies that directly target mTOR signaling show promise in treating TSC-associated conditions^{?????}, significant challenges remain. Indeed, some TSC tumors are unaffected by treatment with rapamycin^{????}, and both seizures[?] and tumors^{??} develop resistance to these therapies with long-term treatment. Thus, the identification of alternative or adjunct therapies for TSC is critical for long-term disease management. Previous work supports strategies targeting mTOR-independent pathways. For example, genetic studies in patients with TSC led to work in a mouse model demonstrating that treatment with IFN- γ in combination with rapamycin was more effective in reducing tumor growth and improving survival more than rapamycin alone[?]. Identification of this and other alternative pathways^{???} through genetic modifier screens, may be critical for treatment of refractory TSC.

Our preliminary data and prior reports definitively demonstrate the existence of genetic modifiers of both neurological and tumor outcomes in haploinsufficiency mouse models of TSC, providing a tractable experimental platform for genetic discovery. Here, we propose to use natural genetic variation in a population-based model of TSC to identify additional molecular pathways that drive variation in TSC outcomes.

Genetic modifiers of seizure outcomes exist in a haploinsufficiency mouse model of TSC. In **Aim 1** we will genetically map modifiers of neurological outcomes in a mouse population model of *Tsc1/2* haploinsufficiency. Our recently completed study (DOD TS180087) tested whether epilepsy outcomes in *Tsc1*^{+/-} mice were influenced by genetic background (Fig. ??). We crossed a C57BL/6J-*Tsc1*^{+/-} mouse to multiple wild-type genetic backgrounds—C57BL/6J (B6), DBA/2J (D2), and BXD87/Rww (BXD87)—allowing us to test the relationship between strain background and disease outcomes (Fig. ??A). The BXD87 strain is a recombinant inbred strain whose genome is a mosaic of the B6 and D2 genomes (Fig. ??A), which we chose because wild-type BXD87 mice are outliers for both low hippocampal volume and poor cognitive performance^{??}, suggesting a developmentally susceptible state to poor TSC outcomes. Because the carrier parent was heterozygous for the *Tsc1* knockout, the F1 progeny could inherit either one knockout allele (HET) or the B6 wild-type allele (WT) (Fig. ??A). We performed five-day video-electroencephalogram (EEG) monitoring (starting around postnatal day 70) in N = 48 mice (Fig. ??B). We found that only BXD87-HET mice developed spontaneous, tonic-clonic seizures when carrying a single copy of the *Tsc1* null allele. Over the course of our experiment 4 out of 7 BXD87 females had a total of 61 seizures (Fig. ??C-D), averaging 1.7 seizures per BXD87 female mouse per day. This is the first time that a mouse model has been observed to develop chronic epilepsy without an induced second hit to *Tsc1/2*, demonstrating that a single germline mutation of *Tsc1* can cause chronic seizures in mice in a manner that is modified by genetic background. We note that haploinsufficiency models of spontaneous seizures have been reported before; however, in these models early-life epilepsy resolved by post-natal day 19[?], well before adulthood, and hence does not recapitulate the chronic epilepsy of the most severely affected human patients.

We can make two important observations about the genetics of epilepsy in our BXD87-HET model. First, the appearance of seizures in BXD87, but not the parent strains B6 and D2, implies that there must be at least two genetic modifiers in BXD87 responsible for epilepsy; If there were only one, we would have observed seizures in either B6-HET or D2-HET mice. Second, we observed seizures only in around half of the female BXD87-HET mice. Based on our cross design with a B6-*Tsc1*^{+/-} dam (Fig. ??A), we infer that one of the modifier alleles may be on the X chromosome and that incomplete penetrance may be due to random X inactivation^{??}. We are currently testing this hypothesis, which is out of the scope of this project. However, to maximize our chance of detecting an X-linked modifier in this study, we will reverse the cross so that the sire is the inbred carrier and the dam is the genetically diverse, non-carrier. In this design, progeny of all sexes will have genetic variation on the X chromosome.

Genetic modifiers of tumor outcomes exist in haploinsufficiency mouse models of TSC. In **Aim 2** we will genetically map modifiers of tumor outcomes in a population model of *Tsc1/2* haploinsufficiency. Unlike epilepsy, there is substantial prior work in rodents showing that genetic variation is an important driver of tumor outcomes in *Tsc1/2* haploinsufficiency[?]. In Eker rats, which carry a null allele of *Tsc2*, variation in kidney tumor size and number have been mapped to alleles on rat chromosomes 3[?] and 5[?]. In mice carrying a null *Tsc2* allele, tumor number, extent, and location vary by strain background. Specifically, A/J-*Tsc2*^{+/-} mice had a substantially higher burden of kidney tumors than 129S4/SvJae-*Tsc2*^{+/-} mice[?], while 129S4/SvJae-*Tsc2*^{+/-} mice had a higher burden of liver tumors compared to six other strains tested[?]. Similarly, a null allele of *Tsc1* caused high incidence of renal carcinoma with some lung metastases in BALB/c mice compared to other strains, while the same allele caused early-life mortality in one quarter of B6 pups[?]. Together these studies demonstrate that tumor outcomes in haploinsufficiency models of TSC are highly heritable.

Rationale for a DO-TSC population model. The heritability of TSC outcomes in inbred strains proves the existence of modifier alleles, but does not identify them. Here we propose to map these alleles using the Diversity Outbred (DO) mouse population. The DO mice are a highly recombined, outbred population derived from eight inbred founder strains[?], which capture 89% of the known genetic variation in laboratory mice[?]. Three of the DO founders—

with highly mature software tools for genetic mapping[?] (Fig. 1E), gene network inference[?] (Fig. 1F), and gene expression mediation analysis[?] (Fig. 1G). We have successfully used DO mice to map individual QTLs and infer genetic networks[?] (Fig. 1E-F). Others have successfully mapped tumor outcomes using modifier screens similar to this proposal[?]. We have also thoroughly studied the effects of population structure on mapping studies in this population[?]. This prior work combined with our strong preliminary data demonstrate that the DO-TSC population will be a singularly powerful genetic discovery resource for TSC.

Artificial intelligence (AI) approaches for scalable, automated phenotyping. One of the persistent problems for genetic mapping has been the requirement for high-throughput phenotyping at the scale of hundreds of individuals. This has traditionally meant either simple summaries of complex observations (e.g., case vs. control) or surrogate biomarkers that may not capture the complete phenotype. However, recent advances in AI and machine learning have dramatically lowered the barriers to subtle phenotyping at scale, including scoring days-long home-cage behavior videos[?], non-invasive seizure monitoring[?], and analyzing whole-slide images of tissues[?] (Fig. 1C-D). We have pioneered applications of AI-based image analysis for quantitative trait genetics. In a study of the *Far2* gene, which alters kidney aging phenotypes[?], we demonstrated that AI approaches could recapitulate known kidney phenotypes and detect novel phenotypes using only genotype as a training label[?]. In an extension of this work, Dr. Mahoney has an active R01 on histological imaging genetics to further develop these tools. Taken together, this prior work and our feasibility data (described below) demonstrate that we can use AI models to robustly score seizure, behavioral, and histopathological phenotypes at scale, and genetically map AI outputs as biologically meaningful traits. Thus, our innovative AI-based phenotyping pipeline overcomes one of the primary technical limitations of genetic mapping.

B. Hypothesis and Objective The **objective** of this project is to identify novel genetic modifiers of TSC outcomes and identify the gene expression signatures that mediate these effects. To achieve this, we will perform a systems genetics analysis of a novel DO-TSC population model. We hypothesize that: 1) DO-TSC mice will vary significantly in seizure, behavioral, and tumor outcomes; 2) each outcome will have modifiers mapping to specific locations in the genome, and 3) we will identify significant modifier pathways using gene-expression mediation analysis.

C. Specific Aims

Aim 1: Identify genetic modifiers of epilepsy and TANDs in TSC by: a) mapping genetic modifiers of epilepsy and TANDs in our DO-TSC model and b) performing mediation analysis with brain gene expression to identify the causal pathways driving poor neurological outcomes.

Aim 2: Identify genetic modifiers of tumors in TSC by: a) mapping genetic modifiers of tumor outcomes in brain, kidney, and lung in our DO-TSC model and b) performing mediation analysis with kidney gene expression to identify the causal pathways driving poor tumor outcomes.

D. Research Strategy and Feasibility

Aim 1: Identify genetic modifiers of epilepsy and TANDs in TSC

Experimental approach:

Generation of experimental mice: We will generate two subpopulations of DO-TSC mice, which we designate as DO-*Tsc1* and DO-*Tsc2* (Fig. 1A-B). We will phenotype both sub-populations in both aims (Fig. 1C-D). Each mouse will be an F1 hybrid between a genetically unique female DO mouse and a male inbred carrier mouse. Each F1 hybrid will also be genetically unique, because of recombination of the DO dam's chromosomes during meiosis. For the DO-*Tsc1* subpopulation, we will use B6-*Tsc1*^{+/-} mice as the carrier sire. We will generate these mice as in our preliminary data, using a cross between our B6-*Tsc1*/fl/fl mouse line and the CMV-Cre line, which expresses Cre recombinase at the germline, causing germline heterozygosity of *Tsc1* LoF. For the DO-*Tsc2* subpopulation, we will use A/J-*Tsc2*^{+/-} mice as the carrier sire. These mice are not commercially available, so the JAX Genetic Engineering Technologies (GET) group will generate this line de novo using CRISPR/Cas9 to knock out exons

5 through 8 in the A/J background, matching the construct used in prior reports^{??}. In both subpopulations, we will genotype all pups for the TSC mutation before weaning using standard PCR and retain pups with a *Tsc1/2* mutation for phenotyping. We will also maintain detailed records of litter sizes and *Tsc1/2* mutation frequency as an additional phenotype. TSC mutations can cause fetal death in humans^{??}, and prenatal death would result in a significant deviation from the expected 50% frequency of *Tsc1/2* mutations among pups.

Seizure and behavior monitoring. For high-throughput seizure and behavioral phenotyping, we will use a novel home-cage behavior system developed by TLR Ventures. This home-cage behavior system was developed in collaboration with the Digital In Vivo Alliance (DIVA), a consortium that includes JAX and supports the development of automated home-cage technologies. DIVA cages have two cameras, one overhead and another at a side angle, that stream video directly to cloud-based computing centers. Once data are in the cloud, DIVA cage software automatically applies AI models to identify animal position (segmentation), identify key points on the body (key point detection), predict respiratory rate, and identify loss of righting reflex (LORR) during seizures. These data are stored for further processing in a cloud-based data science environment, and user-interface tools allow researchers to review videos, define segments of interest, and train models *de novo*. Starting around postnatal day 70, we will put up to three littermate DO-TSC mice into the DIVA cages for three consecutive days of continuous monitoring to measure the frequency and duration of seizures, as primary outcomes. We will also measure frequencies and durations of abnormal behaviors, such as backflipping, abnormal grooming, or other stereotypies that have been associated with ASD, as secondary outcomes. We will manually review periods of high predicted probability of LORR to confirm seizures and fine-tune the seizure detection model using manual annotations. Similarly, we will apply and fine-tune existing models to detect periods of highly stereotyped behavior, abnormal grooming activity, and identify backflipping^{??????}.

Tissue collection: At 14 months of age, we will collect brain, kidney, lung, liver, and heart tissue from each DO-TSC mouse. All tissues will be fixed with formalin and embedded in paraffin (FFPE) for long-term storage. We will separate brains along the corpus callosum and use one hemisphere for RNA sequencing and the other for histology.

RNA sequencing: To identify the gene expression signatures that mediate the effects of genetic modifiers on epilepsy and TANDs, we will perform bulk gene expression of brain tissue. Gene expression analysis of all mouse brains from this study is beyond the scope of this project. Thus, we will focus on DO-*Tsc1*, which has a high probability of manifesting heterogeneous neurological outcomes. We will deliver DO-*Tsc1* brain samples to the JAX Genome Technologies Service for 50 base pair, paired-end bulk RNA sequencing. Fastq files will be delivered to our group for analysis. We will perform read alignment and gene expression quantification using custom, publicly available RNA-seq pipelines developed by JAX Computational Sciences[?]. This pipeline performs read trimming to remove sequencing barcodes, quantifies gene expression using the RNA-Seq by Expectation Maximization (RSEM) algorithm[?], and delivers sample- and batch-level summary and quality-control statistics, including Picard alignment metrics[?] and a MultiQC[?] report. Gene expression levels will be normalized using a variance-stabilizing transformation^{??} for downstream processing.

Genotyping and reconstruction of ancestral haplotypes: For background genotyping, we will use miniMUGA SNP genotyping arrays[?] to generate SNP-level genotypes for each mouse. We will then compute the DO founder haplotypes using HaploQA[?]. We will use haplotypes for genetic mapping[?].

Genetic mapping and power analysis: We will perform two complementary forms of genetic mapping—linkage mapping and combined analysis of pleiotropy and epistasis (CAPE)—each of which has unique strengths. In linkage mapping, we correlate the haplotypes at individual genetic markers with variation in quantitative traits, such as seizure frequency. Linkage mapping seeks loci with strong marginal effects, i.e., strong effects across many genetic backgrounds. We will perform linkage mapping using R/qt12[?] and define statistical significance for QTLs using standard permutation-based statistical thresholds[?]. We will use sex and TSC-genotype as interacting covariates to identify potential sex- and genotype-specific modifiers. We will also use linear mixed models to account for the effects of variable relatedness among individuals[?]. Because the DO mice are highly recombined, we expect

that mapping in this population will yield QTLs encompassing small genomic regions (< 1 megabase (Mb)) with a small number of positional candidate genes. With $n = 400$ mice total, we will have 80% power to detect loci explaining 10% of the variance in any individual phenotype[?]. This population size conforms to prior successful mapping studies in DO-F1 hybrid mice[?]. Because linkage mapping is the most statistically stringent form of mapping, we have powered our study for this analysis. Using gene expression from RNA sequencing, we will also perform expression QTL (eQTL) analysis to identify genetic loci regulating gene expression levels. We will map eQTLs exactly as described above, once for each transcript. The effect sizes for eQTLs are generally much more significant than QTLs for clinical traits due to strong cis-regulatory effects. Thus, with $n = 200$ mice with gene expression, we expect large numbers of eQTLs. By identifying loci that influence both gene expression and traits, we can more easily link trait variation to molecular mechanisms[?]. In addition to mapping to individual QTLs, we will use CAPE, developed by Dr. Tyler[?], to map trait variation to epistatically interacting QTLs (genetic network inference). No gene acts in isolation, and many genetic modifiers of TSC may depend on interactions with the rest of the genome. CAPE leverages information across multiple traits to infer directed epistatic interactions affecting all traits simultaneously. The simultaneous analysis of multiple traits increases power to detect QTLs over classical linkage mapping[?]. Moreover, CAPE allows inference of directed interactions indicating which variant acts upstream of another variant in an interacting pair, thereby aiding in interpretation of the interaction and prioritization of causal positional candidate genes^{??}. Furthermore, the interactions identified are consistent across all analyzed traits, allowing us to relate clinical traits to genetic interactions through endophenotypes, such as the expression of genes in the mTOR signaling pathway, which we will include as traits in the analysis^{???}. As with linkage mapping, we will use sex and TSC-genotype as interactive covariates to ascertain sex- and genotype-specific effects. We will also use linear mixed models to account for interindividual relatedness[?] and perform permutation testing to assess interaction significance[?]. We have experience using CAPE to analyze seizure outcomes in mouse models of absence epilepsy[?], and complex traits in a DO population[?] comparable to the one proposed here. We expect that in this population we will be able to generate multiple testable molecular hypotheses about genetic variants that influence multiple clinical outcomes simultaneously.

Transcriptomic mediation analysis: One limitation to all forms of genetic mapping is that the effects of individual loci must be large enough to detect statistically. However, there is now extensive evidence that most complex traits are highly polygenic, and this polygenic effect is mediated through changes in transcriptional profiles^{??}. Thus, as a complement to genetic mapping per se, we will also use gene expression to identify the transcriptomic profiles driving heritable variation in neurological outcomes, independent of individual genomic loci. To do this, we will perform high-dimensional mediation analysis[?], which is an extension of canonical correlation analysis that, in our case, seeks a transcriptomic signature that is simultaneously correlated with underlying genetics and phenotypic outcomes. Formally, if G , T , and P , represent our genotype, transcriptome, and phenotype data matrices, we define composite genetic, transcriptomic, and phenotypic scores, $s_g = G \cdot l_g$, $s_t = T \cdot l_t$, and $s_p = P \cdot l_p$, respectively, through matrix multiplication of the data matrices with loading vectors l_g , l_t , and l_p that define each variable's importance to the corresponding score. Using the theory of causal graphical models^{??}, we maximize the likelihood function $L = -\log |S| - \text{tr}(\Sigma S^{-1})$, where Σ is the correlation matrix of the scores (s_g , s_t , and s_p), S is the model-implied covariance matrix for the perfect mediation model, and $|\cdot|$ and tr denote the determinant and trace of a matrix, respectively[?]. This model encodes the causal flow of information from genotype, through transcription, to phenotype. The outputs of this analysis are the optimal scores and loadings that define the causal influence of genetic background on TSC outcomes. We will rigorously define the significance of mediation by permuting the transcriptomes relative to the genotypes and phenotypes to generate a null distribution for the path coefficient from genotype to phenotype. To identify the significant pathways in the transcriptomic score, we will rank all individual transcripts by their loadings, l_t , and perform gene set enrichment analysis (GSEA)[?]. In this way, the transcript rankings are a powerful complement to genetic mapping because they allow inference of the causal importance of genes and pathways, even when background modifier effects are highly polygenic.

Feasibility: As feasibility data for this proposal, we trained an AI model *de novo* to detect LORR in manually coded frames from videos of 25 mice during pentylenetetrazole-induced seizures (Fig. ??A). The model achieved greater than 95% accuracy detecting LORR in testing data (Fig. ??B). As an example readout, the model computes

a time-varying probability (Fig. ??C, blue trace) that predicts when the mouse has fallen vs. is upright. Time intervals with high predicted probability were strongly correlated with expert annotations of LORR (Fig. ??C, pink boxes). These data demonstrate that the DIVA cages can robustly detect and quantify seizures at scale. The DIVA cages also compute animal segmentations and pose-estimation phenotypes from which we can score standard home cage behaviors, such as grooming.




Figure 3: AI classification of seizures. A) Still images of normal mice (top row) and mice experiencing seizures (bottom row). The bottom row shows the loss-of-righting reflex (LORR) that is classified by the AI as a seizure. B) Confusion matrix showing the counts of correct and incorrect classifications by the AI along with calculated accuracy, recall, and precision. C) A time-varying probability of LORR predicted by the AI at each frame of a video (blue trace). High probabilities indicate frames for which the AI predicts that the mouse has fallen over. Pink areas highlight frames for which an expert reviewer has indicated that the mouse has fallen over.

Potential problems and alternative strategies: We anticipate three potential issues. First, we may not detect any QTLs at the genome-wide significance level using any of the proposed strategies, thereby limiting candidate gene prioritization from genetic mapping. While we stress that our combined approach is designed to mitigate this possibility, if it occurs, we anticipate that it will be due to both the presence of a large number of modifiers and limited statistical power to detect effects. If this transpires, we will use false discovery rate (FDR) approaches[?] to lower the burden of multiple testing for our associations. We also note that high-dimensional mediation analysis is an explicitly polygenic analysis that supports gene prioritization independent of QTLs. Second, we may not have sufficient genetic resolution to nominate candidate genes by mapping alone. While current DO generations have linkage blocks that are substantially below 1 Mb, it is possible that our analyses may not resolve individual candidate genes. If this transpires, we will pursue candidate gene prioritization within QTLs using network-based approaches that Drs. Mahoney and Tyler have extensively developed^{??}. Third, we may find that DO-TSC mice have heterogeneous seizure presentations that are difficult to track with our existing AI tools. Because we expect all animals having tonic-clonic seizures to have LORR, we anticipate that thresholding the AI-generated probabilities at a low level, e.g. 20%, will guide us to epochs that are enriched for seizure activity (see Fig. ??). If we identify misclassified epochs, we will first attempt to fine-tune the LORR model by manually annotating a subset of these epochs as ground-truth labels. Barring that, following Geschwind et al.[?], we will perform unsupervised analyses that cluster animal behaviors into groups, independent of annotations, and manually annotate clusters corresponding to seizures and other abnormal behaviors. In the worst case, we will review all videos in their entirety, visually

screening for seizures.

Aim 2: Identify genetic modifiers of tumors in TSC.

Experimental approach: For this aim we will use the same 400 DO-TSC mice developed in **Aim 1**. We will perform tissue pre-processing, genotyping, genetic mapping, and mediation analysis exactly as described above.

Histological imaging: We will deliver FFPE brain, kidney, and lung tissue blocks to the JAX Histology Service for sectioning and staining with hematoxylin and eosin (H&E). Stained slides will then be delivered to the JAX Microscopy Service for slide scanning with a 40X objective. Whole-slide image files will be delivered to our team for analysis. At the conclusion of this study, we will make all histological images available to the public.

Image processing and genetic mapping: To generate histological phenotypes at the scale required for genetic mapping, we will use an AI-based computer vision approach to automatically score the extent of tissue abnormalities without requiring extensive human annotation. From each tissue image, we will extract tiles, or sub-images, of 299x299 pixels, which we will input into the Inception v3 deep neural network[?]. The Inception v3 model is a top-performing network for natural image classification[?]. We will perform transfer learning[?], where we remove the last layers of the model and retain the numerical features at the second-to-last layer as a black-box feature vector encoding histological image structure. This transfer learning approach has been applied heavily in the cancer field with remarkable success[?]. From the black-box feature vector, we will perform principal components analysis (PCA) to identify the dominant axes of variation across all images of the same tissue. For each sample, we will average the PCA scores over tiles to compute mouse-level scores, which we will map as quantitative traits, using the techniques described in **Aim 1**. We will visualize the learned tissue features from our model using example montages, where we compare high- and low-scoring tiles to detect differences.

RNA sequencing and analysis: To identify the gene expression signatures that mediate the effects of genetic modifiers on tumors, we will perform bulk gene expression of kidney tissue. Gene expression analysis of all mouse tissues from this study is beyond the scope of this project. Thus, we will focus on DO-*Tsc2* kidneys, which have a high probability of manifesting heterogeneous tumor outcomes.

Tissue biobanking: We will biobank the brain, lung, and kidney tissue not used for histology or RNA sequencing, as well as all liver and heart samples, from each DO-TSC mouse. We expect our results from this project to provide a strong rationale to generate comprehensive histological and gene expression data for these remaining samples. We will also make the biobanked tissues available to the wider TSC research community upon reasonable request.

Feasibility: In ongoing work for Dr. Mahoney's R01, we have analyzed kidney aging in a population of 500 DO mice using the strategy described above (in preparation; Fig. ??). Using whole-slide images of periodic acid-Schiff-stained kidney sections, we extracted features from ~150k individual glomeruli and performed PCA to automatically identify age-related tissue features across the population. We genetically mapped these AI-derived scores and identified a significant QTL on chr. 17 (Fig. ??A). Examples of high- and low-scoring glomeruli show marked differences in mesangial matrix expansion, an age-associated histopathology (Fig. ??B), and cellularity, both of which we have validated quantitatively (data not shown). The allele effects for this AI-derived trait show that inheriting the A/J allele leads to worse outcomes while the NZO/LtJ allele leads to better outcomes (Fig. ??C). We have nominated high-confidence candidate genes within this locus—*Xdh* and *Birc6*—which have variants specific to the A/J and NZO/LtJ backgrounds, respectively, and strong evidence for association to kidney phenotypes[?]. We have ongoing experiments with CRISPR knock-in mice to validate these effects. While this study does not speak to the hypotheses of this project, it demonstrates feasibility for mapping genetic modifiers of histological outcomes using AI-derived traits.

Potential problems and alternative strategies: We can anticipate two potential problems. First, bulk RNA sequencing of kidney tissues necessarily mixes healthy tissue and tumor tissue, possibly limiting our ability to detect causally important transcriptomic changes relative to “passenger” changes. We can detect this by performing mediation analysis in reverse, with phenotypes as the mediator and gene expression as the outcome. Individual transcripts that are highly ranked in both models (i.e., with transcriptome as a mediator and an outcome) will




Figure 4: Previous study demonstrating feasibility of mapping novel histological phenotypes generated by AI. A) An age-related QTI-generated histological feature mapped to a genome-wide significant QTL on mouse chromosome 17. The solid and dashed red lines indicate genome-wide significance of 1% and 5% respectively. B) Montage of images comparing glomeruli that scored high and low for this mapped feature. High- and low-scoring glomeruli differ in mesangial matrix expansion (white areas? XXX), cellularity (blue areas? XXX). C) Founder allele effects at the QTL for this feature. Each founder is represented by one color. NZO alleles (light blue) were associated with better outcomes than A/J alleles (yellow).

be considered confounded. If this occurs, we will cross-reference our kidney expression to existing eQTL data from DO kidneys in mice without a TSC insult[?]. This will give added confidence in specific, genetically encoded transcript changes that cannot be due to tumor-composition effects. These data will also provide a strong rationale for follow-up proposals to generate single-cell and spatial transcriptomic atlases from our biobanked samples, from which we can deconvolve our bulk expression computationally[?]. A second potential problem is that we may fail to detect overt lesions using an unsupervised PCA approach. While we consider this extremely unlikely, if this occurs, we anticipate that it will be due to the heterogeneity of normal tissue structure in the DO obscuring the effects of relatively rarer lesional tissue. If our PCA montages (see Fig. ??B) do not contrast normal tissue with lesional tissue, then we will consult with JAX's on-site veterinary pathologist to both annotate the learned, non-lesional features and provide annotations of lesions that were missed by our analysis. With ground-truth labels of novel features, we can train models *de novo* to detect such features directly, which we will use as an additional phenotype. If lesions are extremely rare, we will apply outlier detection methodologies[?] to identify image tiles that exist far outside the distribution of healthy tissue. If all else fails, we will review each whole-slide image in its entirety, manually annotating abnormalities. These images will become public upon project completion for researchers in the wider community to apply their expertise and define novel phenotypes.

Overall timeline

Integration of aims and project growth: The aims of this project are conceptually and operationally distinct, but form an integrated whole. If both are successful, we will be positioned to perform a joint analysis of neurological and tumor outcomes. Indeed, we anticipate that many genetic modifiers will be pleiotropic across both classes

of outcomes[?]. By mapping pleiotropic modifiers, we may identify universal disease-modifying targets. Going forward, we believe that the DO-TSC model from this Idea Development proposal will provide a platform for multiple, critical follow-up research programs, including an extended search for disease modifiers, pharmacogenetic studies of resistance to therapies (*e.g.*, rapalogs), and pre-clinical drug screening in genetically diverse animals, minimizing the probability of false-positive hits. Thus, by developing the DO-TSC population model, we will enable a significant new paradigm for model systems studies of TSC.

References

- Al-Shaer, A. E., Pal, A., Shi, Q., Carson, M. S., Regan, J., Behee, M., Buddenbaum, N., Drawdy, C., Davis, T., Virk, R., and Shaikh, S. R. (2022). Modeling human heterogeneity of obesity with diversity outbred mice reveals a fat mass-dependent therapeutic window for resolvin E1. *FASEB J*, 36(6):e22354.
- Aydin, S., Pham, D. T., Zhang, T., Keele, G. R., Skelly, D. A., Paulo, J. A., Pankratz, M., Choi, T., Gygi, S. P., Reinholdt, L. G., Baker, C. L., Churchill, G. A., and Munger, S. C. (2023). Genetic dissection of the pluripotent proteome through multi-omics data integration. *Cell Genom*, 3(4):100283.
- Backer, G., Eddy, S., Sheehan, S. M., Takemon, Y., Reznichenko, A., Savage, H. S., Kretzler, M., and Korstanje, R. (2018). FAR2 is associated with kidney disease in mice and humans. *Physiol Genomics*, 50(8):543–552.
- Bagley, J. R., Bailey, L. S., Gagnon, L. H., He, H., Philip, V. M., Reinholdt, L. G., Tarantino, L. M., Chesler, E. J., and Jentsch, J. D. (2022). Behavioral phenotypes revealed during reversal learning are linked with novel genetic loci in diversity outbred mice. *Addict Neurosci*, 4.
- Bagley, J. R., Chesler, E. J., Philip, V. M., and Jentsch, J. D. (2021). Heritability of ethanol consumption and pharmacokinetics in a genetically diverse panel of collaborative cross mouse strains and their inbred founders. *Alcohol Clin Exp Res*, 45(4):697–708.
- Blair, R. H., Kliebenstein, D. J., and Churchill, G. A. (2012). What can causal networks tell us about metabolic pathways? *PLoS Comput Biol*, 8(4):e1002458.
- Bollen, K. A. (1989). *Structural equations with latent variables*, volume 210. John Wiley & Sons.
- Boyle, E. A., Li, Y. I., and Pritchard, J. K. (2017). An Expanded View of Complex Traits: From Polygenic to Omnigenic. *Cell*, 169(7):1177–1186.
- Brabec, J. L., Lara, M. K., Tyler, A. L., and Mahoney, J. M. (2021). System-Level Analysis of Alzheimer’s Disease Prioritizes Candidate Genes for Neurodegeneration. *Front Genet*, 12:625246.
- Broad Institute (2023). Picard tools.
- Broman, K. W., Gatti, D. M., Simecek, P., Furlotte, N. A., Prins, P., Sen, S., Yandell, B. S., and Churchill, G. A. (2019). R/qt12: Software for Mapping Quantitative Trait Loci with High-Dimensional Data and Multiparent Populations. *Genetics*, 211(2):495–502.
- Bubier, J. A., He, H., Philip, V. M., Roy, T., Hernandez, C. M., Bernat, R., Donohue, K. D., O’Hara, B. F., and Chesler, E. J. (2020). Genetic variation regulates opioid-induced respiratory depression in mice. *Sci Rep*, 10(1):14970.
- Campanella, G., Hanna, M. G., Geneslaw, L., Miraflor, A., Werneck Krauss Silva, V., Busam, K. J., Brogi, E., Reuter, V. E., Klimstra, D. S., and Fuchs, T. J. (2019). Clinical-grade computational pathology using weakly supervised deep learning on whole slide images. *Nat Med*, 25(8):1301–1309.
- Canpolat, M., Gumus, H., Kumandas, S., Coskun, A., and Per, H. (2018). The use of rapamycin in patients with tuberous sclerosis complex: Long-term results. *Epilepsy Behav*, 88:357–364.
- Caylor, R. C., Grote, L., Thiffault, I., Farrow, E. G., Willig, L., Soden, S., Amudhavalli, S. M., Nopper, A. J., Horii, K. A., Fleming, E., Jenkins, J., Welsh, H., Ilyas, M., Engleman, K., Abdelmoity, A., and Saunders, C. J. (2018). Incidental diagnosis of tuberous sclerosis complex by exome sequencing in three families with subclinical findings. *Neurogenetics*, 19(3):205–213.
- Chella Krishnan, K., El Hachem, E. J., Keller, M. P., Patel, S. G., Carroll, L., Vegas, A. D., Gerdes Gyuricza, I., Light, C., Cao, Y., Pan, C., Kaczor-Urbanowicz, K. E., Shravah, V., Anum, D., Pellegrini, M., Lee, C. F., Seldin, M. M., Rosenthal, N. A., Churchill, G. A., Attie, A. D., Parker, B., James, D. E., and Lusi, A. J. (2023). Genetic architecture of heart mitochondrial proteome influencing cardiac hypertrophy. *Elife*, 12.

- Cheng, N., Ren, Y., Zhou, J., Zhang, Y., Wang, D., Zhang, X., Chen, B., Liu, F., Lv, J., Cao, Q., Chen, S., Du, H., Hui, D., Weng, Z., Liang, Q., Su, B., Tang, L., Han, L., Chen, J., and Shao, C. (2022). Deep Learning-Based Classification of Hepatocellular Nodular Lesions on Whole-Slide Histopathologic Images. *Gastroenterology*, 162(7):1948–1961.
- Chick, J. M., Munger, S. C., Simecek, P., Huttlin, E. L., Choi, K., Gatti, D. M., Raghupathy, N., Svenson, K. L., Churchill, G. A., and Gygi, S. P. (2016). Defining the consequences of genetic variation on a proteome-wide scale. *Nature*, 534(7608):500–505.
- Churchill, G. A., Gatti, D. M., Munger, S. C., and Svenson, K. L. (2012). The Diversity Outbred mouse population. *Mamm Genome*, 23(9-10):713–718.
- Civelek, M. and Lusis, A. J. (2014). Systems genetics approaches to understand complex traits. *Nat Rev Genet*, 15(1):34–48.
- Coffey, A. R., Smallwood, T. L., Albright, J., Hua, K., Kanke, M., Pomp, D., Bennett, B. J., and Sethupathy, P. (2017). Systems genetics identifies a co-regulated module of liver microRNAs associated with plasma LDL cholesterol in murine diet-induced dyslipidemia. *Physiol Genomics*, 49(11):618–629.
- Cornes, B. K., Paisie, C., Swanzy, E., Fields, P. D., Schile, A., Brackett, K., Reinholdt, L. G., and Srivastava, A. (2023). Protein coding variation in the J:ARC and J:DO outbred laboratory mouse stocks provides a molecular basis for distinct research applications. *G3 (Bethesda)*, 13(4).
- Crammer, K. and Chechik, G. (2004). A needle in a haystack: local one-class optimization. In *Proceedings of the twenty-first international conference on Machine learning*, page 26.
- Crino, P. B., Nathanson, K. L., and Henske, E. P. (2006). The tuberous sclerosis complex. *N Engl J Med*, 355(13):1345–1356.
- Crouse, W. L., Keele, G. R., Gastonguay, M. S., Churchill, G. A., and Valdar, W. (2022). A Bayesian model selection approach to mediation analysis. *PLoS Genet*, 18(5):e1010184.
- Dalle Pezze, P., Sonntag, A. G., Thien, A., Prentzell, M. T., del, M., Fischer, S., Neumann-Haefelin, E., Huber, T. B., Baumeister, R., Shanley, D. P., and Thedieck, K. (2012). A dynamic network model of mTOR signaling reveals TSC-independent mTORC2 regulation. *Sci Signal*, 5(217):ra25.
- Delprato, A., Bonheur, B., o, M. P., Murillo, A., Dhawan, E., Lu, L., Williams, R. W., and Crusio, W. E. (2018). A quantitative trait locus on chromosome 1 modulates intermale aggression in mice. *Genes Brain Behav*, 17(7):e12469.
- Delprato, A., Bonheur, B., o, M. P., Rosay, P., Lu, L., Williams, R. W., and Crusio, W. E. (2015). Systems genetic analysis of hippocampal neuroanatomy and spatial learning in mice. *Genes Brain Behav*, 14(8):591–606.
- Delprato, A., o, M. P., Bonheur, B., Bubier, J. A., Lu, L., Williams, R. W., Chesler, E. J., and Crusio, W. E. (2017). QTL and systems genetics analysis of mouse grooming and behavioral responses to novelty in an open field. *Genes Brain Behav*, 16(8):790–799.
- Doerge, R. W. and Churchill, G. A. (1996). Permutation tests for multiple loci affecting a quantitative character. *Genetics*, 142(1):285–294.
- Dutta, P., Akhter Sathi, K., and Saiful Islam, M. (2022). Multi-classification of brain tumor images using transfer learning based deep neural network. In *International Conference on Artificial Intelligence for Smart Community: AISC 2020, 17–18 December, Universiti Teknologi Petronas, Malaysia*, pages 927–933. Springer.
- Eggermann, T., Spengler, S., Denecke, B., Zerres, K., and Mache, C. J. (2013). Multi-exon deletion in the XDH gene as a cause of classical xanthinuria. *Clin Nephrol*, 79(1):78–80.

- Ewels, P., Magnusson, M., Lundin, S., and Iler, M. (2016). MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*, 32(19):3047–3048.
- Farooq, M. A., Khatoon, A., Varkarakis, V., and Corcoran, P. (2020). Advanced deep learning methodologies for skin cancer classification in prodromal stages. *arXiv preprint arXiv:2003.06356*.
- Fox, J., Ben-Shachar, S., Uliel, S., Svirsky, R., Saitsu, H., Matsumoto, N., and Fattal-Valevski, A. (2017). Rare familial TSC2 gene mutation associated with atypical phenotype presentation of Tuberous Sclerosis Complex. *Am J Med Genet A*, 173(3):744–748.
- Gatti, D. M., Svenson, K. L., Shabalin, A., Wu, L. Y., Valdar, W., Simecek, P., Goodwin, N., Cheng, R., Pomp, D., Palmer, A., Chesler, E. J., Broman, K. W., and Churchill, G. A. (2014). Quantitative trait locus mapping methods for diversity outbred mice. *G3 (Bethesda)*, 4(9):1623–1633.
- Gerdes Gyuricza, I., Chick, J. M., Keele, G. R., Deighan, A. G., Munger, S. C., Korstanje, R., Gygi, S. P., and Churchill, G. A. (2022). Genome-wide transcript and protein analysis highlights the role of protein homeostasis in the aging mouse heart. *Genome Res*, 32(5):838–852.
- Geuther, B., Chen, M., Galante, R. J., Han, O., Lian, J., George, J., Pack, A. I., and Kumar, V. (2022). High-throughput visual assessment of sleep stages in mice using machine learning. *Sleep*, 45(2).
- Geuther, B. Q., Peer, A., He, H., Sabnis, G., Philip, V. M., and Kumar, V. (2021). Action detection using a neural network elucidates the genetics of mouse grooming behavior. *Elife*, 10.
- Gilboa, T., Segel, R., Zeligson, S., Alterescu, G., and Ben-Pazi, H. (2018). Ganglioglioma, Epilepsy, and Intellectual Impairment due to Familial TSC1 Deletion. *J Child Neurol*, 33(7):482–486.
- Glassbrook, J. E., Hackett, J. B., iz, M. C., Bross, M., Dyson, G., Movahhedine, N., Ullrich, A., and Gibson, H. M. (2023). Host genetic background regulates the capacity for anti-tumor antibody-dependent phagocytosis. *bioRxiv*.
- Gomez, M., Sampson, J., and Whittemore, V. (1999). *Tuberous Sclerosis Complex*. Developmental Perspectives in Psychiatry. Oxford University Press.
- Goto, T. and Monk, M. (1998). Regulation of X-chromosome inactivation in development in mice and humans. *Microbiol Mol Biol Rev*, 62(2):362–378.
- Gould, R. L., Craig, S. W., McClatchy, S., Churchill, G. A., and Pazdro, R. (2021). Genetic mapping of renal glutathione suggests a novel regulatory locus on the murine X chromosome and overlap with hepatic glutathione regulation. *Free Radic Biol Med*, 174:28–39.
- Graving, J. M., Chae, D., Naik, H., Li, L., Koger, B., Costelloe, B. R., and Couzin, I. D. (2019). DeepPoseKit, a software toolkit for fast and robust animal pose estimation using deep learning. *Elife*, 8.
- Gschwind, T., Zeine, A., Raikov, I., Markowitz, J. E., Gillis, W. F., Felong, S., Isom, L. L., Datta, S. R., and Soltesz, I. (2023). Hidden behavioral fingerprints in epilepsy. *Neuron*, 111(9):1440–1452.
- Gu, T., Gatti, D. M., Srivastava, A., Snyder, E. M., Raghupathy, N., Simecek, P., Svenson, K. L., Dotu, I., Chuang, J. H., Keller, M. P., Attie, A. D., Braun, R. E., and Churchill, G. A. (2016). Genetic Architectures of Quantitative Variation in RNA Editing Pathways. *Genetics*, 202(2):787–798.
- Gupta, N., Singh, N., Sarangi, S., Dalmia, S., and Mittal, S. (2008). Fetal cardiac rhabdomyoma with maternal tuberous sclerosis complicating pregnancy. *Arch Gynecol Obstet*, 278(2):169–170.
- Gupta, Y., Ernst, A. L., Vorobyev, A., Beltsiou, F., Zillikens, D., Bieber, K., Sanna-Cherchi, S., Christiano, A. M., Sadik, C. D., Ludwig, R. J., and Sezin, T. (2023). Impact of diet and host genetics on the murine intestinal mycobiome. *Nat Commun*, 14(1):834.

- Hackett, J. B., Glassbrook, J. E., iz, M. C., Bross, M., Fielder, A., Dyson, G., Movahhedin, N., McCasland, J., McCarthy-Leo, C., and Gibson, H. M. (2022). A diversity outbred F1 mouse model identifies host-intrinsic genetic regulators of response to immune checkpoint inhibitors. *Oncoimmunology*, 11(1):2064958.
- Hession, L. E., Sabnis, G. S., Churchill, G. A., and Kumar, V. (2022). A machine-vision-based frailty index for mice. *Nat Aging*, 2(8):756–766.
- Huda, M. N., VerHague, M., Albright, J., Smallwood, T., Bell, T. A., Que, E., Miller, D. R., Roshanravan, B., Allayee, H., Manuel de Villena, F. P., and Bennett, B. J. (2020). Dissecting the Genetic Architecture of Cystatin C in Diversity Outbred Mice. *G3 (Bethesda)*, 10(7):2529–2541.
- Huynh, K. D. and Lee, J. T. (2005). X-chromosome inactivation: a hypothesis linking ontogeny and phylogeny. *Nat Rev Genet*, 6(5):410–418.
- Jacob, J. B., Wei, K. C., Bepler, G., Reyes, J. D., Cani, A., Polin, L., White, K., Kim, S., Viola, N., McGrath, J., Guastella, A., Yin, C., Mi, Q. S., Kidder, B. L., Wagner, K. U., Ratner, S., Phillips, V., Xiu, J., Parajuli, P., and Wei, W. Z. (2023). Identification of actionable targets for breast cancer intervention using a diversity outbred mouse model. *iScience*, 26(4):106320.
- Kang, H. M., Zaitlen, N. A., Wade, C. M., Kirby, A., Heckerman, D., Daly, M. J., and Eskin, E. (2008). Efficient control of population structure in model organism association mapping. *Genetics*, 178(3):1709–1723.
- Karbowniczek, M., Cash, T., Cheung, M., Robertson, G. P., Astrinidis, A., and Henske, E. P. (2004). Regulation of B-Raf kinase activity by tuberin and Rheb is mammalian target of rapamycin (mTOR)-independent. *J Biol Chem*, 279(29):29930–29937.
- Karbowniczek, M., Robertson, G. P., and Henske, E. P. (2006). Rheb inhibits C-raf activity and B-raf/C-raf heterodimerization. *J Biol Chem*, 281(35):25447–25456.
- Keele, G. R., Zhang, T., Pham, D. T., Vincent, M., Bell, T. A., Hock, P., Shaw, G. D., Paulo, J. A., Munger, S. C., de Villena, F. P., Ferris, M. T., Gygi, S. P., and Churchill, G. A. (2021). Regulation of protein abundance in genetically diverse mouse populations. *Cell Genom*, 1(1).
- Keenan, B. T., Galante, R. J., Lian, J., Zhang, L., Guo, X., Veatch, O. J., Chesler, E. J., O'Brien, W. T., Svenson, K. L., Churchill, G. A., and Pack, A. I. (2021). The dihydropyrimidine dehydrogenase gene contributes to heritable differences in sleep in mice. *Curr Biol*, 31(23):5238–5248.
- Keller, M. P., Gatti, D. M., Schueler, K. L., Rabaglia, M. E., Stapleton, D. S., Simecek, P., Vincent, M., Allen, S., Broman, A. T., Bacher, R., Kendzierski, C., Broman, K. W., Yandell, B. S., Churchill, G. A., and Attie, A. D. (2018). Genetic Drivers of Pancreatic Islet Function. *Genetics*, 209(1):335–356.
- Kenerson, H., Dundon, T. A., and Yeung, R. S. (2005). Effects of rapamycin in the Eker rat model of tuberous sclerosis complex. *Pediatr Res*, 57(1):67–75.
- Kikuchi, Y., Sudo, A., Mitani, H., and Hino, O. (2004). Presence of a modifier gene(s) affecting early renal carcinogenesis in the Tsc2 mutant (Eker) rat model. *Int J Oncol*, 24(1):75–80.
- Kim, M., Huda, M. N., Evans, L. W., Que, E., Gertz, E. R., Maeda-Smithies, N., and Bennett, B. J. (2023). Integrative analysis of hepatic transcriptional profiles reveals genetic regulation of atherosclerosis in hyperlipidemic Diversity Outbred-F1 mice. *Sci Rep*, 13(1):9475.
- King, J. A. and Stamilio, D. M. (2005). Maternal and fetal tuberous sclerosis complicating pregnancy: a case report and overview of the literature. *Am J Perinatol*, 22(2):103–108.
- Koyuncu, D., Niazi, M. K. K., Tavolara, T., Abeijon, C., Ginese, M. L., Liao, Y., Mark, C., Specht, A., Gower, A. C., Restrepo, B. I., Gatti, D. M., Kramnik, I., Gurcan, M., Yener, B., and Beamer, G. (2021). CXCL1: A

new diagnostic biomarker for human tuberculosis discovered using Diversity Outbred mice. *PLoS Pathog*, 17(8):e1009773.

- [] Kwiatkowski, D. J. (2010). Animal models of lymphangioleiomyomatosis (LAM) and tuberous sclerosis complex (TSC). *Lymphat Res Biol*, 8(1):51–57.
- [] Lee, L., Sudentas, P., and Dabora, S. L. (2006). Combination of a rapamycin analog (CCI-779) and interferon-gamma is more effective than single agents in treating a mouse model of tuberous sclerosis complex. *Genes Chromosomes Cancer*, 45(10):933–944.
- [] Lee, L., Sudentas, P., Donohue, B., Asrican, K., Worku, A., Walker, V., Sun, Y., Schmidt, K., Albert, M. S., El-Hashemite, N., Lader, A. S., Onda, H., Zhang, H., Kwiatkowski, D. J., and Dabora, S. L. (2005). Efficacy of a rapamycin analog (CCI-779) and IFN-gamma in tuberous sclerosis mouse models. *Genes Chromosomes Cancer*, 42(3):213–227.
- [] Li, B. and Dewey, C. N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics*, 12:323.
- [] Li, C., Lee, P. S., Sun, Y., Gu, X., Zhang, E., Guo, Y., Wu, C. L., Auricchio, N., Priolo, C., Li, J., Csibi, A., Parkhitko, A., Morrison, T., Planaguma, A., Kazani, S., Israel, E., Xu, K. F., Henske, E. P., Blenis, J., Levy, B. D., Kwiatkowski, D., and Yu, J. J. (2014a). Estradiol and mTORC2 cooperate to enhance prostaglandin biosynthesis and tumorigenesis in TSC2-deficient LAM cells. *J Exp Med*, 211(1):15–28.
- [] Li, C., Zhang, E., Sun, Y., Lee, P. S., Zhan, Y., Guo, Y., Osorio, J. C., Rosas, I. O., Xu, K. F., Kwiatkowski, D. J., and Yu, J. J. (2014b). Rapamycin-insensitive up-regulation of adipocyte phospholipase A2 in tuberous sclerosis and lymphangioleiomyomatosis. *PLoS One*, 9(10):e104809.
- [] Li, M., Zhou, Y., Chen, C., Yang, T., Zhou, S., Chen, S., Wu, Y., and Cui, Y. (2019). Efficacy and safety of mTOR inhibitors (rapamycin and its analogues) for tuberous sclerosis complex: a meta-analysis. *Orphanet J Rare Dis*, 14(1):39.
- [] Liu, X., Li, Y. I., and Pritchard, J. K. (2019). Trans Effects on Gene Expression Can Drive Omnigenic Inheritance. *Cell*, 177(4):1022–1034.
- [] Liu, X. Y., Zhang, F. R., Shang, J. Y., Liu, Y. Y., Lv, X. F., Yuan, J. N., Zhang, T. T., Li, K., Lin, X. C., Liu, X., Lei, Q., Fu, X. D., Zhou, J. G., and Liang, S. J. (2018). Renal inhibition of miR-181a ameliorates 5-fluorouracil-induced mesangial cell apoptosis and nephrotoxicity. *Cell Death Dis*, 9(6):610.
- [] Liu, Y., Marks, K., Cowley, G. S., Carretero, J., Liu, Q., Nieland, T. J., Xu, C., Cohoon, T. J., Gao, P., Zhang, Y., Chen, Z., Altabef, A. B., Tchaicha, J. H., Wang, X., Choe, S., Driggers, E. M., Zhang, J., Bailey, S. T., Sharpless, N. E., Hayes, D. N., Patel, N. M., Janne, P. A., Bardeesy, N., Engelman, J. A., Manning, B. D., Shaw, R. J., Asara, J. M., Scully, R., Kimmelman, A., Byers, L. A., Gibbons, D. L., Wistuba, I. I., Heymach, J. V., Kwiatkowski, D. J., Kim, W. Y., Kung, A. L., Gray, N. S., Root, D. E., Cantley, L. C., and Wong, K. K. (2013). Metabolic and functional genomic studies identify deoxythymidylate kinase as a target in LKB1-mutant lung cancer. *Cancer Discov*, 3(8):870–879.
- [] Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*, 15(12):550.
- [] Lozovaya, N., Gataullina, S., Tsintsadze, T., Tsintsadze, V., Pallesi-Pocachard, E., Minlebaev, M., Goriounova, N. A., Buhler, E., Watrin, F., Shityakov, S., Becker, A. J., Bordey, A., Milh, M., Scavarda, D., Bulteau, C., Dorfmueller, G., Delalande, O., Represa, A., Cardoso, C., Dulac, O., Ben-Ari, Y., and Burnashev, N. (2014). Selective suppression of excessive GluN2C expression rescues early epilepsy in a tuberous sclerosis murine model. *Nat Commun*, 5:4563.

- Lyczkowski, D. A., Conant, K. D., Pulsifer, M. B., Jarrett, D. Y., Grant, P. E., Kwiatkowski, D. J., and Thiele, E. A. (2007). Intrafamilial phenotypic variability in tuberous sclerosis complex. *J Child Neurol*, 22(12):1348–1355.
- Markowitz, J. E., Gillis, W. F., Beron, C. C., Neufeld, S. Q., Robertson, K., Bhagat, N. D., Peterson, R. E., Peterson, E., Hyun, M., Linderman, S. W., Sabatini, B. L., and Datta, S. R. (2018). The Striatum Organizes 3D Behavior via Moment-to-Moment Action Selection. *Cell*, 174(1):44–58.
- Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W., and Bethge, M. (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat Neurosci*, 21(9):1281–1289.
- Matsuzaki, G., Lin, T., and Nomoto, K. (1994). Differentiation and function of intestinal intraepithelial lymphocytes. *Int Rev Immunol*, 11(1):47–60.
- Melia, T. and Waxman, D. J. (2019). Sex-Biased lncRNAs Inversely Correlate With Sex-Opposite Gene Coexpression Networks in Diversity Outbred Mouse Liver. *Endocrinology*, 160(5):989–1007.
- Melia, T. and Waxman, D. J. (2020). Genetic factors contributing to extensive variability of sex-specific hepatic gene expression in Diversity Outbred mice. *PLoS One*, 15(12):e0242665.
- Millstein, J. and Volfson, D. (2013). Computationally efficient permutation-based confidence interval estimation for tail-area FDR. *Front Genet*, 4:179.
- Moran, M. M., Ko, F. C., Mesner, L. D., Calabrese, G. M., Al-Barghouthi, B. M., Farber, C. R., and Sumner, D. R. (2022). Intramembranous bone regeneration in diversity outbred mice is heritable. *Bone*, 164:116524.
- n, O. Y., Crainiceanu, C., Ogburn, E. L., Caffo, B. S., Wager, T. D., and Lindquist, M. A. (2018). High-dimensional multivariate mediation with application to neuroimaging data. *Biostatistics*, 19(2):121–136.
- Neto, E. C., Keller, M. P., Attie, A. D., and Yandell, B. S. (2010). CAUSAL GRAPHICAL MODELS IN SYSTEMS GENETICS: A UNIFIED FRAMEWORK FOR JOINT INFERENCE OF CAUSAL NETWORK AND GENETIC ARCHITECTURE FOR CORRELATED PHENOTYPES. *Ann Appl Stat*, 4(1):320–339.
- Northrup, H., Wheless, J. W., Bertin, T. K., and Lewis, R. A. (1993). Variability of expression in tuberous sclerosis. *J Med Genet*, 30(1):41–43.
- Onda, H., Lueck, A., Marks, P. W., Warren, H. B., and Kwiatkowski, D. J. (1999). Tsc2(+/-) mice develop tumors in multiple sites that express gelsolin and are influenced by genetic background. *J Clin Invest*, 104(6):687–695.
- Ouellette, A. R., Neuner, S. M., Dumitrescu, L., Anderson, L. C., Gatti, D. M., Mahoney, E. R., Bubier, J. A., Churchill, G., Peters, L., Huentelman, M. J., Herskowitz, J. H., Yang, H. S., Smith, A. N., Reitz, C., Kunkle, B. W., White, C. C., De Jager, P. L., Schneider, J. A., Bennett, D. A., Seyfried, N. T., Chesler, E. J., Hadad, N., Hohman, T. J., and Kaczorowski, C. C. (2020). Cross-Species Analyses Identify Dlgap2 as a Regulator of Age-Related Cognitive Decline and Alzheimer’s Dementia. *Cell Rep*, 32(9):108091.
- Parker, C. C., Philip, V. M., Gatti, D. M., Kasperek, S., Kreuzman, A. M., Kuffler, L., Mansky, B., Masneuf, S., Sharif, K., Sluys, E., Tattera, D., Taylor, W. M., Thomas, M., Polesskaya, O., Palmer, A. A., Holmes, A., and Chesler, E. J. (2022). Genome-wide association mapping of ethanol sensitivity in the Diversity Outbred mouse population. *Alcohol Clin Exp Res*, 46(6):941–960.
- Pereira, T. D., Aldarondo, D. E., Willmore, L., Kislin, M., Wang, S. S., Murthy, M., and Shaevitz, J. W. (2019). Fast animal pose estimation using deep neural networks. *Nat Methods*, 16(1):117–125.
- Philip, V. M., He, H., Saul, M. C., Dickson, P. E., Bubier, J. A., and Chesler, E. J. (2023). Gene expression genetics of the striatum of Diversity Outbred mice. *bioRxiv*.
- Philip, V. M., Tyler, A. L., and Carter, G. W. (2014). Dissection of complex gene expression using the combined analysis of pleiotropy and epistasis. *Pac Symp Biocomput*, pages 200–211.

- Price, T. R., Stapleton, D. S., Schueler, K. L., Norris, M. K., Parks, B. W., Yandell, B. S., Churchill, G. A., Holland, W. L., Keller, M. P., and Attie, A. D. (2023).) as an enzyme that metabolizes phosphatidylcholine and cardiolipin. *bioRxiv*.
- Que, E., James, K. L., Coffey, A. R., Smallwood, T. L., Albright, J., Huda, M. N., Pomp, D., Sethupathy, P., and Bennett, B. J. (2020). Genetic Architecture Modulates Diet-Induced Hepatic mRNA and miRNA Expression Profiles in Diversity Outbred Mice. *Genetics*, 216(1):241–259.
- Roberts, A., Pardo-Manuel de Villena, F., Wang, W., McMillan, L., and Threadgill, D. W. (2007). The polymorphism architecture of mouse genetic resources elucidated using genome-wide resequencing data: implications for QTL discovery and systems genetics. *Mamm Genome*, 18(6-7):473–481.
- Sanderson, B., Fields, P., and M.W., L. (2023). JAX NGS Operations Nextflow DSL2 Pipelines.
- Sharma, N., Sharma, S., Thiek, J. L., Ahanthem, S. S., Kalita, A., and Lynser, D. (2017). Maternal and Fetal Tuberous Sclerosis: Do We Know Enough as an Obstetrician? *J Reprod Infertil*, 18(2):257–260.
- Sheehan, S., Mawe, S., Cianciolo, R. E., Korstanje, R., and Mahoney, J. M. (2019). Detection and Classification of Novel Renal Histologic Phenotypes Using Deep Neural Networks. *Am J Pathol*, 189(9):1786–1796.
- Sheppard, K., Gardin, J., Sabnis, G. S., Peer, A., Darrell, M., Deats, S., Geuther, B., Lutz, C. M., and Kumar, V. (2022). Stride-level analysis of mouse open field behavior using deep-learning-based pose estimation. *Cell Rep*, 38(2):110231.
- Sigmon, J. S., Blanchard, M. W., Baric, R. S., Bell, T. A., Brennan, J., Brockmann, G. A., Burks, A. W., Calabrese, J. M., Caron, K. M., Cheney, R. E., Ciavatta, D., Conlon, F., Darr, D. B., Faber, J., Franklin, C., Gershon, T. R., Gralinski, L., Gu, B., Gaines, C. H., Hagan, R. S., Heimsath, E. G., Heise, M. T., Hock, P., Ideraabdullah, F., Jennette, J. C., Kafri, T., Kashfeen, A., Kulis, M., Kumar, V., Linnertz, C., Livraghi-Butrico, A., Lloyd, K. C. K., Lutz, C., Lynch, R. M., Magnuson, T., Matsushima, G. K., McMullan, R., Miller, D. R., Mohlke, K. L., Moy, S. S., Murphy, C. E. Y., Najarian, M., O'Brien, L., Palmer, A. A., Philpot, B. D., Randell, S. H., Reinholdt, L., Ren, Y., Rockwood, S., Rogala, A. R., Saraswatula, A., Sassetti, C. M., Schisler, J. C., Schoenrock, S. A., Shaw, G. D., Shorter, J. R., Smith, C. M., St Pierre, C. L., Tarantino, L. M., Threadgill, D. W., Valdar, W., Vilen, B. J., Wardwell, K., Whitmire, J. K., Williams, L., Zylka, M. J., Ferris, M. T., McMillan, L., and Manuel de Villena, F. P. (2020). Content and Performance of the MiniMUGA Genotyping Array: A New Tool To Improve Rigor and Reproducibility in Mouse Research. *Genetics*, 216(4):905–930.
- Skelly, D. A., Czechanski, A., Byers, C., Aydin, S., Spruce, C., Olivier, C., Choi, K., Gatti, D. M., Raghupathy, N., Keele, G. R., Stanton, A., Vincent, M., Dion, S., Greenstein, I., Pankratz, M., Porter, D. K., Martin, W., O'Connor, C., Qin, W., Harrill, A. H., Choi, T., Churchill, G. A., Munger, S. C., Baker, C. L., and Reinholdt, L. G. (2020). Mapping the Effects of Genetic Variation on Chromatin State and Gene Expression Reveals Loci That Control Ground State Pluripotency. *Cell Stem Cell*, 27(3):459–469.
- Soneson, C. and Delorenzi, M. (2013). A comparison of methods for differential expression analysis of RNA-seq data. *BMC Bioinformatics*, 14:91.
- Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., Paulovich, A., Pomeroy, S. L., Golub, T. R., Lander, E. S., and Mesirov, J. P. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*, 102(43):15545–15550.
- Szegedy, C., Vanhoucke, V., Ioffe, S., Shlens, J., and Wojna, Z. (2016). Rethinking the inception architecture for computer vision. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pages 2818–2826.
- Takemon, Y., Chick, J. M., Gerdes Gyuricza, I., Skelly, D. A., Devuyst, O., Gygi, S. P., Churchill, G. A., and

- Korstanje, R. (2021a). Proteomic and transcriptomic profiling reveal different aspects of aging in the kidney. *Elife*, 10.
- [] Takemon, Y., Wright, V., Davenport, B., Gatti, D. M., Sheehan, S. M., Letson, K., Savage, H. S., Lennon, R., and Korstanje, R. (2021b). Uncovering Modifier Genes of X-Linked Alport Syndrome Using a Novel Multiparent Mouse Model. *J Am Soc Nephrol*, 32(8):1961–1973.
- [] Tang, Y., Kwiatkowski, D. J., and Henske, E. P. (2022). Midkine expression by stem-like tumor cells drives persistence to mTOR inhibition and an immune-suppressive microenvironment. *Nat Commun*, 13(1):5018.
- [] Taveira-DaSilva, A. M. and Moss, J. (2012). Optimizing treatments for lymphangioleiomyomatosis. *Expert Rev Respir Med*, 6(3):267–276.
- [] The Jackson Laboratory (2022). Haploqa.
- [] Tyler, A., Mahoney, J. M., and Carter, G. W. (2020). Genetic Interactions Affect Lung Function in Patients with Systemic Sclerosis. *G3 (Bethesda)*, 10(1):151–163.
- [] Tyler, A. L., Donahue, L. R., Churchill, G. A., and Carter, G. W. (2016). Weak Epistasis Generally Stabilizes Phenotypes in a Mouse Intercross. *PLoS Genet*, 12(2):e1005805.
- [] Tyler, A. L., El Kassaby, B., Kolishovski, G., Emerson, J., Wells, A. E., Mahoney, J. M., and Carter, G. W. (2021). Effects of kinship correction on inflation of genetic interaction statistics in commonly used mouse populations. *G3 (Bethesda)*, 11(7).
- [] Tyler, A. L., Ji, B., Gatti, D. M., Munger, S. C., Churchill, G. A., Svenson, K. L., and Carter, G. W. (2017). Epistatic Networks Jointly Influence Phenotypes Related to Metabolic Disease and Gene Expression in Diversity Outbred Mice. *Genetics*, 206(2):621–639.
- [] Tyler, A. L., Lu, W., Hendrick, J. J., Philip, V. M., and Carter, G. W. (2013). CAPE: an R package for combined analysis of pleiotropy and epistasis. *PLoS Comput Biol*, 9(10):e1003270.
- [] Tyler, A. L., McGarr, T. C., Beyer, B. J., Frankel, W. N., and Carter, G. W. (2014). A genetic interaction network model of a complex neurological disease. *Genes Brain Behav*, 13(8):831–840.
- [] Tyler, A. L., Raza, A., Krementsov, D. N., Case, L. K., Huang, R., Ma, R. Z., Blankenhorn, E. P., Teuscher, C., and Mahoney, J. M. (2019). Network-Based Functional Prediction Augments Genetic Association To Predict Candidate Genes for Histamine Hypersensitivity in Mice. *G3 (Bethesda)*, 9(12):4223–4233.
- [] Tyler, A. L., Spruce, C., Kursawe, R., Haber, A., Ball, R. L., Pitman, W. A., Fine, A. D., Raghupathy, N., Walker, M., Philip, V. M., Baker, C. L., Mahoney, J. M., Churchill, G. A., Trowbridge, J. J., Stitzel, M. L., Paigen, K., Petkov, P. M., and Carter, G. W. (2023). Variation in histone configurations correlates with gene expression across nine inbred strains of mice. *Genome Res*.
- [] Wang, X., Park, J., Susztak, K., Zhang, N. R., and Li, M. (2019). Bulk tissue cell type deconvolution with multi-subject single-cell expression reference. *Nat Commun*, 10(1):380.
- [] Wei, W. Z., Gibson, H. M., Jacob, J. B., Frelinger, J. A., Berzofsky, J. A., Maeng, H., Dyson, G., Reyes, J. D., Pilon-Thomas, S., Ratner, S., and Wei, K. C. (2020). Diversity Outbred Mice Reveal the Quantitative Trait Locus and Regulatory Cells of HER2 Immunity. *J Immunol*, 205(6):1554–1563.
- [] Williams, E. G., Pfister, N., Roy, S., Statzer, C., Haverty, J., Ingels, J., Bohl, C., Hasan, M., uklina, J., hlmann, P., Zamboni, N., Lu, L., Ewald, C. Y., Williams, R. W., and Aebersold, R. (2022). Multiomic profiling of the liver across diets and age in a diverse mouse population. *Cell Syst*, 13(1):43–57.
- [] Wilson, C., Idziaszczyk, S., Parry, L., Guy, C., Griffiths, D. F., Lazda, E., Bayne, R. A., Smith, A. J., Sampson, J. R., and Cheadle, J. P. (2005). A mouse model of tuberous sclerosis 1 showing background specific early post-natal mortality and metastatic renal cell carcinoma. *Hum Mol Genet*, 14(13):1839–1850.

- Wiltchko, A. B., Johnson, M. J., Iurilli, G., Peterson, R. E., Katon, J. M., Pashkovski, S. L., Abaira, V. E., Adams, R. P., and Datta, S. R. (2015). Mapping Sub-Second Structure in Mouse Behavior. *Neuron*, 88(6):1121–1135.
- Winter, J. M., Gildea, D. E., Andreas, J. P., Gatti, D. M., Williams, K. A., Lee, M., Hu, Y., Zhang, S., Mullikin, J. C., Wolfsberg, T. G., McDonnell, S. K., Fogarty, Z. C., Larson, M. C., French, A. J., Schaid, D. J., Thibodeau, S. N., Churchill, G. A., and Crawford, N. P. (2017). Mapping Complex Traits in a Diversity Outbred F1 Mouse Population Identifies Germline Modifiers of Metastasis in Human Prostate Cancer. *Cell Syst*, 4(1):31–45.
- Xenakis, J. G., Douillet, C., Bell, T. A., Hock, P., Farrington, J., Liu, T., Murphy, C. E. Y., Saraswatula, A., Shaw, G. D., Nativio, G., Shi, Q., Venkatratnam, A., Zou, F., Fry, R. C., blo, M., and Pardo-Manuel de Villena, F. (2022). An interaction of inorganic arsenic exposure with body weight and composition on type 2 diabetes indicators in Diversity Outbred mice. *Mamm Genome*, 33(4):575–589.
- Xu, K. F., Tian, X., Yang, Y., and Zhang, H. (2018). Rapamycin for lymphangioleiomyomatosis: optimal timing and optimal dosage. *Thorax*, 73(4):308–310.
- Xu, Y., Hosny, A., Zeleznik, R., Parmar, C., Coroller, T., Franco, I., Mak, R. H., and Aerts, H. J. W. L. (2019). Deep Learning Predicts Lung Cancer Treatment Response from Serial Medical Imaging. *Clin Cancer Res*, 25(11):3266–3275.
- Yeung, R. S., Gu, H., Lee, M., and Dundon, T. A. (2001). Genetic identification of a locus, *Mot1*, that affects renal tumor size in the rat. *Genomics*, 78(3):108–112.
- Yu, J., Deng, Y., Liu, T., Zhou, J., Jia, X., Xiao, T., Zhou, S., Li, J., Guo, Y., Wang, Y., Zhou, J., and Chang, C. (2020). Lymph node metastasis prediction of papillary thyroid carcinoma based on transfer learning radiomics. *Nat Commun*, 11(1):4807.