

Supplementary File 1

S1.1 Obligatory criteria

Our differential expression analyses required non-normalised RNA-seq data for the hybrid and both parental species in each allopolyploid or homoploid hybrid complex. Only datasets that included at least two biological replicates of each species were considered. A genome or gene sequences for at least one of the parental species was needed for the utilisation of the HyLiTE pipeline (Duchemin et al., 2015) in the differential expression analyses. The implementation of GO for functional analyses required genome annotations for at least one parental species, however, both a genome and genome annotation were not required, therefore if a dataset only had gene annotations available it was still considered. The RNA-seq data for the parents and hybrid had to be extracted from the same tissue, or from cells grown in the same medium, where applicable, and extracted under the same conditions.

S1.2 Preferential criteria

We gave preference to datasets where the respective hybrid species from which RNA-seq data was obtained were naturally occurring (it is common in the literature to see synthetic hybrids; commonly formed via artificial crosses for homoploid hybrids or colchicine treatment to obtain allopolyploids), to mitigate the influence of synthetic formation on any observed gene expression patterns. We also gave preference to allopolyploid and homoploid hybrid complexes that were as taxonomically close as possible, to minimise taxon-specific differences in observed gene expression patterns. Moreover, we preferred that both parental species were extant, where possible, as opposed to extant relatives of the extinct parents.

A broad search of the literature and online databases was performed with the above criteria considered. These are the resulting options and their justifications for use, or omission, in differential gene expression analysis.

S1.3 Animals

Achondrostoma oligolepis x Pseudochondrostoma polylepis

Naturally-occurring homoploid hybrid of two Iberian Leuciscinae fish (Pereira et al., 2014). No comparative transcriptomic studies are available of this parent-hybrid complex.

Ambystoma

McElroy et al. (2017) studied an allotriploid *Ambystoma* (mole salamander) with three parental species. This would have complicated our analyses as the classes used in the categorisation of differential gene expression are based on a two parent system (Cox et al., 2014).

In other publications, it was unclear as to exactly how many species were parents (parental species could potentially range from 2-4) of a given hybrid, or if the hybrid had a duplicated genome (Bogart et al., 2007; Schmid et al., 2015).

We additionally searched NCBI BioProjects for *Ambystoma*. There were some hybrids amongst the results, but they did not specify polyploidy/homoploidy, and as no papers have given specific hybrid/parent complexes other than McElroy et al, *Ambystoma* were not used in the analyses.

Bacillus

Stick insect genus purportedly containing allopolyploid species (Mallet, 2007). Subsequent searches of Google to identify specific species were unsuccessful and no data could be found on NCBI SRA.

Bufo viridis

The *B. viridis* subgroup contains a complex of diploid, triploid and tetraploid green toad species from Central Asia. The parental species of the triploid and tetraploid species are unclear (Stöck et al., 2005; Betto-Colliard et al., 2015).

Bulinus truncates

Allopolyploid freshwater snail briefly mentioned in (Mallet, 2007). Searches of NCBI SRA found no data.

Carassius auratus x *Cyprinus carpio*

Both parental species of this allopolyploid fish (carp) had genome sequences and annotations; the issue was a lack of homoploid hybrid complex from within the carp or carp-like fishes. Due to the lack of an accompanying publication, it was also unclear if this allopolyploid (NCBI BioProject PRJNA86641) was naturally formed.

Carassius auratus gibelio

Naturally-occurring allopolyploid fish (carp) formed from *Carassius auratus* x *Cyprinus carpio* pairing (Li et al., 2014). Only an mtDNA genome was available and limited gene models (13), however, there were extensive genome annotations available for the parental species. The issue was a lack of homoploid hybrid complex from within the carp fishes.

Carassius auratus red var. x *Cyprinus carpio*

Allopolyploid fish (carp) formed via artificial cross (Wang et al., 2015).

Carassius auratus red var. x *Megalobrama amblycephala*

Allopolyploid fish (carp) with genome sequence, however, only the transcriptome sequence for one biological replicate (Qin et al., 2016).

Cichlids

Within the literature, there exist a number of studies on cichlids (fish, family: Cichlidae), e.g. (Stelkens et al., 2009; Genner & Turner, 2011; Selz et al., 2014; Santos et al., 2016), however, none produced RNA-seq data from a comparative analysis between hybrid and parents.

Cobitis taenia x *Colbitis elongatoides*

These two diploid fish hybridised to form both diploid and polyploid lineages. RNA-seq data was only available for one parental species (*C. taenia*) (Juchno et al., 2007; Juchno & Boroń, 2018).

Cottus gobio

Homoploid hybrid fish derived from *C. perifretum* x *C. rhenanum* (Mallet, 2007; Stemshorn et al., 2011). No RNA-seq data was available for the hybrid.

Ctenopharyngodon idellus x *Megalobrama amblycephala*

This interspecific diploid fish pairing can form both diploid and triploid hybrids (He et al., 2013), however, there is no transcriptomic study that encompasses both parental species and the hybrids in the literature.

Two transcriptome accessions for triploid *Ctenopharyngodon idellus* x *Megalobrama amblycephala* hybrids can be found under NCBI BioProject PRJNA240314.

1 *Danio rerio* x *Danio nigrofasciatus*

2 A homoploid hybrid zebrafish complex (Spiewak et al., 2018). Although extensive genomic
3 information is available for *D. rerio*, there are no comparative transcriptomic data available
4 for this complex.

5 Artificial triploid *D. rerio* can be generated, however these are not allopolyploid (Mizgireuv
6 et al., 2004).

7 *Daphnia*

8 The water flea genus *Daphnia* does contain hybrid species, such as *D. galeata* × *D. hyalina*
9 (Wolinska et al., 2006) and some allopolyploids within the *D. pulex* complex (Dufresne,
10 2011). Comparative RNA-seq data could not be found for either complex.

11 *Drosophila*

12 Interspecific homoploid hybrids are common in the fruit fly genus *Drosophila* (Goulielmos &
13 Alahiotis, 1989; Landry et al., 2005; Kelleher et al., 2012). Some *Drosophila* species do
14 exhibit mosaic polyploidy (Fox et al., 2010), but there are no entirely allopolyploid species.

15 *Epinephelus coioides* x *Epinephelus lanceolatus*

16 Under NCBI BioProject PRJNA431058 there are transcriptome accessions for diploid and
17 triploid hybrids from the *Epinephelus coioides* x *Epinephelus lanceolatus* (grouper fish)
18 pairing. However, there are no accompanying parental transcriptomes, and only one
19 biological replicate at each ploidy level with two different tissues sampled.

20 BioProject PRJNA497240 has transcriptomic sequences for *E. coioides*, however, there is
21 only one replicate subjected to different treatment condition. Likewise, BioProject
22 PRJNA413272 has transcriptomic sequences for a single replicate of *E. lanceolatus*, sampled
23 over a time series, post-hatch.

24 *Gasterosteus nipponicus* x *Gasterosteus aculeatus*

25 Parent-hybrid fish complex with 6 or 7 RNA-seq runs available for each species, found under
26 NCBI BioProject PRJDB7242. Genome with gene models also available for *G. aculeatus*
27 (three-spined stickleback fish). However, as data was unpublished, it was unclear if the
28 hybrid was homoploid and if it was naturally or synthetically formed.

1 *Gila seminuda*

2 Homoploid hybrid fish formed from the hybridisation of *Gila robusta* x *Gila elegans* (Mallet,
3 2007; Mavárez & Linares, 2008). Only an organellar genome is available for *G. robusta* on
4 NCBI, in addition to 13 gene models. Further, no comparative RNA-seq data is available for
5 this hybrid complex.

6 *Megalobrama amblycephala* x *Cyprinus carpio*

7 RNA-seq data for the hybrid was unavailable (Wang et al., 2017). The research group was
8 contacted but no response was obtained.

9 *Morone chrysops* x *Morone saxatilis*

10 Six transcriptome accessions of homoploid hybrid striped bass available under NCBI
11 BioProject PRJNA382266. There were no accompanying parental transcriptome accessions.
12 A single transcriptomic replicate for *M. chrysops* and *M. saxatilis* were found under
13 BioSample SRR1187194 and SRR1187193, respectively.

14 *Oncorhynchus*

15 This genus in the Salmonidae family contains a number of hybrid species, including
16 *Oncorhynchus keta* x *Oncorhynchus gorbuscha* (Zhivotovsky et al., 2016), and diploid and
17 triploid *Oncorhynchus kisutch* (Withler et al., 1995). No comparative transcriptomic analyses
18 were found for either an allopolyploid or homoploid *Oncorhynchus* complex.

19 *Pelophylax esculentus*

20 The hybrid water frog of *Pelophylax lessonae* x *Pelophylax ridibundus* (Doležálková-
21 Kaštánková et al., 2018). Searches of NCBI database and <https://amphibiaweb.org> did not
22 find any genome sequences or comparative RNA-seq analyses.

23 *Poecilia formosa*

24 Naturally-occurring allopolyploid fish (molly) formed from *Poecilia Mexicana* x *Poecilia*
25 *latipinna* (Schedina et al., 2014). Genome sequences and annotations were available for both
26 parental species and the allopolyploid. The issue was a lack of homoploid hybrid complex
27 within the mollies.

28 *Salmo salar*

Atlantic salmon can occur as natural diploid and artificial triploid hybrids (Refstie & Gjedrem, 1975; Vera et al., 2017; Murray et al., 2018). However, there are no comparative transcriptomic analyses of parental species and hybrids in the literature.

Squalius

A genus of Cyprinidae fish with allopolyploid and homoploid hybrids produced from the same parental pairing. *S. alburnoides* exists as natural populations of hybrid allotriploid PAA, homodiploid hybrid PA, and the parental-like diploid AA genotypes. The parental species of *S. alburnoides* are *S. pyrenaicus* (PP) and an extinct species related to *Anaocypris hispanica* (Pala et al., 2010; Matos et al., 2015).

There are genome sequences and annotations for *S. pyrenaicus* available at ftp://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/001/403/095/GCA_001403095.1_pp_cds_nonRedundant, there is also a GenBank record (accession CVRK000000000) for the *S. pyrenaicus* whole genome shotgun sequencing project, but it contains no sequence data.

Pala et al. (2008) performed cDNA sequencing of allotriploid *S. alburnoides*, but only for 6 genes.

There is comparative small RNA data available for this complex under GEO Series GSE38691 (Inácio et al., 2012).

Comparative RNA-seq data for allopolyploid and homoploid *Squalius* complexes can be found under ArrayExpress E-MTAB-3174 (Matos et al., 2015, 2019). This genus is a strong candidate for our homoploid and allopolyploid animal representatives.

Takifugu rubripes x *Takifugu flavidus*

An artificially formed homoploid hybrid fish (pufferfish), with transcriptome accessions only available for one replicate each of the parental species and hybrid (Gao et al., 2013).

Tigriopus californicus

Copepod species whose hybridisations are limited to being inter-population only (Pritchard et al., 2013; Barreto et al., 2014).

Xenopus laevis

Allopolyploid African clawed frog with unknown parental species (Session et al., 2016).

SI.4 Fungi

Aspergillus flavus x *Aspergillus parasiticus*

These Ascomycete molds form interspecific hybrid strains through artificial crosses, some with evidence of allopolyploidisation (Olarie et al., 2015). Genome sequences and annotations for both parental species are available through the NCBI database. No comparative transcriptomic studies available.

Blumeria graminis f. sp. *triticales*

A homoploid species of powdery mildew formed from the hybridisation of *B. g.* f. sp. *tritici* and *B. g.* f. sp. *secalis*. There is an annotated genome available for *B. g. tritici* and RNA-seq data for all members of this complex available. However, these transcriptomic data did not originate from a single study, so the conditions around RNA extraction could not be guaranteed as controlled across all samples, therefore we would not use this as a first option dataset (Menardo et al., 2016; Praz et al., 2018).

Additional comparative transcriptomic analyses relating to this complex involve only *B. g. tritici* (Hu et al., 2018).

Epichloë canadensis

Allopolyploid endophyte hybrid of *E. elymi* x *E. amarillans* with RNA-seq data with replicates for all members of the complex, in addition to genomes and gene sequences for both the parental species (data unpublished). A strong candidate for our allopolyploid fungal complex.

Neotyphodium lolii x *Epichloë typhina* Lp1

Allopolyploid endophyte. RNA-seq data with replicates available for all members of the complex, in addition to gene sequences for both the parental species (Cox et al., 2014; Campbell et al., 2017).

Ophiostoma ulmi x *Ophiostoma novo-ulmi*

Transient hybrids of *Ophiostoma ulmi* x *Ophiostoma novo-ulmi* do occur in nature (Brasier, 2001). No comparative transcriptomic analyses of this homoploid hybrid complex are

1 available; only a single study in which the *O. novo-ulmi* transcriptome was compared with
2 that of *Candida albicans* and *Histoplasma capsulatum* (Nigg et al., 2015).

3 *Saccharomyces cerevisiae* x *Saccharomyces bayanus*

4 Homoploid hybrid yeast. Genome annotations for both parental species are available at
5 [http://www.saccharomycessensustricto.org/cgi-](http://www.saccharomycessensustricto.org/cgi-bin/s3.cgi?data=Annotations&version=current)
6 [bin/s3.cgi?data=Annotations&version=current](http://www.saccharomycessensustricto.org/cgi-bin/s3.cgi?data=Annotations&version=current), in addition to RNA-seq reads from a
7 comparative transcriptomic analysis of this complex. The hybrids were generated
8 synthetically through mating a single haploid individual of each parental species, growing a
9 clonal population of the diploid, and checking it by PCR (Schraiber et al., 2013).

10 *Saccharomyces cerevisiae* x *Saccharomyces kudriavzevii*

11 There is a comparative transcriptomic study available that used two biological replicates of
12 this homoploid hybrid yeast, however, it used a microarray and only compared the hybrids at
13 varying temperatures (Tronchoni et al., 2017).

14 *Saccharomyces cerevisiae* x *Saccharomyces paradoxus*

15 Homoploid hybrid yeast. RNA-seq-based transcriptomic analyses of multiple replicates are
16 available. The hybrids were generated synthetically through mating a single haploid
17 individual of each parental species, growing a clonal population of the diploid, and checking
18 it by PCR (Schraiber et al., 2013). Genome annotations for both parental species available at
19 [http://www.saccharomycessensustricto.org/cgi-](http://www.saccharomycessensustricto.org/cgi-bin/s3.cgi?data=Annotations&version=current)
20 [bin/s3.cgi?data=Annotations&version=current](http://www.saccharomycessensustricto.org/cgi-bin/s3.cgi?data=Annotations&version=current). A strong candidate for our homoploid fungal
21 complex.

22 Swain Lenz et al. (2014) also did a comparative transcriptomic analysis of this complex,
23 however, due to the genetic manipulations performed during their synthetic crossing of the
24 parental species, we have opted to use the data from Schraiber et al.

25 *Saccharomyces cerevisiae* x *Saccharomyces uvarum*

26 Synthetically formed homoploid hybrids (Pfliegler et al., 2012). Although genome
27 annotations are available for both parental species on
28 [http://www.saccharomycessensustricto.org/cgi-](http://www.saccharomycessensustricto.org/cgi-bin/s3.cgi?data=Annotations&version=current)
29 [bin/s3.cgi?data=Annotations&version=current](http://www.saccharomycessensustricto.org/cgi-bin/s3.cgi?data=Annotations&version=current), the only comparative transcriptomic studies

1 available relating to this complex are of a single hybrid replicate grown at varying
2 temperatures, available under NCBI BioProject PRJNA505697.

3 *Saccharomyces pastorianus* (syn. *Saccharomyces carlsbergensis*)

4 This allopolyploid yeast is formed from the mating of *S. cerevisiae* and *S. eubayanus*. The
5 allopolyploid does not have a genome sequence or annotations available, but both parental
6 species do through the NCBI database. A comparative transcriptomic analysis of this
7 complex was performed by (Gibson et al., 2010), however, they used a microarray. It was
8 unclear whether the *S. pastorianus* they used was a naturally occurring allopolyploid, so the
9 authors were emailed to clarify. Their response was that their experimental procedure
10 involved an industrial strain that had been in use for decades, thus its ancestry is not entirely
11 certain. However, the author did not think it had originated from a synthetic protocol. The
12 literature also suggests that *S. pastorianus* arose through instantaneous speciation during an
13 interspecific hybridisation event (Dunn & Sherlock, 2008).

14 *Zygosaccharomyces parabailii*

15 A homoploid hybrid species of *Z. bailii sensu stricto* and an unidentified *Zygosaccharomyces*
16 species. There is an RNA-seq study of the response to lactic acid stress in *Z. parabailii*, but it
17 does not involve comparisons to its parental species (Ortiz-Merino et al., 2018).

18 *Zygosaccharomyces rouxii*

19 Yeast with allopolyploid strains. *Zygosaccharomyces rouxii* NBRC 1876 draft genome
20 sequence is available (Sato et al., 2017). RNA-seq data for this species can be found under
21 NCBI BioProject PRJNA437612. However, no comparative transcriptomic studies were
22 found through NCBI SRA.

23 *Zymoseptoria pseudotritici*

24 *Zymoseptoria pseudotritici* is a fungal homoploid hybrid grass pathogen with unknown
25 parental species (Stukenbrock et al., 2012).

26 ***S1.5 Plants***

27 *Arabidopsis suecica*

1 Natural allopolyploid of *A. thaliana* x *A. arenosa* (Jeffrey Chen et al., 2004). Comparative
2 RNA-seq data of this complex is available under NCBI BioProject PRJNA393427 with two
3 biological replicates for each member, in addition to genome annotations available through
4 NCBI SRA. The main issue was no suitable *Arabidopsis* homoploid hybrid complex.

5 *Arabidopsis thaliana* x *Arabidopsis lyrata*

6 Artificial homoploid hybrid with two RNA-seq replicates available for the hybrid and *A.*
7 *lyrata* (Zhu et al., 2017). The issue was that this study did not perform RNA-seq for *A.*
8 *thaliana*, so there was no guarantee that the conditions under which extraction was performed
9 were held constant across all species.

10 *Arachis hypogaea*

11 Peanut is an allopolyploid of *A. ipaensis* and *A. duranensis*. RNA-seq data, genomes and
12 genome annotations for the parental species and the allopolyploid are available (Bertioli et
13 al., 2019).

14 Further RNA-seq data for allopolyploid *A. hypogaea* is also available, without biological
15 replicates (Han et al., 2017).

16 The major issue was a lack on a homoploid hybrid complex from within this genus.

17 *Brassica juncea*

18 Allotetraploid hybrid of *B. rapa* and *B. nigra*. (Yang et al., 2016) did RNA-seq analysis of *B.*
19 *juncea* only, in addition to a genome annotation of *B. juncea*. The genome annotation of *B.*
20 *rapa* is available through NCBI SRA.

21 *Brassica napus*

22 An allotetraploid hybrid of *B. rapa* x *B. oleracea*. RNA-seq data with three biological
23 replicates for each member of the complex is available under NCBI BioProject
24 PRJNA449400 (Wu et al., 2018). All members have genes and genomes available through
25 NCBI SRA. This is a possible option for an allopolyploid complex.

26 *Brassica napus* x *Brassica carinata*

27 Homoploid hybrid artificially produced via hand pollination and embryo rescue. Comparative
28 RNA-seq analysis of this homoploid hybrid complex with three biological replicates for each

member, and functional gene annotations is available (Chu et al., 2014). Extensive genome annotations are available for *B. napus* through NCBI SRA. Limited gene models are available for *B. carinata* through NCBI SRA and <http://brassicadb.org>. This is a possible option for a homoploid hybrid plant complex.

Brassica napus x *Brassica rapa*

Comparative RNA-seq analysis of this homoploid hybrid complex (Zhang et al., 2015), with no biological replicates.

Camellia

Camellia reticulata can exist as an allopolyploid (Gu & Xiao, 2003), but there are no comparative transcriptomic studies on the allopolyploid complex. The only related RNA-seq data on NCBI SRA is of five different *C. reticulata* tissues (Yao et al., 2016).

A homoploid hybrid complex exists in *Camellia azalea* x *Camellia amplexicaulis*, with RNA-seq data on each member available. There are no biological replicates; the RNA libraries were formed from pooled RNA of three individuals (Zhang et al., 2018).

Capsella bursa-pastoris

Natural allopolyploid of *C. orientalis* and *C. grandiflora*. Multiple replicates of RNA-seq data are available for the allopolyploid and parental species (Kryvokhyzha et al., 2019).

Capsella rubella x *Capsella grandiflora*

Reciprocal homoploid hybrids with RNA-seq data, however, no accompanying RNA-seq data is available for the parental species (Rebernig et al., 2015).

Chrysanthemum nankingense x *Tanacetum vulgare*

Homoploid hybrid with comparative RNA-seq data available for hybrid and parental species (Wang et al., 2013; Qi et al., 2018). Qi et al. also generated allopolyploids through colchicine treatment of homoploid hybrids in addition to performing functional annotations of genes, making this a possible option for our analyses.

Coffea arabica

1 Natural allopolyploid of *C. eugenoides* and *C. canephora*. Data from a microarray-based
2 comparative analysis of this complex are available (Bardil et al., 2011), as well as an RNA-
3 seq-based comparative analysis of *C. arabica* grown under different temperatures (Combes et
4 al., 2013).

5 *Glycine*

6 No records of an RNA-seq-based study of *Glycine* homoploid hybrids. However, an
7 intraspecific comparative transcriptome analysis of homoploid *G. max* hybrids has been done
8 (Zhang et al., 2017).

9 *Glycine dolichocarpa*

10 A naturally formed recent allopolyploid of *G. syndetika* and *G. tomentella*. RNA-seq data for
11 only the allopolyploid is available (Coate et al., 2014).

12 A comparative transcriptomic study of the allopolyploid and its parents is found here, with 3
13 biological replicates per member (Ilut et al., 2012). <https://soybase.org> only has *G. max* and
14 *G. soja* genome and gene sequences. NCBI SRA has 127 gene models available for each of
15 *G. syndetika*, *G. tomentella* and *G. dolichocarpa*.

16 Transcriptomic analyses of this complex can also be found here (Coate et al., 2012).

17 *Gossypium*

18 Microarray-based analysis of natural allopolyploid cotton (*G. hirsutum*) and synthetic
19 homoploid hybrid progeny of *G. arboreum* and *G. raimondii* (Flagel et al., 2008).

20 RNA-seq based analysis of natural allopolyploid (*G. hirsutum*) and synthetic homoploid
21 hybrid progeny of *G. arboreum* and *G. raimondii* (Yoo et al., 2013). There are 3 biological
22 replicates of RNA-seq data available for each member under NCBI BioProject
23 PRJNA171342. The synthetic hybrid was formed by emasculation of the A2 parent in the
24 evening and subsequent cross-pollination the following morning. They are a strong option for
25 our homoploid and allopolyploid plant representatives.

26 *Helianthus*

The sunflower genus *Helianthus* contains several homoploid hybrid species (Ungerer et al., 2006). RNA-seq data is available (Renaut et al., 2014), in addition to a reference genome for the parental species *H. annuus* (Badouin et al., 2017).

There are some purportedly allopolyploid *Helianthus* species (Espinasse et al., 1995), but data on these are very limited, especially RNA-seq.

Mimulus

Mimulus peregrinus is a naturally formed recent allopolyploid of *M. luteus* x *M. guttatus*. Genomes and genome annotations for both parents found at <https://datadryad.org/resource/doi:10.5061/dryad.d4vr0>. Edger et al. (2017) performed a comparative RNA-seq analysis of this complex, in addition to an artificially formed *M. x robertsii* of the same parent pairing. *Mimulus x robertsii* was formed by hand pollination via diallel cross. They also did GO and KEGG. The issue was there were no biological replicates for any of the RNA-seq data.

Oryza

Comparative RNA-seq analysis of intraspecific rice hybrids and their derived allopolyploids (Xu et al., 2014).

Comparative RNA-seq analysis of a synthetic interspecific rice hybrid and its two parental species (Wu et al., 2016). However, the hybrid is a triploid produced from a diploid and tetraploid, so its genome size relative to one parent has increased and to the other, decreased, and thus it would confound the interpretation of our analyses.

Saccharum

Modern allopolyploid and homoploid sugarcane cultivars are commonly formed from pairings between domesticated *S. officinarum* and wild *S. spontaneum* (Song et al., 2016). RNA-seq data for *S. officinarum*, *S. spontaneum*, and their F₁ hybrid are available under NCBI BioProject PRJNA335885. *Saccharum intermedium*, an allopolyploid of *S. angustifolium* and *S. villosum*, has recently been described (Welker et al., 2017), but the complex lacks accompanying RNA-seq data.

Senecio

Comparative transcriptomic analyses of both an allopolyploid (*S. cambrensis* = *S. vulgaris* x *S. squalidus*) and homoploid hybrid (*S. x baxteri* = *S. vulgaris* x *S. squalidus*) of this ragwort genus are available (Hegarty et al., 2006), however, they are microarray-based. Further microarray-based transcriptomic analyses of this genus are also available (Hegarty et al., 2005; Hegarty et al., 2008, 2009).

Spartina

The cordgrass genus contains a recently formed natural allopolyploid, *S. anglica* (Ainouche et al., 2004), two interspecific hybrids *S. x neyrautii* and *S. x townsendii*, and the parental species *S. maritima* and *S. alterniflora*. Non-normalised Illumina data is available for these complexes as assembled reference transcriptomes (Boutte et al., 2016), but comparative RNA-seq data is as yet unavailable.

Microarray data for allopolyploid, homoploid hybrid and parental *Spartina* are also available (Chelaifa et al., 2010).

Tragopogon

There is RNA-seq data with biological replicates for the allopolyploid species *T. mirus* and *T. miscellus* and their diploid parents available under NCBI BioProject PRJNA210897.

A study of *T. miscellus*, its parental species, and their synthetically produced homoploid hybrid has been done, but it was not RNA-seq based, rather surveying expression of 144 duplicated gene pairs derived from the parental species (Buggs et al., 2011).

Triticeae

Synthetic allopolyploid of *Aegilops sharonensis* and *Triticum monococcum* ssp. *aegilopoides* obtained through colchicine treatment of F₁ hybrids. Gene expression was analysed through cDNA-AFLP (Kashkush et al., 2002).

Synthetic allopolyploid of *Aegilops longissimi* and *Triticum urartu*. Gene expression of the allopolyploid and its diploid parental species was analysed through RNA-seq (Wang et al., 2016). The main issue was that the allopolyploid was synthetic, as well as the lack of RNA-seq data for a homoploid hybrid complex within these genera.

Vigna reflexo-pilosa var. *glabra*

1 Allopolyploid mungbean species with hypothetical parental species, one of which is thought
2 to be extinct. RNA-seq assemblies are available for the allopolyploid (Kang et al., 2014)

3 *Zea mays*

4 Maize is purportedly of allotetraploid origin that has subsequently undergone rediploidisation
5 (Gaut et al., 2000). Thus, it would not be suitable for investigating the impact of genome
6 doubling on gene expression.

7 *Zea mays* x *Tripsacum dactyloides*

8 Interspecific hybrid of maize and gamagrass (Shavrukov & Sokolov, 2015) with
9 transcriptomic studies of each parent available (Wu et al., 2017; Gault et al., 2018), albeit
10 none for the entire complex.

11

SI.6 References

- Ainouche, M. L., Baumel, A., & Salmon, A. (2004). *Spartina anglica* C. E. Hubbard: a natural model system for analysing early evolutionary changes that affect allopolyploid genomes. *Biological Journal of the Linnean Society*, 82(4), 475-484. 10.1111/j.1095-8312.2004.00334.x
- Badouin, H., Gouzy, J., Grassa, C. J., Murat, F., Staton, S. E., Cottret, L., . . . Langlade, N. B. (2017). The sunflower genome provides insights into oil metabolism, flowering and Asterid evolution. *Nature*, 546, 148-152. 10.1038/nature22380
- Bardil, A., de Almeida, J. D., Combes, M. C., Lashermes, P., & Bertrand, B. (2011). Genomic expression dominance in the natural allopolyploid *Coffea arabica* is massively affected by growth temperature. *New Phytologist*, 192(3), 760-774. 10.1111/j.1469-8137.2011.03833.x
- Barreto, F. S., Pereira, R. J., & Burton, R. S. (2014). Hybrid dysfunction and physiological compensation in gene expression. *Molecular Biology and Evolution*, 32(3), 613-622. 10.1093/molbev/msu321
- Bertioli, D. J., Jenkins, J., Clevenger, J., Dudchenko, O., Gao, D., Seijo, G., . . . Schmutz, J. (2019). The genome sequence of segmental allotetraploid peanut *Arachis hypogaea*. *Nature Genetics*, 51(5), 877-884. 10.1038/s41588-019-0405-z
- Betto-Colliard, C., Sermier, R., Litvinchuk, S., Perrin, N., & Stöck, M. (2015). Origin and genome evolution of polyploid green toads in Central Asia: evidence from microsatellite markers. *Heredity*, 114(3), 300-308. 10.1038/hdy.2014.100
- Bogart, J. P., Bi, K., Fu, J., Noble, D. W. A., & Niedzwiecki, J. (2007). Unisexual salamanders (genus *Ambystoma*) present a new reproductive mode for eukaryotes. *Genome*, 50(2), 119-136. 10.1139/G06-152
- Boutte, J., Ferreira de Carvalho, J., Rousseau-Gueutin, M., Poulain, J., Da Silva, C., Wincker, P., . . . Salmon, A. (2016). Reference transcriptomes and detection of duplicated copies in hexaploid and allododecaploid *Spartina* species (Poaceae). *Genome Biology and Evolution*, 8(9), 3030-3044. 10.1093/gbe/evw209

1 Brasier, C. M. (2001). Rapid evolution of introduced plant pathogens via interspecific
2 hybridization: hybridization is leading to rapid evolution of Dutch elm disease and other
3 fungal plant pathogens. *BioScience*, 51(2), 123-133. 10.1641/0006-
4 3568(2001)051[0123:REOIPP]2.0.CO;2

5 Buggs, Richard J. A., Zhang, L., Miles, N., Tate, Jennifer A., Gao, L., Wei, W., . . . Soltis,
6 Douglas E. (2011). Transcriptomic shock generates evolutionary novelty in a newly formed,
7 natural allopolyploid plant. *Current Biology*, 21(7), 551-556. 10.1016/j.cub.2011.02.016

8 Campbell, M. A., Tapper, B. A., Simpson, W. R., Johnson, R. D., Mace, W., Ram, A., . . .
9 Cox, M. P. (2017). *Epichloë hybrida*, sp. nov., an emerging model system for investigating
10 fungal allopolyploidy. *Mycologia*, 109(5), 715-729. 10.1080/00275514.2017.1406174

11 Chelaifa, H., Monnier, A., & Ainouche, M. (2010). Transcriptomic changes following recent
12 natural hybridization and allopolyploidy in the salt marsh species *Spartina* × *townsendii*
13 and *Spartina anglica* (Poaceae). *New Phytologist*, 186(1), 161-174. 10.1111/j.1469-
14 8137.2010.03179.x

15 Chu, P., Liu, H., Yang, Q., Wang, Y., Yan, G., & Guan, R. (2014). An RNA-seq
16 transcriptome analysis of floral buds of an interspecific *Brassica* hybrid between *B. carinata*
17 and *B. napus*. *Plant Reproduction*, 27(4), 225-237. 10.1007/s00497-014-0253-z

18 Coate, J. E., Bar, H., & Doyle, J. J. (2014). Extensive translational regulation of gene
19 expression in an allopolyploid (*Glycine dolichocarpa*). *The Plant Cell*, 26(1), 136.
20 10.1105/tpc.113.119966

21 Coate, J. E., Powell, A. F., Owens, T. G., & Doyle, J. J. (2012). Transgressive physiological
22 and transcriptomic responses to light stress in allopolyploid *Glycine dolichocarpa*
23 (Leguminosae). *Heredity*, 110, 160. 10.1038/hdy.2012.77

24 Combes, M.-C., Dereeper, A., Severac, D., Bertrand, B., & Lashermes, P. (2013).
25 Contribution of subgenomes to the transcriptome and their intertwined regulation in the
26 allopolyploid *Coffea arabica* grown at contrasted temperatures. *New Phytologist*, 200(1),
27 251-260. 10.1111/nph.12371

1 Cox, M. P., Dong, T., Shen, G., Dalvi, Y., Scott, D. B., & Ganley, A. R. D. (2014). An
2 interspecific fungal hybrid reveals cross-kingdom rules for allopolyploid gene expression
3 patterns. *PLOS Genetics*, 10(3), e1004180. 10.1371/journal.pgen.1004180

4 Doležálková-Kaštánková, M., Pruvost, N. B. M., Plötner, J., Reyer, H.-U., Janko, K., &
5 Choleva, L. (2018). All-male hybrids of a tetrapod *Pelophylax esculentus* share its origin and
6 genetics of maintenance. *Biology of Sex Differences*, 9(1), 13. 10.1186/s13293-018-0172-z

7 Duchemin, W., Dupont, P. Y., Campbell, M. A., Ganley, A. R., & Cox, M. P. (2015).
8 HyLiTE: accurate and flexible analysis of gene expression in hybrid and allopolyploid
9 species. *BMC Bioinformatics*, 16(8). 10.1186/s12859-014-0433-8

10 Dufresne, F. (2011). The history of the *Daphnia pulex* complex: asexuality, hybridization,
11 and polyploidy. In C. Held, S. Koenemann, & C. Schubart (Eds.), *Phylogeography and*
12 *population genetics in crustacea* (pp. 217-232): CRC Press.

13 Dunn, B., & Sherlock, G. (2008). Reconstruction of the genome origins and evolution of the
14 hybrid lager yeast *Saccharomyces pastorianus*. *Genome Research*, 18(10), 1610-1623.
15 10.1101/gr.076075.108

16 Edger, P. P., Smith, R., McKain, M. R., Cooley, A. M., Vallejo-Marin, M., Yuan, Y., . . .
17 Puzey, J. R. (2017). Subgenome dominance in an interspecific hybrid, synthetic
18 allopolyploid, and a 140-year-old naturally established neo-allopolyploid monkeyflower. *The*
19 *Plant Cell*, 29(9), 2150. 10.1105/tpc.17.00010

20 Espinasse, A., Foueillassar, J., & Kimber, G. (1995). Cytogenetical analysis of hybrids
21 between sunflower and four wild relatives. *Euphytica*, 82(1), 65-72. 10.1007/BF00028710

22 Flagel, L., Udall, J., Nettleton, D., & Wendel, J. (2008). Duplicate gene expression in
23 allopolyploid *Gossypium* reveals two temporally distinct phases of expression evolution.
24 *BMC Biology*, 6(1), 16. 10.1186/1741-7007-6-16

25 Fox, D. T., Gall, J. G., & Spradling, A. C. (2010). Error-prone polyploid mitosis during
26 normal *Drosophila* development. *Genes & Development*, 24(20), 2294-2302.
27 10.1101/gad.1952710

1 Gao, Y., Zhang, H., Gao, Q., Wang, L., Zhang, F., Siva, V. S., . . . Zhang, S. (2013).
2 Transcriptome analysis of artificial hybrid pufferfish *Jiyan-1* and its parental species:
3 implications for pufferfish heterosis. *PLOS One*, 8(3), e58453. 10.1371/journal.pone.0058453

4 Gault, C. M., Kremling, K. A., & Buckler, E. S. (2018). De novo transcriptome assemblies
5 reveal parallel gene evolution with maize after ancient polyploidy. *The Plant Genome*, 11(3),
6 180012. 10.3835/plantgenome2018.02.0012

7 Gaut, B. S., d'Ennequin, M. L. T., Peek, A. S., & Sawkins, M. C. (2000). Maize as a model
8 for the evolution of plant nuclear genomes. *Proceedings of the National Academy of Sciences*
9 *of the United States of America*, 97(13), 7008-7015. 10.1073/pnas.97.13.7008

10 Genner, M. J., & Turner, G. F. (2011). Ancient hybridization and phenotypic novelty within
11 Lake Malawi's cichlid fish radiation. *Molecular Biology and Evolution*, 29(1), 195-206.
12 10.1093/molbev/msr183

13 Gibson, B. R., Graham, N. S., Boulton, C. A., Box, W. G., Lawrence, S. J., Linforth, R. S. T.,
14 . . . Smart, K. A. (2010). Differential yeast gene transcription during brewery propagation.
15 *Journal of the American Society of Brewing Chemists*, 68(1), 21-29. 10.1094/ASBCJ-2009-
16 1123-01

17 Goulielmos, G. N., & Alahiotis, S. N. (1989). Interspecific hybridization between *Drosophila*
18 species group *melanogaster* sibling species: isozymic patterns and reproductive relationships.
19 *Genome*, 32(1), 146-154. 10.1139/g89-422

20 Gu, Z., & Xiao, H. (2003). Physical mapping of the 18S-26S rDNA by fluorescent in situ
21 hybridization (FISH) in *Camellia reticulata* polyploid complex (Theaceae). *Plant Science*,
22 164(2), 279-285. 10.1016/S0168-9452(02)00410-7

23 Han, S., Liu, H., Yan, M., Qi, F., Wang, Y., Sun, Z., . . . He, G. (2017). Differential gene
24 expression in leaf tissues between mutant and wild-type genotypes response to late leaf spot
25 in peanut (*Arachis hypogaea* L.). *PLOS One*, 12(8), e0183428.
26 10.1371/journal.pone.0183428

27 He, W., Xie, L., Li, T., Liu, S., Xiao, J., Hu, J., . . . Liu, Y. (2013). The formation of diploid
28 and triploid hybrids of female grass carp \times male blunt snout bream and their 5S rDNA
29 analysis. *BMC Genetics*, 14, 110. 10.1186/1471-2156-14-110

1 Hegarty, M. J., Barker, G. L., Brennan, A. C., Edwards, K. J., Abbott, R. J., & Hiscock, S. J.
2 (2008). Changes to gene expression associated with hybrid speciation in plants: further
3 insights from transcriptomic studies in *Senecio*. *Philosophical Transactions of the Royal*
4 *Society B: Biological Sciences*, 363(1506), 3055-3069. 10.1098/rstb.2008.0080

5 Hegarty, M. J., Barker, G. L., Brennan, A. C., Edwards, K. J., Abbott, R. J., & Hiscock, S. J.
6 (2009). Extreme changes to gene expression associated with homoploid hybrid speciation.
7 *Molecular Ecology*, 18(5), 877-889. 10.1111/j.1365-294X.2008.04054.x

8 Hegarty, M. J., Barker, G. L., Wilson, I. D., Abbott, R. J., Edwards, K. J., & Hiscock, S. J.
9 (2006). Transcriptome shock after interspecific hybridization in *Senecio* is ameliorated by
10 genome duplication. *Current Biology*, 16(16), 1652-1659. 10.1016/j.cub.2006.06.071

11 Hegarty, M. J., Jones, J. M., Wilson, I. D., Barker, G. L., Coghill, J. A., Sanchez-Baracaldo,
12 P., . . . Hiscock, S. J. (2005). Development of anonymous cDNA microarrays to study
13 changes to the *Senecio* floral transcriptome during hybrid speciation. *Molecular Ecology*,
14 14(8), 2493-2510. 10.1111/j.1365-294x.2005.02608.x

15 Hu, Y., Liang, Y., Zhang, M., Tan, F., Zhong, S., Li, X., . . . Luo, P. (2018). Comparative
16 transcriptome profiling of *Blumeria graminis* f. sp. *tritici* during compatible and incompatible
17 interactions with sister wheat lines carrying and lacking *Pm40*. *PLOS One*, 13(7), e0198891.
18 10.1371/journal.pone.0198891

19 Ilut, D. C., Coate, J. E., Luciano, A. K., Owens, T. G., May, G. D., Farmer, A., & Doyle, J. J.
20 (2012). A comparative transcriptomic study of an allotetraploid and its diploid progenitors
21 illustrates the unique advantages and challenges of RNA-seq in plant species. *American*
22 *Journal of Botany*, 99(2), 383-396. 10.3732/ajb.1100312

23 Inácio, A., Pinho, J., Pereira, P. M., Comai, L., & Coelho, M. M. (2012). Global analysis of
24 the small RNA transcriptome in different ploidies and genomic combinations of a vertebrate
25 complex – the *Squalius alburnoides*. *PLOS One*, 7(7), e41158. 10.1371/journal.pone.0041158

26 Jeffrey Chen, Z., Wang, J., Tian, L., Lee, H.-S., Wang, J. J., Chen, M., . . . Osborn, T. C.
27 (2004). The development of an *Arabidopsis* model system for genome-wide analysis of
28 polyploidy effects. *Biological Journal of the Linnean Society*, 82(4), 689-700.
29 10.1111/j.1095-8312.2004.00351.x

- 1 Juchno, D., & Boroń, A. (2018). Histological evidence that diploid hybrids of *Cobitis taenia*
2 and *C. elongatoides* (Teleostei, Cobitidae) develop into fertile females and sterile males.
3 *Hydrobiologia*, 814(1), 147-159. 10.1007/s10750-018-3530-2
- 4 Juchno, D., Boroń, A., & Gołaszewski, J. (2007). Comparative morphology and histology of
5 the ovaries of the spined loach *Cobitis taenia* L. and natural allopolyploids of *Cobitis*
6 (Cobitidae). *Journal of Fish Biology*, 70(5), 1392-1411. 10.1111/j.1095-8649.2007.01419.x
- 7 Kang, Y. J., Kim, S. K., Kim, M. Y., Lestari, P., Kim, K. H., Ha, B.-K., . . . Lee, S.-H.
8 (2014). Genome sequence of mungbean and insights into evolution within *Vigna* species.
9 *Nature Communications*, 5, 5443. 10.1038/ncomms6443
- 10 Kashkush, K., Feldman, M., & Levy, A. A. (2002). Gene loss, silencing and activation in a
11 newly synthesized wheat allotetraploid. *Genetics*, 160(4), 1651-1659.
- 12 Kelleher, E. S., Edelman, N. B., & Barbash, D. A. (2012). *Drosophila* interspecific hybrids
13 phenocopy piRNA-pathway mutants. *PLOS Biology*, 10(11), e1001428.
14 10.1371/journal.pbio.1001428
- 15 Kryvokhyzha, D., Milesi, P., Duan, T., Orsucci, M., Wright, S. I., Glémin, S., & Lascoux, M.
16 (2019). Towards the new normal: transcriptomic convergence and genomic legacy of the two
17 subgenomes of an allopolyploid weed (*Capsella bursa-pastoris*). *PLOS Genetics*, 15(5),
18 e1008131. 10.1371/journal.pgen.1008131
- 19 Landry, C. R., Wittkopp, P. J., Taubes, C. H., Ranz, J. M., Clark, A. G., & Hartl, D. L.
20 (2005). Compensatory cis-trans evolution and the dysregulation of gene expression in
21 interspecific hybrids of *Drosophila*. *Genetics*, 171(4), 1813-1822.
22 10.1534/genetics.105.047449
- 23 Li, C.-Y., Li, J.-T., Kuang, Y.-Y., Xu, R., Zhao, Z.-X., Hou, G.-Y., . . . Sun, X.-W. (2014).
24 The transcriptomes of the crucian carp complex (*Carassius auratus*) provide insights into the
25 distinction between unisexual triploids and sexual diploids. *International Journal of*
26 *Molecular Sciences*, 15(6), 9386-9406. 10.3390/ijms15069386
- 27 Mallet, J. (2007). Hybrid speciation. *Nature*, 446(7133), 279-283. 10.1038/nature05706

- 1 Matos, I., Machado, M. P., Scharl, M., & Coelho, M. M. (2015). Gene expression dosage
2 regulation in an allopolyploid fish. *PLOS One*, 10(3), e0116309.
3 10.1371/journal.pone.0116309
- 4 Matos, I., Machado, M. P., Scharl, M., & Coelho, M. M. (2019). Allele-specific expression
5 variation at different ploidy levels in *Squalius alburnoides*. *Scientific Reports*, 9(1), 3688.
6 10.1038/s41598-019-40210-8
- 7 Mavárez, J., & Linares, M. (2008). Homoploid hybrid speciation in animals. *Molecular*
8 *Ecology*, 17(19), 4181-4185. 10.1111/j.1365-294X.2008.03898.x
- 9 McElroy, K. E., Denton, R. D., Sharbrough, J., Bankers, L., Neiman, M., & Lisle Gibbs, H.
10 (2017). Genome expression balance in a triploid trihybrid vertebrate. *Genome Biology and*
11 *Evolution*, 9(4), 968-980. 10.1093/gbe/evx059
- 12 Menardo, F., Praz, C. R., Wyder, S., Ben-David, R., Bourras, S., Matsumae, H., . . . Keller,
13 B. (2016). Hybridization of powdery mildew strains gives rise to pathogens on novel
14 agricultural crop species. *Nature Genetics*, 48, 201-205. 10.1038/ng.3485
- 15 Mizgireuv, I. V., Majorova, I. G., Gorodinskaya, V. M., Khudoley, V. V., & Revskoy, S. Y.
16 (2004). Carcinogenic effect of N-nitrosodimethylamine on diploid and triploid zebrafish
17 (*Danio rerio*). *Toxicologic Pathology*, 32(5), 514-518. 10.1080/01926230490496311
- 18 Murray, D. S., Kainz, M. J., Hebberecht, L., Sales, K. R., Hindar, K., & Gage, M. J. G.
19 (2018). Comparisons of reproductive function and fatty acid fillet quality between triploid
20 and diploid farm Atlantic salmon (*Salmo salar*). *Royal Society Open Science*, 5(8), 180493.
21 10.1098/rsos.180493
- 22 Nigg, M., Laroche, J., Landry, C. R., & Bernier, L. (2015). RNAseq analysis highlights
23 specific transcriptome signatures of yeast and mycelial growth phases in the Dutch elm
24 disease fungus *Ophiostoma novo-ulmi*. *G3 (Bethesda)*, 5(11), 2487-2495.
25 10.1534/g3.115.021022
- 26 Olarte, R. A., Worthington, C. J., Horn, B. W., Moore, G. G., Singh, R., Monacell, J. T., . . .
27 Carbone, I. (2015). Enhanced diversity and aflatoxigenicity in interspecific hybrids of
28 *Aspergillus flavus* and *Aspergillus parasiticus*. *Molecular Ecology*, 24(8), 1889-1909.
29 10.1111/mec.13153

1 Ortiz-Merino, R. A., Kuanyshev, N., Byrne, K. P., Varela, J. A., Morrissey, J. P., Porro, D., .
2 . . Branduardi, P. (2018). Transcriptional response to lactic acid stress in the hybrid yeast
3 *Zygosaccharomyces parabailii*. *Applied and Environmental Microbiology*, 84(5), e02294-
4 e02310. 10.1128/AEM.02294-17

5 Pala, I., Coelho, M. M., & Scharl, M. (2008). Dosage compensation by gene copy silencing
6 in a triploid hybrid fish. *Current Biology*, 18(1), 38, 10.1016/j.cub.2008.07.096

7 Pala, I., Scharl, M., Brito, M., Malta Vacas, J., & Coelho, M. M. (2010). Gene expression
8 regulation and lineage evolution: the North and South tale of the hybrid polyploid *Squalius*
9 *alburnoides* complex. *Proceedings. Biological sciences*, 277(1699), 3519-3525.
10 10.1098/rspb.2010.1071

11 Pereira, C. S. A., Aboim, M. A., Ráb, P., & Collares-Pereira, M. J. (2014). Introgressive
12 hybridization as a promoter of genome reshuffling in natural homoploid fish hybrids
13 (Cyprinidae, Leuciscinae). *Heredity*, 112(3), 343-350. 10.1038/hdy.2013.110

14 Pfliegler, W. P., Antunovics, Z., & Sipiczki, M. (2012). Double sterility barrier between
15 *Saccharomyces* species and its breakdown in allopolyploid hybrids by chromosome loss.
16 *FEMS Yeast Research*, 12(6), 703-718. 10.1111/j.1567-1364.2012.00820.x

17 Praz, C. R., Menardo, F., Robinson, M. D., Müller, M. C., Wicker, T., Bourras, S., & Keller,
18 B. (2018). Non-parent of origin expression of numerous effector genes indicates a role of
19 gene regulation in host adaption of the hybrid triticale powdery mildew pathogen. *Frontiers*
20 *in Plant Science*, 9, 49. 10.3389/fpls.2018.00049

21 Pritchard, V. L., Knutson, V. L., Lee, M., Zieba, J., & Edmands, S. (2013). Fitness and
22 morphological outcomes of many generations of hybridization in the copepod *Tigriopus*
23 *californicus*. *Journal of Evolutionary Biology*, 26(2), 416-433. 10.1111/jeb.12060

24 Qi, X., Wang, H., Song, A., Jiang, J., Chen, S., & Chen, F. (2018). Genomic and
25 transcriptomic alterations following intergeneric hybridization and polyploidization in the
26 *Chrysanthemum nankingense*×*Tanacetum vulgare* hybrid and allopolyploid (Asteraceae).
27 *Horticulture Research*, 5, 5. 10.1038/s41438-017-0003-0

- 1 Qin, Q., Lai, Z., Cao, L., Xiao, Q., Wang, Y., & Liu, S. (2016). Rapid genomic changes in
2 allopolyploids of *Carassius auratus red var.* (♀) × *Megalobrama amblycephala* (♂).
3 *Scientific Reports*, 6, 34417. 10.1038/srep34417
- 4 Rebernig, C. A., Lafon-Placette, C., Hatorangan, M. R., Slotte, T., & Köhler, C. (2015). Non-
5 reciprocal interspecies hybridization barriers in the *Capsella* genus are established in the
6 endosperm. *PLOS Genetics*, 11(6), e1005295. 10.1371/journal.pgen.1005295
- 7 Refstie, T., & Gjedrem, T. (1975). Hybrids between Salmonidae species. Hatchability and
8 growth rate in the freshwater period. *Aquaculture*, 6(4), 333-342. 10.1016/0044-
9 8486(75)90112-X
- 10 Renaut, S., Rowe, H. C., Ungerer, M. C., & Rieseberg, L. H. (2014). Genomics of homoploid
11 hybrid speciation: diversity and transcriptional activity of long terminal repeat
12 retrotransposons in hybrid sunflowers. *Philosophical Transactions of the Royal Society B:*
13 *Biological Sciences*, 369(1648), 20130345. 10.1098/rstb.2013.0345
- 14 Santos, M. E., Baldo, L., Gu, L., Boileau, N., Musilova, Z., & Salzburger, W. (2016).
15 Comparative transcriptomics of anal fin pigmentation patterns in cichlid fishes. *BMC*
16 *Genomics*, 17(1), 712. 10.1186/s12864-016-3046-y
- 17 Sato, A., Matsushima, K., Oshima, K., Hattori, M., & Koyama, Y. (2017). Draft genome
18 sequencing of the highly halotolerant and allopolyploid yeast *Zygosaccharomyces rouxii*
19 NBRC 1876. *Genome Announcements*, 5(7), e01610-e01616. 10.1128/genomeA.01610-16
- 20 Schedina, I. M., Hartmann, S., Groth, D., Schlupp, I., & Tiedemann, R. (2014). Comparative
21 analysis of the gonadal transcriptomes of the all-female species *Poecilia formosa* and its
22 maternal ancestor *Poecilia mexicana*. *BMC Research Notes*, 7, 249. 10.1186/1756-0500-7-
23 249
- 24 Schmid, M., Evans, B. J., & Bogart, J. P. (2015). Polyploidy in Amphibia. *Cytogenetic and*
25 *Genome Research*, 145(3-4), 315-330. 10.1159/000431388
- 26 Schraiber, J. G., Mostovoy, Y., Hsu, T. Y., & Brem, R. B. (2013). Inferring evolutionary
27 histories of pathway regulation from transcriptional profiling data. *PLOS Computational*
28 *Biology*, 9(10), e1003255. 10.1371/journal.pcbi.1003255

1 Selz, O. M., Thommen, R., Maan, M. E., & Seehausen, O. (2014). Behavioural isolation may
2 facilitate homoploid hybrid speciation in cichlid fish. *Journal of Evolutionary Biology*, 27(2),
3 275-289. 10.1111/jeb.12287

4 Session, A. M., Uno, Y., Kwon, T., Chapman, J. A., Toyoda, A., Takahashi, S., . . . Rokhsar,
5 D. S. (2016). Genome evolution in the allotetraploid frog *Xenopus laevis*. *Nature*, 538, 336.
6 10.1038/nature19840

7 Shavrukov, Y., & Sokolov, V. (2015). Maize-gamagrass interspecific hybrid, *Zea mays* x
8 *Tripsacum dactyloides*, shows better salinity tolerance and higher Na⁺ exclusion than maize
9 and sorghum. *International Journal of Latest Research in Science and Technology*, 4(1), 128-
10 133.

11 Song, J., Yang, X., Resende, M. F. R., Neves, L. G., Todd, J., Zhang, J., . . . Wang, J. (2016).
12 Natural allelic variations in highly polyploidy *Saccharum* complex. *Frontiers in Plant*
13 *Science*, 7, 804. 10.3389/fpls.2016.00804

14 Spiewak, J. E., Bain, E. J., Liu, J., Kou, K., Sturiale, S. L., Patterson, L. B., . . . Parichy, D.
15 M. (2018). Evolution of Endothelin signaling and diversification of adult pigment pattern in
16 *Danio* fishes. *PLOS Genetics*, 14(9), e1007538. 10.1371/journal.pgen.1007538

17 Stelkens, R. B., Schmid, C., Selz, O., & Seehausen, O. (2009). Phenotypic novelty in
18 experimental hybrids is predicted by the genetic distance between species of cichlid fish.
19 *BMC Evolutionary Biology*, 9, 283. 10.1186/1471-2148-9-283

20 Stemshorn, K. C., Reed, F. A., Nolte, A. W., & Tautz, D. (2011). Rapid formation of distinct
21 hybrid lineages after secondary contact of two fish species (*Cottus* sp.). *Molecular Ecology*,
22 20(7), 1475-1491. 10.1111/j.1365-294X.2010.04997.x

23 Stöck, M., Steinlein, C., Lamatsch, D. K., Scharl, M., & Schmid, M. (2005). Multiple origins
24 of tetraploid taxa in the Eurasian *Bufo viridis* subgroup. *Genetica*, 124(2), 255-272.
25 10.1007/s10709-005-3085-9

26 Stukenbrock, E. H., Christiansen, F. B., Hansen, T. T., Dutheil, J. Y., & Schierup, M. H.
27 (2012). Fusion of two divergent fungal individuals led to the recent emergence of a unique
28 widespread pathogen species. *Proceedings of the National Academy of Sciences of the United*
29 *States of America*, 109(27), 10954-10959. 10.1073/pnas.1201403109

- 1 Swain Lenz, D., Riles, L., & Fay, J. C. (2014). Heterochronic meiotic misexpression in an
2 interspecific yeast hybrid. *Molecular Biology and Evolution*, 31(6), 1333-1342.
3 10.1093/molbev/msu098
- 4 Tronchoni, J., García-Ríos, E., Guillamón, J. M., Querol, A., & Pérez-Torrado, R. (2017).
5 Transcriptomic analysis of *Saccharomyces cerevisiae* x *Saccharomyces kudriavzevii* hybrids
6 during low temperature winemaking. *F1000Research*, 6, 679.
7 10.12688/f1000research.11550.3
- 8 Ungerer, M. C., Strakosh, S. C., & Zhen, Y. (2006). Genome expansion in three hybrid
9 sunflower species is associated with retrotransposon proliferation. *Current Biology*, 16(20),
10 R872-R873. 10.1016/j.cub.2006.09.020
- 11 Vera, L. M., Metochis, C., Taylor, J. F., Clarkson, M., Skjærven, K. H., Migaud, H., &
12 Tocher, D. R. (2017). Early nutritional programming affects liver transcriptome in diploid
13 and triploid Atlantic salmon, *Salmo salar*. *BMC Genomics*, 18(1), 886. 10.1186/s12864-017-
14 4264-7
- 15 Wang, H., Jiang, J., Chen, S., Fang, W., Guan, Z., Liao, Y., & Chen, F. (2013). Rapid
16 genomic and transcriptomic alterations induced by wide hybridization: *Chrysanthemum*
17 *nankingense* × *Tanacetum vulgare* and *C. crassum* × *Crossostephium chinense* (Asteraceae).
18 *BMC Genomics*, 14, 902. 10.1186/1471-2164-14-902
- 19 Wang, J., Ye, L. H., Liu, Q. Z., Peng, L. Y., Liu, W., Yi, X. G., . . . Liu, S. J. (2015). Rapid
20 genomic DNA changes in allotetraploid fish hybrids. *Heredity*, 114(6), 601-609.
21 10.1038/hdy.2015.3
- 22 Wang, S., Ye, X., Wang, Y., Chen, Y., Lin, B., Yi, Z., . . . Liu, S. (2017). A new type of
23 homodiploid fish derived from the interspecific hybridization of female common carp × male
24 blunt snout bream. *Scientific Reports*, 7(1), 4189. 10.1038/s41598-017-04582-z
- 25 Wang, X., Zhang, H., Li, Y., Zhang, Z., Li, L., & Liu, B. (2016). Transcriptome asymmetry
26 in synthetic and natural allotetraploid wheats, revealed by RNA-sequencing. *New*
27 *Phytologist*, 209(3), 1264-1277. 10.1111/nph.13678
- 28 Welker, C. A. D., Souza-Chies, T. T., Peichoto, M. C., Oliveira, R. P., Carvalho, L. C.,
29 Muccillo, V., B. S., . . . Kaltchuk-Santos, E. (2017). A new allopolyploid species of

1 *Saccharum* (Poaceae - Andropogoneae) from South America, with notes on its cytogenetics.
2 *Systematic Biology*, 42(3), 507-515. 10.1600/036364417X696005

3 Withler, R. E., Beacham, T. D., Solar, I. I., & Donaldson, E. M. (1995). Freshwater growth,
4 smolting, and marine survival and growth of diploid and triploid coho salmon (*Oncorhynchus*
5 *kisutch*). *Aquaculture*, 136(1), 91-107. 10.1016/0044-8486(95)01036-X

6 Wolinska, J., Bittner, K., Ebert, D., & Spaak, P. (2006). The coexistence of hybrid and
7 parental *Daphnia*: the role of parasites. *Proceedings of the Royal Society B: Biological*
8 *Sciences*, 273(1596), 1977-1983. 10.1098/rspb.2006.3523

9 Wu, J., Lin, L., Xu, M., Chen, P., Liu, D., Sun, Q., . . . Wang, Y. (2018). Homoeolog
10 expression bias and expression level dominance in resynthesized allopolyploid *Brassica*
11 *napus*. *BMC Genomics*, 19(1), 586. 10.1186/s12864-018-4966-5

12 Wu, L., Li, M., Tian, L., Wang, S., Wu, L., Ku, L., . . . Chen, Y. (2017). Global transcriptome
13 analysis of the maize (*Zea mays* L.) inbred line 08LF during leaf senescence initiated by
14 pollination-prevention. *PLOS One*, 12(10), e0185838. 10.1371/journal.pone.0185838

15 Wu, Y., Sun, Y., Wang, X., Lin, X., Sun, S., Shen, K., . . . Liu, B. (2016). Transcriptome
16 shock in an interspecific F1 triploid hybrid of *Oryza* revealed by RNA sequencing. *Journal of*
17 *Integrative Plant Biology*, 58(2), 150-164. 10.1111/jipb.12357

18 Xu, C., Bai, Y., Lin, X., Zhao, N., Hu, L., Gong, Z., . . . Liu, B. (2014). Genome-wide
19 disruption of gene expression in allopolyploids but not hybrids of rice subspecies. *Molecular*
20 *Biology and Evolution*, 31(5), 1066-1076. 10.1093/molbev/msu085

21 Yang, J., Liu, D., Wang, X., Ji, C., Cheng, F., Liu, B., . . . Zhang, M. (2016). The genome
22 sequence of allopolyploid *Brassica juncea* and analysis of differential homoeolog gene
23 expression influencing selection. *Nature Genetics*, 48, 1225-1232. 10.1038/ng.3657

24 Yao, Q.-Y., Huang, H., Tong, Y., Xia, E.-H., & Gao, L.-Z. (2016). Transcriptome analysis
25 identifies candidate genes related to triacylglycerol and pigment biosynthesis and
26 photoperiodic flowering in the ornamental and oil-producing plant, *Camellia reticulata*
27 (Theaceae). *Frontiers in Plant Science*, 7, 163. 10.3389/fpls.2016.00163

- 1 Yoo, M. J., Szadkowski, E., & Wendel, J. F. (2013). Homeolog expression bias and
2 expression level dominance in allopolyploid cotton. *Heredity*, *110*, 171-180.
3 10.1038/hdy.2012.94
- 4 Zhang, C., Lin, C., Fu, F., Zhong, X., Peng, B., Yan, H., . . . Zhao, L. (2017). Comparative
5 transcriptome analysis of flower heterosis in two soybean F1 hybrids by RNA-seq. *PLOS*
6 *One*, *12*(7), e0181061. 10.1371/journal.pone.0181061
- 7 Zhang, J., Li, G., Li, H., Pu, X., Jiang, J., Chai, L., . . . Jiang, L. (2015). Transcriptome
8 analysis of interspecific hybrid between *Brassica napus* and *B. rapa* reveals heterosis for oil
9 rape improvement. *International Journal of Genomics*, *2015*, 11. 10.1155/2015/230985
- 10 Zhang, M., Liu, X.-K., Fan, W., Yan, D.-F., Zhong, N.-S., Gao, J.-Y., & Zhang, W.-J. (2018).
11 Transcriptome analysis reveals hybridization-induced genome shock in an interspecific F1
12 hybrid from *Camellia*. *Genome*, *61*(7), 477-485. 10.1139/gen-2017-0105
- 13 Zhivotovsky, L., G. Tochilina, T., Shaikhaev, E., P. Pogodin, V., V. Malinina, T., & J.
14 Gharrett, A. (2016). Hybrids between chum *Oncorhynchus keta* and pink *Oncorhynchus*
15 *gorbuscha* salmon: age, growth and morphology and effects on salmon production. *Journal*
16 *of Fish Biology*, *89*(4) 10.1111/jfb.13118
- 17 Zhu, W., Hu, B., Becker, C., Doğan, E. S., Berendzen, K. W., Weigel, D., & Liu, C. (2017).
18 Altered chromatin compaction and histone methylation drive non-additive gene expression in
19 an interspecific *Arabidopsis* hybrid. *Genome Biology*, *18*(1) 10.1186/s13059-017-1281-4