

Geometric Transformations of the Plane

We find standard matrices for classic transformations of the plane such as scalings, shears, rotations and reflections.

Introduction


Digital image manipulation apps continue to increase in popularity. To manipulate a digital image, we treat every pixel of the image as a point or a vector in \mathbb{R}^2 . A transformation is applied to each pixel, and the output pixel is colored the same color as the input pixel. The figure below shows the result of a non-linear transformation.



Many familiar transformations, such as rotations, reflections and shears, are linear. Every pixel (x, y) of a digital image is treated as a vector $\begin{bmatrix} x \\ y \end{bmatrix}$. To perform a linear transformation, we multiply the vector by a matrix. The figure below shows the result of a linear transformation applied to the photo of a building. Linear transformations keep the origin fixed, and map lines to lines. (See Practice Problem ??) ([reference](#))

Learning outcomes:

Author(s): Anna Davis and Paul Zachlin




Special Transformations of the Plane

We now consider several basic linear transformations and the standard matrices associated with them. The key concept is that if we want to understand what a linear transformation does, it is enough to understand what it does to basis vectors, such as standard basis vectors \mathbf{i} and \mathbf{j} . Caution: we can get into trouble if we try to construct a standard matrix for a non-linear transformation by tracking images of \mathbf{i} and \mathbf{j} , as you will see in one of the Practice Problems!

Horizontal and Vertical Scaling

Exploration Problem 1. *Let us attempt to find the standard matrix M of a transformation T that stretches the image in Figure ?? vertically by a factor of 2. Assuming that this transformation is linear, we need to find what the transformation does to the standard basis vectors. Once we have a candidate for the standard matrix, we will test it to make sure it accomplishes the stretch.*





Consider what this transformation does to the standard unit vectors (see Figure ??). We observe that $T(\mathbf{i}) = \mathbf{i}$ and $T(\mathbf{j}) = 2\mathbf{j}$. This allows us to construct a candidate for the standard matrix M , by making the images of \mathbf{i} and \mathbf{j} the columns of M . Thus,

$$M = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$$

We can now check to see what this matrix does to an arbitrary point (a, b) . Treating this point as a vector $\begin{bmatrix} a \\ b \end{bmatrix}$, we compute

$$M \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} a \\ 2b \end{bmatrix}$$

Thus, this transformation takes point (a, b) to point $(a, 2b)$. So, the proposed transformation doubles all y -coordinates resulting in a vertical stretch by a factor of 2.

Generalization 1. A vertical stretch (or compression) leaves \mathbf{i} unchanged, and scales the vector \mathbf{j} while preserving its vertical direction. Thus, a vertical stretch (or compression) maps \mathbf{i} to \mathbf{i} , and maps \mathbf{j} to $k\mathbf{j}$ for some positive number k . Similarly, a horizontal stretch (or compression) maps \mathbf{i} to $k\mathbf{i}$, and maps \mathbf{j} to \mathbf{j} .

Formula 1 (Horizontal and Vertical Scaling). A linear transformation that scales objects in the plane vertically by a factor of k is induced by

$$M_v = \begin{bmatrix} 1 & 0 \\ 0 & k \end{bmatrix} \quad (1)$$

A linear transformation that scales objects in the plane horizontally by a factor of k is induced by


$$M_h = \begin{bmatrix} k & 0 \\ 0 & 1 \end{bmatrix} \quad (2)$$

where $k > 0$.

If we were to allow k to be zero, what would the resulting transformations accomplish? In what way would the resulting matrices be fundamentally different from matrices M_v and M_h ? (See Practice Problem ??) What would happen if k were allowed to be negative?

Horizontal and Vertical Shears

A *horizontal shear* is a transformation that takes an arbitrary point (a, b) and maps it to the point $(a + kb, b)$. The effect of this transformation is that all points along a fixed horizontal line slide to the left or to the right by a fixed amount. Note that the higher the point (a, b) is above the x -axis, the greater is the magnitude of kb , resulting in a greater amount of horizontal slide.



Adding a scalar multiple of the y component to the x component can be accomplished by a matrix transformation. Observe that

$$\begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} a + kb \\ b \end{bmatrix}$$

A *vertical shear* is a transformation that takes an arbitrary point (a, b) and maps it to the point $(a, b + ka)$. This too, is a matrix transformation.

$$\begin{bmatrix} 1 & 0 \\ k & 1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} a \\ b + ka \end{bmatrix}$$


Formula 2 (Horizontal and Vertical Shears). *A linear transformation that shears the plane horizontally is induced by*

$$M_{hs} = \begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix} \quad (3)$$


A linear transformation that shears the plane vertically is induced by

$$M_{vs} = \begin{bmatrix} 1 & 0 \\ k & 1 \end{bmatrix} \quad (4)$$

Example 1. Find the standard matrix M_{vs} of a linear transformation T that shears the image of a seagull as shown in the figure below.



Explanation. Consider what this transformation does to the standard unit vectors.



The tip of the vector \mathbf{i} slides up a vertical line and its x -component remains the same. Vector \mathbf{j} stays fixed. We observe that $T(\mathbf{i}) = \begin{bmatrix} 1 \\ \tan 30^\circ \end{bmatrix} = \begin{bmatrix} 1 \\ \sqrt{3}/3 \end{bmatrix}$ and $T(\mathbf{j}) = \mathbf{j}$. This allows us to construct M_{vs} , by making the images of \mathbf{i} and \mathbf{j} the columns of M_{vs} . Thus,

$$M_{vs} = \begin{bmatrix} 1 & 0 \\ \sqrt{3}/3 & 1 \end{bmatrix}$$


Rotations about the Origin

Recall that to prove that a transformation T is linear, we have to show that for all vectors \mathbf{u} and \mathbf{v} and scalars k_1 and k_2 we have

$$T(k_1\mathbf{u} + k_2\mathbf{v}) = k_1T(\mathbf{u}) + k_2T(\mathbf{v}) \quad (5)$$

We are used to going through this verification algebraically. In some situations, however, it is instructive to think about this property geometrically. Consider a transformation R_θ that rotates every point in the plane counter-clockwise through angle θ about the origin. Is R_θ a linear transformation?


Figure ?? illustrates the left-hand side of Equation ?. Scalar multiples of \mathbf{u} and \mathbf{v} are added in the domain, then the sum is rotated through angle θ by R_θ .



The above figure illustrates the right-hand side of Equation ?. First, vectors \mathbf{u} and \mathbf{v} are rotated through angle θ , then their images are scaled and added together. Because the diagonal of a parallelogram rotates with the parallelogram, it is clear that

$$R_\theta(k_1\mathbf{u} + k_2\mathbf{v}) = k_1R_\theta(\mathbf{u}) + k_2R_\theta(\mathbf{v})$$


So, intuition tells us that R_θ is linear.




Before we consider the general standard matrix of R_θ , let's take a look at a specific example.

Example 2. Find the standard matrix M_θ of a linear transformation R_θ that rotates the image by 45° counterclockwise.

Geometric Transformations of the Plane




Explanation. Consider the action of R_θ on the standard unit vectors.



We observe that $R_\theta(\mathbf{i}) = \begin{bmatrix} \cos 45^\circ \\ \sin 45^\circ \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} \end{bmatrix}$ and $R_\theta(\mathbf{j}) = \begin{bmatrix} -\sin 45^\circ \\ \cos 45^\circ \end{bmatrix} = \begin{bmatrix} -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} \end{bmatrix}$. This allows us to construct the matrix M_θ , by making the images of \mathbf{i} and \mathbf{j} the columns of M_θ . Thus,

$$M_\theta = \begin{bmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \end{bmatrix}$$

Generalization 2. We find the standard rotation matrix by determining the images of vectors \mathbf{i} and \mathbf{j} .




Formula 3 (Counterclockwise Rotation). *A linear transformation that rotates the plane counterclockwise through angle θ about the origin is induced by*

$$M_\theta = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \quad (6)$$

Reflections about Lines of the Form $y = mx$

When a point is reflected about a line, its image is located on the opposite side of the line and the same distance away from the line as the original point.

For example, Figure ?? shows the reflection of point A about line l . Note that the reflection lies on a line through A perpendicular to l .



Arguing in a manner similar to our discussion of linearity of rotations, we can see that reflections are also linear. Our task is to find the matrix of a reflection of the plane about an arbitrary line through the origin.

We will start by finding reflections about the axes. You can easily do this on your own by finding the images of vectors \mathbf{i} and \mathbf{j} .

Geometric Transformations of the Plane

We will start with the reflection about the x -axis.

$$\mathbf{i} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{maps to} \quad \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\mathbf{j} = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad \text{maps to} \quad \begin{bmatrix} 0 \\ -1 \end{bmatrix}$$

So, the standard matrix that induces the reflection about the x -axis is

$$\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

Next, we will consider the reflection about the y -axis.

$$\mathbf{i} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{maps to} \quad \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$


$$\mathbf{j} = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \quad \text{maps to} \quad \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Thus, the standard matrix that induces the reflection about the y -axis is

$$\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$

Now we will turn our attention to transformations that reflect the plane about the line $y = mx$. We will assume that $m \neq 0$.

Consider the vector \mathbf{i} and its reflection. (See Figure ??)



Observe that the head of the image vector, $T(\mathbf{i})$, will lie on the line that passes through $(1, 0)$ and is perpendicular to the line $y = mx$. The equation of this line is given by

$$y = -\frac{1}{m}x + \frac{1}{m} \quad (7)$$

The head of $T(\mathbf{i})$ will also lie on the circle with equation

$$x^2 + y^2 = 1$$

To find the image of \mathbf{i} we need to determine where the line $y = -\frac{1}{m}x + \frac{1}{m}$ intersects the circle. Substitution gives us

$$x^2 + \left(-\frac{1}{m}x + \frac{1}{m}\right)^2 = 1$$

After a little algebra we get

$$\left(1 + \frac{1}{m^2}\right)x^2 - \left(\frac{2}{m^2}\right)x + \left(\frac{1}{m^2} - 1\right) = 0$$

The quadratic formula yields

$$x = 1 \quad \text{and} \quad x = \frac{1 - m^2}{m^2 + 1}$$


The solution $x = 1$ corresponds to the head of the vector \mathbf{i} . So, the x -component of $T(\mathbf{i})$ is $x = \frac{1 - m^2}{m^2 + 1}$. We find the y -component of $T(\mathbf{i})$ by substituting $x = \frac{1 - m^2}{m^2 + 1}$ into Equation ??.

$$y = -\frac{1}{m} \left(\frac{1 - m^2}{m^2 + 1}\right) + \frac{1}{m} = \frac{2m}{m^2 + 1}$$

Thus, the image of \mathbf{i} under this reflection is given by

$$T(\mathbf{i}) = \begin{bmatrix} \frac{1 - m^2}{m^2 + 1} \\ \frac{2m}{m^2 + 1} \end{bmatrix}$$

Next we need to find the image of \mathbf{j} . The head of $T(\mathbf{j})$ is located at one of the intersections of line $y = -\frac{1}{m}x + 1$ and the circle $x^2 + y^2 = 1$.



We leave it to the reader to verify that

$$T(\mathbf{j}) = \begin{bmatrix} \frac{2m}{m^2 + 1} \\ \frac{m^2 - 1}{m^2 + 1} \end{bmatrix} \quad (8)$$


The standard matrix of this reflection is then given by

$$M_{y=mx} = \begin{bmatrix} \frac{1-m^2}{m^2+1} & \frac{2m}{m^2+1} \\ \frac{2m}{m^2+1} & \frac{m^2-1}{m^2+1} \end{bmatrix} = \frac{1}{1+m^2} \begin{bmatrix} 1-m^2 & 2m \\ 2m & m^2-1 \end{bmatrix}$$

Formula 4 (Reflection about the line $y = mx$). A linear transformation that reflects the plane about the line $y = mx$ is induced by

$$M_{y=mx} = \frac{1}{1+m^2} \begin{bmatrix} 1-m^2 & 2m \\ 2m & m^2-1 \end{bmatrix} \quad (9)$$

Example 3. Find the standard matrix M of a linear transformation that reflects the image in Figure ?? about the line $y = \frac{3}{5}x$.



Explanation.

$$M = \frac{1}{1 + 9/25} \begin{bmatrix} 1 - 9/25 & 6/5 \\ 6/5 & 9/25 - 1 \end{bmatrix} = \frac{25}{34} \begin{bmatrix} 16/25 & 6/5 \\ 6/5 & -16/25 \end{bmatrix} = \begin{bmatrix} 8/17 & 15/17 \\ 15/17 & -8/17 \end{bmatrix}$$

Note that the eye of the duck in Figure ?? is located on the line $y = \frac{3}{5}x$. The reflection leaves the eye fixed in place. The eye is an example of a *fixed point*. In Practice Problem ?? [reference](#) you will be asked to show that every point along the line $y = \frac{3}{5}x$ is a fixed point.

Composition of Linear Transformations

If a linear transformation is followed by another linear transformation, the resulting transformation can be represented by a product of the two matrices that induce the individual transformations. Thus, if T_1 is induced by M_1 and T_2 is induced by M_2 , then $T = T_2 \circ T_1$ is induced by $M = M_2 M_1$.

$$\mathbb{R}^2 \xrightarrow{T_1} \mathbb{R}^2 \xrightarrow{T_2} \mathbb{R}^2$$

\curvearrowright
 T

Remember that matrix multiplication is not commutative, so the order in which the matrices are multiplied is of utmost importance.

Exploration Problem 2. *In this problem we will consider compositions of two reflections and use geometry to illustrate non-commutativity of matrix multiplication. Let*

$$T_{y=-2x} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

be a reflection about the line $y = -2x$. Let


$$T_{y=x} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

be a reflection about the line $y = x$. We will denote the standard matrices for these transformations by $M_{y=-2x}$ and $M_{y=x}$, and use geometry to demonstrate that $M_{y=x} M_{y=-2x} \neq M_{y=-2x} M_{y=x}$.

To do this, consider transformations $T_1 = T_{y=x} \circ T_{y=-2x}$ and $T_2 = T_{y=-2x} \circ T_{y=x}$. Transformation T_1 is induced by $M_{y=x} M_{y=-2x}$, and T_2 is induced by $M_{y=-2x} M_{y=x}$.

Figure ?? illustrates the action of T_1 on a single point A . First, A is reflected about the line $y = -2x$, then A is reflected about the line $y = x$.

Figure ?? also shows the action of T_2 on the same point A . The point is first reflected about the line $y = x$, followed by a reflection about the line $y = -2x$. The final images of point A under T_1 and T_2 are clearly different.



Since $T_1 \neq T_2$, we conclude that $M_{y=x}M_{y=-2x} \neq M_{y=-2x}M_{y=x}$.

Example 4. Some pairs of matrices do commute. For example, geometry makes it is easy to see that two rotation matrices commute.

Inverse of a Linear Transformation

Recall that two linear transformations are inverses of each other if their composition is the identity transformation. If a linear transformation T induced by the matrix M is invertible, then T^{-1} is induced by M^{-1} .

Geometry can help find the inverse of certain matrices. For example, we can easily see that the inverse of the rotation transformation $R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ with standard matrix M_θ has the inverse $R_{360^\circ-\theta} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ with standard matrix $M_{360^\circ-\theta}$.

Practice Problems


Problem 1 Consider matrices M_v and M_h in (??) and (??).

- If we were to allow k to be zero, what would the resulting transformations accomplish?
- In what way would the resulting matrices be fundamentally different from matrices M_v and M_h ($k \neq 0$)?
- Do M_v and M_h ($k \neq 0$) have inverses? What about M_v and M_h ($k = 0$)?
- What would happen if we allowed k to be negative?

Problem 2 Find the standard matrix M of a linear transformation that would double the length of a photo horizontally, and triple the height of the photo.


$$M = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$$

Problem 3 (Sheared Sheep) Find the standard matrix of the linear transformation shown in the figure.



Problem 4 In the beginning of this module we claimed that linear transformations leave the origin fixed and map lines to lines. Prove this claim.

Problem 5 Suppose a 1 by 1 photo of a chipmunk was shifted as shown in the figure.



Suppose we tried to construct a standard matrix M for this transformation by making the images of \mathbf{i} and \mathbf{j} the columns of M . We would obtain

$$M = \begin{bmatrix} 1.4 & 0.4 \\ 0.4 & 1.4 \end{bmatrix}$$

Does this matrix describe the transformation? If so, prove it. If not, explain why not.

Problem 6 A transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ that shifts all points in the plane horizontally or vertically by a fixed amount is called a *translation*. Is T a linear transformation? Prove your claim.

Problem 7 A reflection about the line $y = mx$ followed by another reflection about the same line, amounts to the identity transformation. Prove this using matrix multiplication.

Problem 8 Verify Equation (??).

Problem 9 Prove that every point along the line $y = \frac{3}{5}x$ in Example ?? is a fixed point.

Problem 10 The figure below shows a sequence of two linear transformations that accomplishes a reflection about the line $y = \frac{3}{5}x$. The first transformation is a reflection of the plane about the x -axis. The second transformation is a rotation of the plane about the origin. Find the standard matrices for the two transformations and verify that their product (in the correct order) is the reflection matrix of Example ??.

Geometric Transformations of the Plane

