


Flow Annealed Importance Sampling Bootstrap Meets Differentiable Particle Physics




Annalena Kofler^{1,2,3}, Vincent Stimper^{1,4,5}, Mikhail Mikhaseenko^{6,7}, Michael Kagan⁸, Lukas Heinrich³

¹MPI for Intelligent Systems Tübingen, ²MPG for Gravitational Physics Potsdam, ³Technical University of Munich, ⁴Isomorphic Labs, ⁵University of Cambridge, ⁶Ruhr University Bochum, ⁷ORIGINS Excellence Cluster Munich, ⁸SLAC National Accelerator Laboratory

Generation of Data in Particle Physics




Normalizing Flow




→ Generate samples x and evaluate density $q_\theta(x)$


Training Approaches



→ FAB uses HMC for Annealed Importance Sampling (AIS)
→ Differentiable target distribution $p(x)$ required for rKLD and FAB



Distribution $\Lambda_c^+ \rightarrow pK^-\pi^+$




Sampling Efficiency ϵ

$$\epsilon = \frac{1}{N} \frac{\left(\sum_i w_i \right)^2}{\sum_i w_i^2} \in [0,1]$$

with importance weight

$$w_i = \frac{p(x_i)}{q_\theta(x_i)}$$



$q_\theta(x) \neq p(x) \rightarrow \text{Var}[w_i] \text{ is large} \rightarrow \epsilon \ll 1$

Efficiency with more target evaluations

